

Surgical Periodontics: Resective Procedures

Policy Number: DCP013.15
Effective Date: May 1, 2026

[Instructions for Use](#)

Table of Contents	Page
Coverage Rationale	1
Definitions	2
Applicable Codes	3
Description of Services	4
Clinical Evidence	4
References	5
Policy History/Revision Information	6
Instructions for Use	6

Related Dental Policies
<ul style="list-style-type: none"> Coronal Splinting Surgical Periodontics: Mucogingival Procedures

Coverage Rationale

Gingivectomy/Gingivoplasty

[Gingivectomy/Gingivoplasty](#) is indicated for the following:

- Elimination of suprabony pockets, exceeding 3mm, if the pocket wall is fibrous and firm and there is an adequate zone of keratinized tissue
- Elimination of gingival enlargements/overgrowth
- Elimination of suprabony periodontal abscesses
- Exposure of soft tissue impacted teeth to aid in eruption
- To reestablish gingival contour following an episode of acute necrotizing ulcerative gingivitis
- To allow restorative access, including root surface caries

Gingivectomy/Gingivoplasty is not indicated for the following:

- When bone surgery is required for infrabony defects, or for the purpose of examining bone shape and morphology
- Situations in which the bottom of the pocket is apical to the mucogingival junction
- In areas with a shallow palatal vault or prominent external oblique ridge

Anatomical Crown Exposure

[Anatomical Crown](#) exposure is indicated in a periodontally healthy area for the following:

- To facilitate the restoration of subgingival caries
- To allow proper contour of restoration
- To allow management of a subgingivally fractured tooth

Flap Procedures

[Gingival Flap](#) and apically positioned flap procedures are indicated for the following:

- The presence of moderate to deep probing depths
- Moderate/severe gingival enlargement or extensive areas of overgrowth
- Loss of attachment
- The need for increased access to root surface and/or alveolar bone when previous non-surgical attempts have been unsuccessful
- The diagnosis of a cracked tooth, fractured root, or external root resorption when this cannot be accomplished by non-invasive methods
- To preserve keratinized tissue in conjunction with Osseous Surgery

Clinical Crown Lengthening – Hard Tissue

Clinical crown lengthening – hard tissue is indicated for the following:

- In an otherwise periodontally healthy area to allow a restorative procedure on a tooth with little to no crown exposure
- To allow preservation of the biological width for restorative procedures

Osseous Surgery

Osseous Surgery is indicated for the following:

- Patients with a diagnosis of Stage III or Stage IV periodontal disease
- When less invasive therapy (i.e., non-surgical periodontal therapy, Flap procedures) has failed to eliminate disease

Osseous Surgery is not indicated for teeth with a hopeless prognosis.

Mesial/Distal Wedge

A mesial/distal wedge procedure is indicated for the following:

- The presence of moderate to deep probing depths (greater than 5mm) on a surface adjacent to an edentulous/terminal tooth area
- The need for increased access to root surface and/or alveolar bone when previous non-surgical attempts have been unsuccessful on a surface adjacent to an edentulous/terminal tooth area
- The diagnosis of a cracked tooth, fractured root, or external root resorption on a surface adjacent to an edentulous/terminal tooth area, when this cannot be accomplished by non-invasive methods

Resective Periodontal Surgical Procedures

Resective periodontal surgical procedures are not indicated for the following:

- Individuals who have been non-compliant with non-surgical periodontal therapies
- For teeth with a hopeless prognosis
- Individuals with an unmanaged medical condition; these conditions include but are not limited to metabolic, cardiovascular, and autoimmune/inflammatory, as well as genetic conditions that affect collagen synthesis
- Individuals taking medications that negatively affects the healing response; these include but are not limited to immunosuppressive agents, corticosteroids, anticoagulants, NSAIDs, and nicotine

Definitions

Anatomical Crown: That portion of tooth normally covered by, and including, enamel. (ADA)

Gingival Flap: A section of the gingiva and/or the mucosa surgically separated from the underlying tissues to provide visibility and access to the bone and root surface. (Reddy)

Gingivectomy: The excision or removal of gingiva. (ADA)

Gingivoplasty: Surgical procedure to reshape gingiva. (ADA)

McGuire Classification of Tooth Prognosis (Levi, 2016):

- Good: Teeth with adequate periodontal support where the etiologic factors can be controlled, including systemic factors.
- Fair: No more than 25% attachment loss with Grade 1 furcation invasion which can be maintained. Plaque control and systemic factors can be maintained.
- Poor: As much as 50% bone loss with Grade II furcation invasions, poor crown: root ratio; Mobility greater than Miller Class I; systemic factors; poor patient participation in treatment.
- Questionable: Teeth with greater than 50% attachment loss; Grade II or III furcation involvements; the tooth is not easily maintained either with professional hygiene or by the patient.
- Hopeless: Inadequate attachment to support the tooth; Class III or IV furcation involvement; Miller Class III Mobility; the tooth cannot be maintained with adequate plaque control by the clinician or by the patient.

Mobility: The movement of a tooth in its socket resulting from an applied force. (AAP) Miller Index of Tooth Mobility (Harpenau 2013):

- Class 0: Normal physiologic tooth movement.
- Class I: First distinguishable signs of movement beyond normal.

- Class II: Tooth movement up to 1mm in any direction.
- Class III: Tooth can be moved more than 1mm in any direction and/or the tooth can be depressed into the socket.

Osseous Surgery: Procedures to modify bone support altered by periodontal disease, either by reshaping the alveolar process to achieve physiologic form without the removal of alveolar supporting, or by the removal of some alveolar bone, thus changing the position of the crestal bone relative to the tooth root. (See: Ostectomy; Osteoplasty) (AAP)

Quadrant: One of the four equal sections into which the dental arches can be divided; begins at the midline of the arch and extends distally to the last tooth. (ADA)

Staging Periodontitis (AAP):

Stage I

- 1-2mm clinical attachment loss (CAL)
- Radiographic bone loss (RBL) of < 15%
- No tooth loss
- Complexity
 - Maximum probing depth ≤ 4mm
 - Mostly horizontal bone loss

Stage II

- 3-4mm interdental CAL
- RBL of 15-33%
- No tooth loss
- Complexity
 - Maximum probing depth ≤ 5mm
 - Mostly horizontal bone loss

Stage III

- ≥ 5mm CAL
- RBL extends to middle third of root and beyond
- Loss of ≤ 4 teeth
- Complexity includes all of criteria for Stage II as well as:
 - Probing depths ≥ 6mm
 - Vertical bone loss ≥ 3mm
 - Class II or III furcation involvement
 - Moderate ridge defects

Stage IV

- ≥ 5mm CAL
- RBL extends to middle third of root and beyond
- Loss of ≥ 5 teeth
- Complexity includes all of criteria for Stage III as well as:
 - The need for complex rehabilitation due to:
 - Masticatory dysfunction
 - Secondary occlusal trauma (tooth obliquity ≥ 2)
 - Severe ridge defects
 - Bite collapse, drifting and/or flaring
 - < 20 remaining teeth (10 opposing pairs)

The extent and distribution for each stage is described as:

- Localized (< 30% of teeth involved); or
- Generalized; or
- Molar/incisor pattern

Applicable Codes

The following list(s) of procedure and/or diagnosis codes is provided for reference purposes only and may not be all inclusive. Listing of a code in this policy does not imply that the service described by the code is a covered or non-covered health service. Benefit coverage for health services is determined by the member specific benefit plan document and applicable laws that may require coverage for a specific service. The inclusion of a code does not imply any right to reimbursement or guarantee claim payment. Other Policies and Guidelines may apply.

CDT Code	Description
D4210	Gingivectomy or gingivoplasty – four or more contiguous teeth or tooth bounded spaces per quadrant
D4211	Gingivectomy or gingivoplasty – one to three contiguous teeth or tooth bounded spaces per quadrant
D4212	Gingivectomy or gingivoplasty to allow access for restorative procedure, per tooth
D4230	Anatomical crown exposure – four or more contiguous teeth or bounded tooth spaces per quadrant
D4231	Anatomical crown exposure one to three teeth or bounded tooth spaces per quadrant
D4240	Gingival flap procedure, including root planing – four or more contiguous teeth or tooth bounded spaces per quadrant
D4241	Gingival flap procedure, including root planing – one to three contiguous teeth or tooth bounded spaces per quadrant
D4245	Apically positioned flap
D4249	Clinical crown lengthening – hard tissue
D4260	Osseous surgery (including flap entry and closure) – four or more contiguous teeth or tooth bounded spaces per quadrant
D4261	Osseous surgery (including flap entry and closure) – one to three contiguous teeth or tooth bounded spaces per quadrant
D4274	Mesial/distal wedge procedure, single tooth (when not performed in conjunction with surgical procedures in the same anatomical area)
D4999	Unspecified periodontal procedure, by report

CDT® is a registered trademark of the American Dental Association

Description of Services

Using non-surgical periodontal therapy, many individuals can be treated and maintained without the need for surgical intervention. However, surgical procedures may be required when periodontal health cannot be achieved or maintained non-surgically, and may be performed by electrosurgery, lasers or surgical scalpels. Resective periodontal surgery procedures are long established standards of care and are indicated to eliminate pockets and recontour osseous bone. They may also be indicated when there is a need to expose or lengthen the clinical crown for the completion of restorative procedures.

Pursuant to CA AB2585: While not common in dentistry, nonpharmacological pain management strategies should be encouraged if appropriate.

Clinical Evidence

Hayakawa et al. (2012) conducted a retrospective study with the aim of investigating the outcome of surgical periodontal therapy during the period of April 2010 through March 2012 at the General Dentistry, Tokyo Dental College Suidobashi Hospital. The main focus is to compare open flap debridement and regenerative treatment modalities. Following initial periodontal therapy, 17 clinicians performed a total of 138 periodontal surgeries in 80 patients with moderate to advanced periodontitis. Open Flap Debridement was the most commonly performed surgery (74%), followed by 29 regenerative procedures, 7 cases of periodontal plastic surgery, and no cases of guided tissue regeneration. Clinical parameters (probing depth, bleeding on probing and clinical attachment levels) were reduced following initial therapy for all cases, with surgical intervention reducing them further. There was a significant gain in clinical attachment level when regenerative therapy was performed on areas with an initial probing depth greater than 8 mm. The authors concluded that while initial non-surgical therapy improves clinical parameters, open flap debridement surgery results in significantly higher gain in clinical attachment level for probing depths over 6 mm, with periodontal regeneration surgery providing higher gain in areas with probing depths exceeding 8mm.

Heitz-Mayfield et al. (2002) conducted a systemic review of the evidence of effectiveness of surgical vs. non-surgical therapy for the treatment of chronic periodontal disease. Sources included the National Library of Medicine computerized bibliographic database MEDLINE, and the Cochrane Oral Health Group (COHG) Specialist Trials Register. The primary outcome measures evaluated were gain in clinical attachment level (CAL) and reduction in probing pocket depth (PPD). Meta-analysis evaluation of these studies indicated that 12 months following treatment, surgical therapy resulted in 0.6

mm more PPD reduction than non-surgical therapy in pockets 6 mm or greater. The authors concluded that both scaling and root planing alone and scaling and root planing combined with flap procedure are effective methods for the treatment of chronic periodontitis in terms of attachment level gain and reduction in gingival inflammation. In the treatment of pocket depths greater than 6 mm, open flap debridement results in greater PPD reduction and clinical attachment gain.

Levy et al. (2002) conducted an investigational study to examine the clinical and microbiologic effects of apically repositioned flap surgery. (This study was intended to extend the findings of a previous study that evaluated the effect of apically repositioned flap surgery on clinical parameters and the composition of the subgingival microbiota at 3 months posttherapy). Eighteen patients with chronic periodontitis received initial preparation (IP) including scaling and root planing followed 3 months later by apically repositioned flap surgery at sites with pocket depth greater than 4 mm. All subjects had at least 20 teeth and at least eight sites with pockets greater than 4 mm and eight sites with attachment loss greater than 3 mm. Subjects were monitored clinically and microbiologically at baseline, 3 months after IP, and at 3-, 6-, 9-, and 12-months post-surgery. Clinical assessments of plaque accumulation, gingival redness, suppuration, bleeding on probing, pocket depth, and attachment level were made at six sites per tooth and the presence and levels of 40 subgingival groups of organisms were determined using checkerboard DNA-DNA hybridization. Significant reductions were seen in mean pocket depth, bacterial colonization and percentage of sites exhibiting gingival redness and bleeding on probing in sites that received IP only and in sites receiving IP followed by surgery. Mean attachment level increased significantly for both sets of sites, but the increase was greater at the surgically treated sites. The study indicated that there were beneficial changes in most clinical parameters accompanied by clear reductions in the post pathogenic organisms associated with periodontal disease. One of the most important aspects of this study was the further improvement at sites that received IP only, once periodontal surgery had been completed at the deeper periodontal pockets. The reduction in pocket depth by surgical means and the associated decrease in reservoirs of periodontal pathogens may be important in achieving sustained periodontal stability. Thus, periodontal surgery appears to be an important part of the armamentarium to control periodontal infections. This study supported and extended the findings of the previous study and described changes not only at sites receiving apically repositioned flap surgery, but also at sites in the same mouth that received IP only. While the major beneficial clinical and microbiologic effect was observed at 3 months after surgery, these beneficial effects were sustained for at least 1 year and conceivably longer.

Serino et al. (2001) performed a clinical trial to determine the initial outcome of non-surgical and surgical periodontal therapy in subjects with advanced disease, as well as the incidence of recurrent disease during 12 years of maintenance following active therapy. There were 64 subjects included in the trial, and all showed signs of generalized gingival inflammation, had a minimum of 12 non-molar teeth with deep pockets (6mm or greater) and with 6mm or greater alveolar bone loss. They were randomly assigned to 2 treatment groups; one surgical (SU) and one non-surgical (SRP). After therapy, all subjects were enrolled in a maintenance care program and were provided with meticulous supportive periodontal therapy and maintenance 3-4 times per year. At these maintenance appointments, sites that bled on gentle probing and had probing depths greater than or equal to 5 mm were treated with subgingival instrumentation. Comprehensive re-examinations were performed after 1, 3, 5, and 13 years of maintenance therapy. It was observed that that surgical therapy was more effective than non-surgical scaling and root planing in reducing the overall mean probing pocket depth and in eliminating deep pockets, and that more non-surgical subjects exhibited signs of advanced disease progression in the 1-3-year period following active therapy than the subjects initially treated surgically. The authors concluded that in subjects with advanced periodontal disease, surgical therapy provides better short and long-term periodontal pocket reduction and may lead to fewer subjects requiring additional adjunctive therapy.

References

- American Academy of Periodontology (AAP). Staging and Grading Periodontitis. 2018. Available at: <https://www.perio.org/wp-content/uploads/2019/08/Staging-and-Grading-Periodontitis.pdf>. Accessed March 2, 2026.
- American Dental Association Glossary of Clinical and Administrative Terms.
- Bezerra B. Newman and Carranza's Clinical Periodontology and Implantology, 14th ed. St. Louis: Elsevier c2023. Chapter 62, Pocket Reduction Therapy-Resective Approach; p. 750-755.
- Cohen, Edward S, DMD. Atlas of Cosmetic and Reconstructive Periodontal Surgery, 3rd ed. Shelton, CT: Peoples Medical Publishing House c2007. Chapter 5, Gingivectomy and Gingivoplasty; p. 39.
- Hayakawa H, Fujinami K, Ida A, et al. Clinical outcome of surgical periodontal therapy: a short-term retrospective study. Bull Tokyo Dent Coll. 2012; 53(4):189-95.
- Heitz-Mayfield LJ, Trombelli L, Heitz F, et al. A systematic review of the effect of surgical debridement vs non-surgical debridement for the treatment of chronic periodontitis. J Clin Periodontol. 2002; 29 Suppl 3:92-102.

Kerr A, Miller C, Nelson R. Little and Falace's Dental Management of the Medically Compromised Patient, 10th ed. St. Louis: Elsevier c2024. Chapter 1, Patient Evaluation, Risk Assessment, and the Diagnostic Process; p. 1-17.

Klokkevold P, Mealy B, Otomo-Corgel J. Newman and Carranza's Clinical Periodontology and Implantology. 14th ed. St. Louis: Elsevier c2023. Chapter 67, Periodontal Treatment of Medically Compromised Patients; p.784-805.

Klokkevold P, Takei H, Carranza F, et al. Newman and Carranza's Clinical Periodontology and Implantology, 14th ed. St. Louis: Elsevier c2023. Chapter 60, Gingival Surgical Techniques; p. 717-729.

Levy RM, Giannobile WV, Feres M, et al. The Effect of Apically Repositioned Flap Surgery on Clinical Parameters and the Composition of the Subgingival Microbiota: 12-Month Data. The International journal of periodontics & restorative dentistry. 2002;22(3):209-219.

Melnick P, Takei H. Newman and Carranza's Clinical Periodontology, 14th ed. St. Louis: Elsevier c2023 Chapter 66, Preparation of the Periodontium for Restorative Dentistry; p.780-83.

Newman M, Do J, Takei H, et al. Newman and Carranza's Clinical Periodontology, 14th ed. St. Louis: Elsevier c2023 Chapter 61, Periodontal Surgical Therapy; p.730-49.

Reddy, S. Essentials of Clinical Periodontology & Periodontics, 5th Edition. JP Medical Ltd c2018; glossary, p.563.

Serino G, Rosling B, Ramberg P, et al. Initial outcome and long-term effect of surgical and non-surgical treatment of advanced periodontal disease. J Clin Periodontol. 2001 Oct; 28(10):910-6.

Policy History/Revision Information

Date	Summary of Changes
05/01/2026	Supporting Information <ul style="list-style-type: none">Updated <i>References</i> section to reflect the most current informationArchived previous policy version DCP013.14

Instructions for Use

This Dental Clinical Policy provides assistance in interpreting UnitedHealthcare standard and Medicare Advantage dental plans. When deciding coverage, the member specific benefit plan document must be referenced as the terms of the member specific benefit plan may differ from the standard dental plan. In the event of a conflict, the member specific benefit plan document governs. Before using this policy, check the member specific benefit plan document and any applicable federal or state mandates. UnitedHealthcare reserves the right to modify its Policies and Guidelines as necessary. This Dental Clinical Policy is provided for informational purposes. It does not constitute medical advice.