

Cosentyx® (Secukinumab)

Policy Number: IEXD0132.04
Effective Date: June 1, 2026

[Instructions for Use](#)

Table of Contents	Page
Applicable States	1
Coverage Rationale	1
Applicable Codes	2
Background	3
Benefit Considerations	3
Clinical Evidence	3
U.S. Food and Drug Administration	4
References	4
Policy History/Revision Information	4
Instructions for Use	5

Related Policies
None

Applicable States

This Medical Benefit Drug Policy applies to Individual Exchange benefit plans in all states except for Nevada. For Nevada, refer to the [UnitedHealthcare Commercial Medical Benefit Drug Policy](#).

Coverage Rationale

[See Benefit Considerations](#)

This policy refers to Cosentyx (secukinumab) for intravenous (IV) injection for administration by a healthcare professional. Cosentyx (secukinumab) for self-administered subcutaneous injection is obtained under the pharmacy benefit.

Psoriatic Arthritis (PsA)

Cosentyx is proven and medically necessary for the treatment of psoriatic arthritis (PsA) when all of the following criteria are met:

- For **initial therapy**, all of the following:
 - Diagnosis of active psoriatic arthritis; **and**
 - **One** of the following:
 - History of failure to a 3 month trial of methotrexate at the maximally indicated dose, unless contraindicated or clinically significant adverse effects are experienced; **or**
 - Patient has been previously treated with a systemic targeted immunomodulator FDA-approved for the treatment of psoriatic arthritis as documented by claims history or submission of medical records (document drug, date, and duration of therapy) [e.g., adalimumab, Bimzelx (bimekizumab-bkzx), Cimzia (certolizumab), Enbrel (etanercept), Orencia (abatacept), Otezla (apremilast), Rinvoq (upadacitinib), Simponi (golimumab), Skyrizi (risankizumab), Taltz (ixekizumab), Tremfya (guselkumab), Xeljanz/Xeljanz XR (tofacitinib), ustekinumab]
- and**
- Cosentyx is initiated and titrated according to U.S. FDA labeled dosing for PsA; **and**
- Patient is not receiving Cosentyx in combination with a systemic targeted immunomodulator [e.g., adalimumab, Bimzelx (bimekizumab-bkzx), Cimzia (certolizumab), Enbrel (etanercept), Orencia (abatacept), Otezla (apremilast), Rinvoq (upadacitinib), Simponi (golimumab), Skyrizi (risankizumab), Taltz (ixekizumab), Tremfya (guselkumab), Xeljanz/Xeljanz XR (tofacitinib), ustekinumab] for treatment of the same indication; **and**
- Prescribed by or in consultation with **one** of the following:

- Rheumatologist; **or**
- Dermatologist
- and**
- Initial authorization will be issued for 12 months
- For **continuation of therapy**, all of the following:
 - Documentation of positive clinical response; **and**
 - Cosentyx is initiated and titrated according to U.S. FDA labeled dosing for PsA; **and**
 - Patient is not receiving Cosentyx in combination with a systemic targeted immunomodulator [e.g., adalimumab, Bimzelx (bimekizumab-bkzx), Cimzia (certolizumab), Enbrel (etanercept), Orencia (abatacept), Otezla (apremilast), Rinvoq (upadacitinib), Simponi (golimumab), Skyrizi (risankizumab), Taltz (ixekizumab), Tremfya (guselkumab), Xeljanz/Xeljanz XR (tofacitinib), ustekinumab] for treatment of the same indication; **and**
 - Authorization will be issued for 12 months

Ankylosing Spondylitis (AS) and Non-Radiographic Axial Spondyloarthritis (nr-axSpA)

Cosentyx is proven and medically necessary for the treatment of ankylosing spondylitis (AS) and non-radiographic axial spondyloarthritis (nr-axSpA) when all of the following criteria are met:

- For **initial therapy**, all of the following:
 - Diagnosis of active ankylosing spondylitis or non-radiographic axial spondyloarthritis; **and**
 - **One** of the following:
 - History of failure to **two** NSAIDs (e.g., ibuprofen, naproxen) at the maximally indicated doses, each used for at least 4 weeks, unless contraindicated or clinically significant adverse effects are experienced; **or**
 - Patient has been previously treated with a systemic targeted immunomodulator FDA-approved for the treatment of ankylosing spondylitis or nr-axSpA [e.g., adalimumab, Bimzelx (bimekizumab-bkzx), Cimzia (certolizumab), Enbrel (etanercept), Rinvoq (upadacitinib), Simponi (golimumab), Taltz (ixekizumab), Xeljanz (tofacitinib)]
 - and**
 - Cosentyx is initiated and titrated according to U.S. FDA labeled dosing for AS or nr-axSpA; **and**
 - Patient is not receiving Cosentyx in combination with a systemic targeted immunomodulator axSpA [e.g., adalimumab, Bimzelx (bimekizumab-bkzx), Cimzia (certolizumab), Enbrel (etanercept), Rinvoq (upadacitinib), Simponi (golimumab), Taltz (ixekizumab), Xeljanz (tofacitinib)] for treatment of the same indication; **and**
 - Prescribed by or in consultation with a rheumatologist; **and**
 - Initial authorization will be issued for 12 months
- For **continuation of therapy**, all of the following:
 - Documentation of positive clinical response; **and**
 - Cosentyx is initiated and titrated according to U.S. FDA labeled dosing for AS or nr-axSpA; **and**
 - Patient is not receiving Cosentyx in combination a systemic targeted immunomodulator [e.g., adalimumab, Bimzelx (bimekizumab-bkzx), Cimzia (certolizumab), Enbrel (etanercept), Rinvoq (upadacitinib), Simponi (golimumab), Taltz (ixekizumab), Xeljanz (tofacitinib)] for treatment of the same indication; **and**
 - Authorization will be issued for 12 months

Cosentyx (secukinumab) for intravenous injection is unproven and not medically necessary for the following (Cosentyx for self-administered subcutaneous injection is obtained under the pharmacy benefit):

- Plaque psoriasis
- Enthesitis-related arthritis
- Hidradenitis suppurativa

Applicable Codes

The following list(s) of procedure and/or diagnosis codes is provided for reference purposes only and may not be all inclusive. Listing of a code in this policy does not imply that the service described by the code is a covered or non-covered health service. Benefit coverage for health services is determined by the member specific benefit plan document and applicable laws that may require coverage for a specific service. The inclusion of a code does not imply any right to reimbursement or guarantee claim payment. Other Policies and Guidelines may apply.

HCPCS Code	Description
J3247	Injection, secukinumab, intravenous, 1 mg

Diagnosis Code	Description
L40.50	Arthropathic psoriasis, unspecified
L40.51	Distal interphalangeal psoriatic arthropathy
L40.52	Psoriatic arthritis mutilans
L40.53	Psoriatic spondylitis
L40.54	Psoriatic juvenile arthropathy
L40.59	Other psoriatic arthropathy
M08.1	Juvenile ankylosing spondylitis
M45.0	Ankylosing spondylitis of multiple sites in spine
M45.1	Ankylosing spondylitis of occipito-atlanto-axial region
M45.2	Ankylosing spondylitis of cervical region
M45.3	Ankylosing spondylitis of cervicothoracic region
M45.4	Ankylosing spondylitis of thoracic region
M45.5	Ankylosing spondylitis of thoracolumbar region
M45.6	Ankylosing spondylitis lumbar region
M45.7	Ankylosing spondylitis of lumbosacral region
M45.8	Ankylosing spondylitis sacral and sacrococcygeal region
M45.9	Ankylosing spondylitis of unspecified sites in spine
M45.A0	Non-radiographic axial spondyloarthritis of unspecified sites in spine
M45.A1	Non-radiographic axial spondyloarthritis of occipito-atlanto-axial region
M45.A2	Non-radiographic axial spondyloarthritis of cervical region
M45.A3	Non-radiographic axial spondyloarthritis of cervicothoracic region
M45.A4	Non-radiographic axial spondyloarthritis of thoracic region
M45.A5	Non-radiographic axial spondyloarthritis of thoracolumbar region
M45.A6	Non-radiographic axial spondyloarthritis of lumbar region
M45.A7	Non-radiographic axial spondyloarthritis of lumbosacral region
M45.A8	Non-radiographic axial spondyloarthritis of sacral and sacrococcygeal region
M45.AB	Non-radiographic axial spondyloarthritis of multiple sites in spine

Background

Cosentyx is a human IgG1 monoclonal antibody that selectively binds to the interleukin-17A (IL-17A) cytokine and inhibits its interaction with the IL-17 receptor. IL-17A is a naturally occurring cytokine that is involved in normal inflammatory and immune responses. Cosentyx inhibits the release of proinflammatory cytokines and chemokines.

Benefit Considerations

Some Certificates of Coverage allow for coverage of experimental/investigational/unproven treatments for life-threatening illnesses when certain conditions are met. The member specific benefit plan document must be consulted to make coverage decisions for this service. Some states mandate benefit coverage for off-label use of medications for some diagnoses or under some circumstances when certain conditions are met. Where such mandates apply, they supersede language in the benefit document or in the medical or drug policy. Benefit coverage for an otherwise unproven service for the treatment of serious rare diseases may occur when certain conditions are met. Refer to the Policy and Procedure addressing the treatment of serious rare diseases.

Clinical Evidence

The effectiveness of intravenous Cosentyx in the treatment of adult patients with active PsA, AS, and nr-axSpA was extrapolated from the established effectiveness of subcutaneous Cosentyx in adult patients with PsA, AS, and nr-axSpA based on pharmacokinetic exposure.

U.S. Food and Drug Administration (FDA)

This section is to be used for informational purposes only. FDA approval alone is not a basis for coverage.

Cosentyx (secukinumab) is a human interleukin-17A antagonist. Cosentyx (secukinumab) for IV injection indicated for the treatment of:

- Adult patients with active psoriatic arthritis
- Adult patients with active ankylosing spondylitis
- Adult patients with active non-radiographic axial spondyloarthritis with objective signs of inflammation

References

1. Cosentyx [package insert]. East Hanover, NJ. Novartis Pharmaceuticals Corp.; August 2025.
2. Ward MM, Deodhar, A, Gensler, LS, et al. 2019 Update of the American College of Rheumatology/Spondylitis Association of America/Spondyloarthritis Research and Treatment Network Recommendations for the Treatment of Ankylosing Spondylitis and Nonradiographic Axial Spondyloarthritis. *Arthritis & Rheumatology*. 2019; 71(10): 1599-1613.
3. Yu, DT, van Tubergen A. Treatment of axial spondyloarthritis (ankylosing spondylitis and nonradiographic axial spondyloarthritis) in adults. Sieper, J (Ed). UpToDate. Waltham, MA: UpToDate Inc. <http://www.uptodate.com>. Accessed on February 21, 2024.
4. Singh, JA, Guyatt, G, et al. 2018 American College of Rheumatology/National Psoriasis Foundation Guideline for the Treatment of Psoriatic Arthritis. *Arthritis & Rheumatology*. 2019; 71(1): 5-32.
5. Menter A, Gottlieb A, Feldman SR, et al. Guidelines of care for the management of psoriasis and psoriatic arthritis: Section 1. Overview of psoriasis and guidelines of care for the treatment of psoriasis with biologics. *J Am Acad Dermatol* 2008; 58(5):826-50.
6. Gottlieb A, Korman NJ, Gordon KB, et al. Guidelines of care for the management of psoriasis and psoriatic arthritis. Psoriatic arthritis: Overview and guidelines of care for treatment with an emphasis on the biologics. *J Am Acad Dermatol* 2008;58(5):851-64.

Policy History/Revision Information

Date	Summary of Changes
06/01/2026	<p>Applicable States Massachusetts and New York</p> <ul style="list-style-type: none">• Removed language indicating this Medical Benefit Drug Policy does not apply to the states of Massachusetts and New York <p>Nevada</p> <ul style="list-style-type: none">• Added instruction to refer to the UnitedHealthcare Commercial policy version for the state of Nevada <p>Coverage Rationale</p> <ul style="list-style-type: none">• Added language to clarify this policy refers to Cosentyx (secukinumab) for intravenous (IV) injection <i>for administration by a healthcare professional</i>• Replaced references to “targeted immunomodulator” with “<i>systemic</i> targeted immunomodulator” <p>Psoriatic Arthritis (PsA)</p> <ul style="list-style-type: none">• Updated list of examples of systemic targeted immunomodulators the patient must not be receiving in combination with Cosentyx for treatment of the same indication:<ul style="list-style-type: none">○ Removed Olumiant (baricitinib)○ Replaced “Xeljanz (tofacitinib)” with “Xeljanz/Xeljanz XR (tofacitinib)”• Updated list of examples of systemic targeted immunomodulators U.S. FDA-approved for the treatment of PsA with which the patient has been previously treated for initial therapy:<ul style="list-style-type: none">○ Added:<ul style="list-style-type: none">▪ Cimzia (certolizumab)▪ Orencia (abatacept)▪ Taltz (ixekizumab)

Date	Summary of Changes
	<p>Ankylosing Spondylitis (AS) and Non-Radiographic Axial Spondyloarthritis (nr-axSpA)</p> <ul style="list-style-type: none"> • Updated list of examples of systemic targeted immunomodulators the patient must not be receiving in combination with Cosentyx for treatment of the same indication: <ul style="list-style-type: none"> ○ Added Taltz (ixekizumab) ○ Removed: <ul style="list-style-type: none"> ▪ Orencia (abatacept) ▪ Olumiant (baricitinib) • Updated list of examples of systemic targeted immunomodulators U.S. FDA-approved for the treatment of AS or nr-axSpA with which the patient has been previously treated for initial therapy: <ul style="list-style-type: none"> ○ Added: <ul style="list-style-type: none"> ▪ Cimzia (certolizumab) ▪ Enbrel (etanercept) ▪ Taltz (ixekizumab) ○ Replaced “Xeljanz/Xeljanz XR (tofacitinib)” with “Xeljanz (tofacitinib)” <p>Supporting Information</p> <ul style="list-style-type: none"> • Updated <i>References</i> section to reflect the most current information • Archived previous policy version IEXD0132.03

Instructions for Use

This Medical Benefit Drug Policy provides assistance in interpreting UnitedHealthcare benefit plans. When deciding coverage, the member specific benefit plan document must be referenced as the terms of the member specific benefit plan may differ from the standard benefit plan. In the event of a conflict, the member specific benefit plan document governs. Before using this policy, check the member specific benefit plan document and any applicable federal or state mandates. UnitedHealthcare reserves the right to modify its Policies and Guidelines as necessary. This Medical Benefit Drug Policy is provided for informational purposes. It does not constitute medical advice.

UnitedHealthcare may also use tools developed by third parties, such as the InterQual[®] criteria, to assist us in administering health benefits. UnitedHealthcare Medical Benefit Drug Policies are intended to be used in connection with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice.