

Bariatric Surgery (for Kentucky Only)

Policy Number: CS007KY.11
Effective Date: May 1, 2025

[Instructions for Use](#)

Table of Contents	Page
Application	1
Coverage Rationale	1
Definitions	1
Applicable Codes	2
Description of Services	4
Clinical Evidence	7
U.S. Food and Drug Administration	31
References	32
Policy History/Revision Information	40
Instructions for Use	40

Related Policies
<ul style="list-style-type: none"> Minimally Invasive Procedures for the Treatment of Upper Gastrointestinal Diseases (for Kentucky Only) Obstructive and Central Sleep Apnea Treatment (for Kentucky Only) Robotic-Assisted Surgery Policy, Professional

Application

This Medical Policy only applies to the state of Kentucky.

Coverage Rationale

For medical necessity clinical coverage criteria, refer to the InterQual® CP: Procedures:

- Bariatric or Metabolic Surgery
- Bariatric or Metabolic Surgery (Adolescent)

[Click here to view the InterQual® criteria.](#)

The following procedures are unproven and not medically necessary for treating obesity due to insufficient evidence of efficacy:

- Bariatric surgery as the primary treatment for any condition other than obesity
- Bariatric interventions for the treatment of obesity including but not limited to:
 - Bariatric artery embolization (BAE)
 - Gastric electrical stimulation with an implantable gastric stimulator (IGS)
 - Gastrointestinal liners
 - Intragastric balloon
 - Laparoscopic greater curvature plication, also known as total gastric vertical plication
 - Mini-gastric bypass (MGB)/laparoscopic mini-gastric bypass (LMGBP)/one-anastomosis gastric bypass (OAGB)
 - Single-anastomosis duodenal switch [also known as duodenal switch with single anastomosis, or stomach intestinal pylorus sparing surgery (SIPS)]
 - Stomach aspiration therapy (AspireAssist®)
 - Transoral endoscopic surgery [includes TransPyloric Shuttle® (TPS®) device, endoscopic sleeve gastropasty]
 - Vagus nerve blocking (VBLOC®)

Definitions

Body Mass Index (BMI): A person's weight in kilograms divided by the square of height in meters. BMI can be used as a screening tool but is not diagnostic of the body fatness or health of an individual [Centers for Disease Control and Prevention (CDC), 2017].

The National Heart, Lung, and Blood Institute's (NHLBI) Practical Guide Identification, Evaluation and Treatment of Overweight and Obesity in Adults classifies the ranges of BMI in adults as follows:

- < 18.5 - Underweight
- 18.5 to 24.9 kg/m² – Normal Weight
- 25-29.9 kg/m² – Overweight
- 30-34.9 kg/m² – Obesity Class I
- 35-39.9 kg/m² – Obesity Class II
- ≥ 40 kg/m² – Obesity Class III

The American Society of Metabolic and Bariatric Surgeons (ASMBS; Pratt et al., 2018), classifies severe obesity in adolescents as follows:

- Class II obesity – 120% of the 95th percentile height, or an absolute BMI of 35-39.9 kg/m², whichever is lower*
- Class III obesity – 140% of the 95th percentile height, or an absolute BMI of ≥ 40 kg/m², whichever is lower

*Also as defined by the American Heart Association (Kelly et al., 2013).

Multidisciplinary: Bariatric center or regimen combining or involving several academic disciplines or professional specializations in an approach to create a well-trained, safe, and effective environment for the complex bariatric patient. Building the Multidisciplinary team includes staff such as the bariatric surgeon, obesity medicine specialist, registered dietician, specialized nursing, behavioral health specialist, exercise specialist and support groups [American Society for Metabolic and Bariatric Surgery (ASMBS) textbook of bariatric surgery].

Obstructive Sleep Apnea (OSA): The American Academy of Sleep Medicine (AASM) defines OSA as a sleep related breathing disorder that involves a decrease or complete halt in airflow despite an ongoing effort to breathe. OSA severity is defined as:

- Mild for AHI or RDI ≥ 5 and < 15
- Moderate for AHI or RDI ≥ 15 and ≤ 30
- Severe for AHI or RDI > 30/hr

For additional information, refer to the Medical Policy titled [Obstructive and Central Sleep Apnea Treatment \(for Kentucky Only\)](#).

Revisional Bariatric Surgery:

- Conversion – A second bariatric procedure that changes the bariatric approach from one procedure to a different type of procedure [e.g., sleeve gastrectomy or adjustable gastric band converted to Roux-en-Y (RYGB)]. Note: This is not to the same as an intraoperative conversion (e.g., converting from laparoscopic approach to an open procedure).
- Corrective – A procedure that corrects or modifies anatomy of a previous bariatric procedure to achieve the original desired outcome or correct a complication. These procedures also address device manipulation (e.g., gastric pouch resizing, re-sleeve gastrectomy, limb length adjustments in RYGB and gastric band replacement).
- Reversal – A procedure that restores original anatomy.

(Mirkin et al., 2021)

Applicable Codes

The following list(s) of procedure and/or diagnosis codes is provided for reference purposes only and may not be all inclusive. Listing of a code in this policy does not imply that the service described by the code is a covered or non-covered health service. Benefit coverage for health services is determined by federal, state, or contractual requirements and applicable laws that may require coverage for a specific service. The inclusion of a code does not imply any right to reimbursement or guarantee claim payment. Other Policies and Guidelines may apply.

Coding Clarification: Utilize CPT code 43775 to report laparoscopic sleeve gastrectomy rather than the unlisted CPT code 43659.

CPT Code	Description
0813T	Esophagogastroduodenoscopy, flexible, transoral, with volume adjustment of intragastric bariatric balloon
43290	Esophagogastroduodenoscopy, flexible, transoral; with deployment of intragastric bariatric balloon

CPT Code	Description
43291	Esophagogastroduodenoscopy, flexible, transoral; with removal of intragastric bariatric balloon(s)
43644	Laparoscopy, surgical, gastric restrictive procedure; with gastric bypass and Roux-en-Y gastroenterostomy (roux limb 150 cm or less)
43645	Laparoscopy, surgical, gastric restrictive procedure; with gastric bypass and small intestine reconstruction to limit absorption
43647	Laparoscopy, surgical; implantation or replacement of gastric neurostimulator electrodes, antrum
43648	Laparoscopy, surgical; revision or removal of gastric neurostimulator electrodes, antrum
43659	Unlisted laparoscopy procedure, stomach
43770	Laparoscopy, surgical, gastric restrictive procedure; placement of adjustable gastric restrictive device (e.g., gastric band and subcutaneous port components)
43771	Laparoscopy, surgical, gastric restrictive procedure; revision of adjustable gastric restrictive device component only
43772	Laparoscopy, surgical, gastric restrictive procedure; removal of adjustable gastric restrictive device component only
43773	Laparoscopy, surgical, gastric restrictive procedure; removal and replacement of adjustable gastric restrictive device component only
43774	Laparoscopy, surgical, gastric restrictive procedure; removal of adjustable gastric restrictive device and subcutaneous port components
43775	Laparoscopy, surgical, gastric restrictive procedure; longitudinal gastrectomy (i.e., sleeve gastrectomy)
43843	Gastric restrictive procedure, without gastric bypass, for morbid obesity; other than vertical-banded gastroplasty
43845	Gastric restrictive procedure with partial gastrectomy, pylorus-preserving duodenoileostomy and ileoileostomy (50 to 100 cm common channel) to limit absorption (biliopancreatic diversion with duodenal switch)
43846	Gastric restrictive procedure, with gastric bypass for morbid obesity; with short limb (150 cm or less) Roux-en-Y gastroenterostomy
43847	Gastric restrictive procedure, with gastric bypass for morbid obesity; with small intestine reconstruction to limit absorption
43848	Revision, open, of gastric restrictive procedure for morbid obesity, other than adjustable gastric restrictive device (separate procedure)
43860	Revision of gastrojejunal anastomosis (gastrojejunostomy) with reconstruction, with or without partial gastrectomy or intestine resection; without vagotomy
43865	Revision of gastrojejunal anastomosis (gastrojejunostomy) with reconstruction, with or without partial gastrectomy or intestine resection; with vagotomy
43881	Implantation or replacement of gastric neurostimulator electrodes, antrum, open
43882	Revision or removal of gastric neurostimulator electrodes, antrum, open
43886	Gastric restrictive procedure, open; revision of subcutaneous port component only
43887	Gastric restrictive procedure, open; removal of subcutaneous port component only
43888	Gastric restrictive procedure, open; removal and replacement of subcutaneous port component only
43999	Unlisted procedure, stomach
64590	Insertion or replacement of peripheral, sacral, or gastric neurostimulator pulse generator or receiver, requiring pocket creation and connection between electrode array and pulse generator or receiver
64595	Revision or removal of peripheral, sacral, or gastric neurostimulator pulse generator or receiver, with detachable connection to electrode array
64999	Unlisted procedure, nervous system

CPT® is a registered trademark of the American Medical Association

Description of Services

Obesity

Obesity is defined clinically using the [Body Mass Index](#) (BMI). Obesity is a significant health concern due to its high prevalence and associated health risks.

Health consequences associated with obesity include hypertension, Type II diabetes, hyperlipidemia, atherosclerosis, heart disease, stroke, diseases of the gallbladder, liver disease, osteoarthritis, Obstructive Sleep Apnea, and other respiratory problems. In addition, certain cancers are more prevalent in obese individuals, including endometrial, ovarian, breast, prostate, colon cancer, renal cell carcinoma, and non-Hodgkin's lymphoma.

The U.S. Preventive Services Task Force (USPSTF) recommends screening all adults for obesity. Clinicians should offer or refer patients with a BMI of 30 kg/m² or higher to intensive, multicomponent behavioral interventions (USPSTF, 2012).

Bariatric Surgery in the Adolescent Population

For adolescents, physical development and maturation may be determined utilizing the [gender specific growth chart and BMI chart](#) developed by the CDC, National Center for Health Statistics.

First-Line Treatments for Obesity

First-line treatments for obesity include dietary therapy, physical activity, behavior modification, and medication management; all of which have often been unsuccessful in long-term weight management for obese individuals (Lannoo and Dillemans, 2014).

Bariatric Surgical Procedures

The goal of surgical treatment for obesity is to induce significant weight loss and, thereby, reduce the incidence or progression of obesity-related comorbidities, as well as to improve quality of life. The purpose of performing bariatric surgery in adolescent individuals is to reduce the lifelong impact of severe obesity.

Surgical treatment of obesity offers two main weight-loss approaches: restrictive and malabsorptive. Restrictive methods are intended to cause weight loss by restricting the amount of food that can be consumed by reducing the size of the stomach. Malabsorptive methods are intended to cause weight loss by limiting the amount of food that is absorbed from the intestines into the body. A procedure can have restrictive features, malabsorptive features, or both. The surgical approach can be open or laparoscopic. The clinical decision on which surgical procedure to use is made based on a medical assessment of the patient's unique situation.

Roux-en-Y Bypass (RYGB)/Gastric Bypass

The RYGB procedure involves creating a stomach pouch out of a small portion of the stomach and attaching it directly to the small intestine, bypassing a large part of the stomach and duodenum.

Laparoscopic Adjustable Gastric Banding (LAGB)

The laparoscopic adjustable gastric banding procedure involves placing an inflatable silicone band around the upper portion of the stomach. The silicone band contains a saline reservoir that can be filled or emptied under fluoroscopic guidance to change the caliber of the gastric opening.

Vertical Sleeve Gastrectomy (VSG)

VSG can be performed as part of a two-staged approach to surgical weight loss or as a stand-alone procedure. A VSG involves the removal of 60-75% of the stomach, leaving a narrow gastric "tube" or "sleeve." This small remaining "tube" cannot hold as much food and produces less of the appetite-regulating hormone ghrelin, lessening a patient's desire to eat. VSG is not a purely malabsorptive procedure, so there is no requirement for lifetime nutritional supplementation (California Technology Assessment Forum, 2015).

Biliopancreatic Diversion With Duodenal Switch (BPD/DS) (also known as the Scopinaro Procedure)

BPD is primarily malabsorptive but has a temporary restrictive component. As in RYGB, three "limbs" of intestine are created: one through which food passes, one that permits emptying of fluids (e.g., bile) from digestive organs, and a common limb through which both food and digestive fluids pass. This procedure involves removal of the greater curvature

of the stomach instead of the distal portion. The two limbs meet in a common channel measuring only 50 to 100 cm, thereby permitting relatively little absorption.

Robotic-Assisted Surgery

Robotic surgery provides surgeons with three-dimensional vision, increased dexterity and precision by downscaling surgeon's movements enabling a fine tissue dissection and filtering out physiological tremor. It overcomes the restraint of torque on ports from thick abdominal wall and minimizes port site trauma by remote center technology (Bindal et al., 2015).

Transoral Endoscopic Surgery

Transoral endoscopic surgery is an option being explored for bariatric surgery. Natural orifice transluminal endoscopic surgery (NOTES) is performed via a natural orifice (e.g., mouth, vagina, etc.), and in some cases eliminates the need for abdominal incisions. This form of surgery is being investigated as an alternative to conventional surgery.

Transoral restorative obesity surgery (ROSE) is another endoscopic procedure. The endoscope with four channels is inserted into the esophagus and then the stomach. Specialized instruments are placed through the channels to create multiple folds around the existing stoma to reduce the diameter.

The Transpyloric Shuttle® (TPS®) device is a non-balloon, space occupying device with a 12-month treatment duration that is proposed as a new endoscopic bariatric therapy. The TPS device is comprised of a spherical silicone bulb connected to a smaller cylindrical silicone bulb by a flexible tether; it is delivered to and removed from the stomach using transluminal endoscopic procedures in the outpatient setting (Marinos, 2014). The device was granted FDA premarket approval on April 16, 2019, and was approved for up to 12 months weight loss therapy in individuals with a BMI of 35.0 kg/m² to 40.0 kg/m² or a BMI of 30.0 kg/m² to 34.9 kg/m² with one or more obesity-related comorbid condition. The device is intended to be used in conjunction with a diet and behavior modification program (ECRI, 2019).

Endoscopic Sleeve Gastroplasty (ESG) is a minimally invasive technique through the mouth that uses an endoscopic suturing device (e.g., OverStitch) to reduce gastric capacity by sealing off most of the stomach, forcing ingested food through an open tube of stomach tissue that connects the esophagus to the small intestine. ESG is similar to a laparoscopic sleeve gastrectomy in which the stomach is manipulated to create a tube-shape; however, no stomach tissue is removed.

Laparoscopic Mini Gastric Bypass (LMGBP)/One-Anastomosis Gastric Bypass (OAGB)

LMGBP/OAGB involves the construction of a gastric tube by dividing the stomach vertically down to the antrum. As in the RYGB, food does not enter the distal stomach. However, unlike gastric bypass surgery, digestive enzymes and bile are not diverted away from the stomach after LMGBP/OAGB. This can lead to bile reflux gastritis which can cause pain that is difficult to treat.

Implantable Gastric Stimulator (IGS)

IGS is a small, battery-powered device similar to a cardiac pacemaker, in a small pocket, created beneath the skin of the abdomen using laparoscopy. The IGS is programmed externally using a controller that sends radiofrequency signals to the device. Although the exact mechanism of action is not yet understood, gastric stimulation is thought to target ghrelin, an appetite-related peptide hormone (Gallas and Fetissov, 2011).

Vagus Nerve Blocking Neurostimulation Therapy (VBLOC)

VBLOC uses an implanted subcutaneous neurostimulator to deliver electrical pulses to the vagus nerve, which may suppress appetite (ECRI, 2016).

VBLOC therapy is designed to target the multiple digestive functions under control of the vagus nerves and to affect the perception of hunger and fullness.

Intragastric Balloon (IGB)

IGBs are acid-resistant balloons that are inserted into the stomach and expanded with saline or air. These space-occupying devices promote weight loss by creating a feeling of fullness, which can lead to reduced consumption of food. The devices are intended as an adjunct to diet, exercise, and behavioral counseling for the treatment of obesity (Hayes,

2021). Available clinical data and manufacturer recommendations indicate six months to be the current standard duration of therapy from insertion to removal (ASMBS, 2016).

Laparoscopic Greater Curvature Plication (LGCP) [also known as Total Gastric Vertical Plication (TGVP)]

LGCP is a restrictive procedure that involves folding and suturing the stomach onto itself to decrease the size of the stomach and requires no resection, bypass, or implantable device. This procedure is a modification of the gastric sleeve which requires surgical resection of stomach.

Stomach Aspiration Therapy

Stomach aspiration therapy, such as with the AspireAssist®, uses a surgically placed tube (endoluminal device) designed to aspirate a portion of the stomach contents after every meal (Hayes, 2021).

The AspireAssist is intended for long-term use in conjunction with lifestyle therapy (to help individuals develop healthier eating habits and reduce caloric intake) and continuous medical monitoring. Individuals must be monitored regularly for weight loss progress, stoma site health, and metabolic and electrolyte balance.

Bariatric Artery Embolization (BAE)

BAE is a minimally invasive procedure which is the percutaneous, catheter-directed, trans-arterial embolization of the left gastric artery (LGA). The procedure is performed by an interventional radiologist and targets the fundus that produces the majority of the hunger-controlling hormone ghrelin. Beads placed inside the vessels purportedly help decrease blood flow and limit the secretion of ghrelin to minimize feelings of hunger to initiate weight loss.

Gastrointestinal Liners

Gastrointestinal liners, such as the EndoBarrier™ system, utilize an endoscopically implanted sleeve into the stomach to reduce the stomach size. The sleeve is then removed after weight loss has been achieved. The EndoBarrier is not approved for use by the U.S. Food and Drug Administration (FDA) in the United States; it is limited by federal law to investigational use only.

Single-Anastomosis Duodenal Switch (SADS)

SADS is also called single-anastomosis loop duodenal switch, single-anastomosis duodenoileal bypass with sleeve gastrectomy, or stomach intestinal pylorus-sparing surgery—is a modification of biliopancreatic diversion with duodenal switch (BPD-DS). SADS consists of a sleeve gastrectomy to remove most of the stomach and an intestinal bypass to shorten the length of the small intestine and to allow bile and pancreatic digestive juices to mix with the food. SADS is typically performed laparoscopically as an inpatient procedure.

Revisional Surgery

The indications for Revisional Bariatric Surgery vary greatly depending on the index procedure performed and the nature of the complication. Some complications may be encountered during the acute postoperative recovery period (leaks, abscesses, fistulae, etc.). Prior to revisional surgery, individuals should undergo a thorough Multidisciplinary assessment and consideration of their individual risks and benefits from revisional surgery (Brethauer et al., 2014). It is important to determine if the poor response to primary bariatric surgery is due to anatomic causes that led to inadequate weight loss or weight regain or to the patient's postoperative behavior, such as not following the prescribed diet and lifestyle changes (e.g., consuming large portions, high-calorie foods, and/or snacks between meals; not exercising). Uncontrollable reflux may be a complication experienced by some individuals; first-line therapy for individuals who experience GERD after bariatric surgery includes dietary and lifestyle modification, alcohol, and smoking cessation, followed by acid-reducing medications (King et al., 2021).

The Metabolic and Bariatric Surgery Accreditation and Quality Improvement Program (MBSAQIP) is a national accreditation standard for bariatric surgery centers. In 2012, the American College of Surgeons (ACS) and the American Society for Metabolic and Bariatric Surgery (ASMBS) combined their individual accreditation programs into a single unified program. MBSAQIP works to advance safe, high-quality care for bariatric surgical patients through the accreditation of bariatric surgical centers. A bariatric surgical center achieves accreditation following a rigorous review process during which it proves that it can maintain certain physical resources, human resources, and standards of practice. All accredited centers report their outcomes to the MBSAQIP database (MBSAQIP, 2019).

Clinical Evidence

Jung et al. (2020) conducted a systematic review and meta-analysis of 22 studies with 2,141 patients to comprehensively evaluate the efficacy of different endoscopic bariatric procedures compared to lifestyle modification in the treatment of morbid obesity. Intra-gastric balloon, duodenal-jejunal bypass liner (DJBL), aspiration therapy, primary obesity surgery endoluminal (POSE) procedure, and botulinum toxin injection to the stomach were included and the meta-analysis determined the percentage of weight loss (%weight loss) and percentage of excess weight loss (%EWL). The results showed that the Obalon Balloon system was shown to have efficacy for both %weight loss and %EWL, its efficacy was not proven due to the small number of studies and comparatively low effect size. Aspiration therapy demonstrated effectiveness for weight reduction when compared to lifestyle modification. Gas-filled balloon and botulinum toxin injection did not show a significant difference in %weight loss or %EWL compared with the control. The authors concluded that all bariatric endoscopic procedures, with the exception of a gas filled balloon and botulinum toxin injection, show superior short-term efficacy compared with lifestyle modification. These findings are limited by lack of long-term efficacy and safety quality data. (The following publications previously cited in this policy, are included in this systematic review: Abu Dayyeh 2015b, Chang 2014, Courcoulas 2017, Gersin 2010, Schouten 2010, Sullivan 2013, Thompson 2017).

O'Brien et al. (2019) performed a systematic review and meta-analysis on 33 reports containing ten or more years of follow-up for patients that underwent bariatric surgery. The authors evaluated the long-term effectiveness of Roux-en-Y gastric bypass (RYGB), laparoscopic adjustable gastric banding (LAGB), or (BPD/DS). Results for gastric bypass surgery showed a weighted mean % EWL of 56.7% at 10 or more years with a mean of 55.4% EWL. Eleven reports addressing BPD/DS showed a mean of 74.1% EWL and two reports for sleeve gastrectomy showed a mean of 57.0% EWL. A longitudinal cohort study for the patients receiving LAGB showed patient weight loss reached a peak at the two-year follow-up and remained relatively stable through the next 18 years with a mean weight loss of 24.8 kg representing 47.2% EWL. The authors concluded that RYGB, LAGB and BPD/DS lead to substantial weight loss which continued for at least 10 years. Due to patient education and lap band design changes, revisional surgery has decreased significantly over the past eleven years. The findings are limited by lack of direct comparison between techniques and lack of comparison groups not undergoing surgical treatments. (The following publications previously cited in this policy, are included in this systematic review: Maciejewski 2016, Salminen 2018, Schauer 2017, Sethi 2016, Sheikh 2017, Topart 2017, Vinzens 2017).

Zhao and Jiao (2019) conducted a systematic review to determine whether LRYGB and LSG are equivalent for mid- and long-term weight loss, resolution of comorbidities and adverse events (AEs). Eleven RCTs were included in the meta-analysis and the authors found no significant difference in excess weight loss between LRYGB and LSG nor any significant difference for T2D improvement. This analysis did identify more postoperative early complications for LRYGB, but no difference between the two procedures in later postoperative period. Future studies should focus on the comparison of complication and comorbidities. Limitations included the variation in sample size among the included studies which may have created a bias, variation of patient age and preoperative BMIs which may have led to heterogeneity, and failure of subgroup analysis for reoperation rate. Additional studies are needed to determine the relative long-term efficacy of different bariatric surgeries. (Publication by Salminen 2018, which was previously cited in this policy, is included in this systematic review).

Lager et al. (2017) retrospectively studied 30-day postoperative complications as well as changes in weight, blood pressure, cholesterol, hemoglobin, hemoglobin A1C, and creatinine from baseline to two, six, 12, and 24 months postoperatively in 383 patients undergoing RYGB and 336 patients undergoing SG. Follow-up rates were 706/719 at two months, 566/719 at six months, 519/719 at 12 months, and 382/719 at 24 months. Baseline characteristics were similar in both groups except for higher weight and BMI in the SG group. The RYGB group experienced greater total body weight loss at 6, 12, and 24 months (41.9 vs. 34.6 kg at 24 months, $p < 0.0001$). Excess weight loss was 69.7 and 51.7% following RYGB and SG respectively at 24 months ($p < 0.0001$). Blood pressure improved significantly in both groups. Surgical complication rates were greater after RYGB (10.1 vs. 3.5%, $p = 0.0007$) with no significant difference in life-threatening or potentially life-threatening complications. Weight loss was greater following RYGB compared to SG at two years. The authors recommend that surgical intervention be tailored to surgical risk, comorbidities, and desired weight loss. Limitations included retrospective design which may have impacted patient selection and other biases, incomplete biochemical data as some patients did not return to clinic for routine blood draws and performed at specific institution.

Polega et al. (2017) conducted a matched cohort study of laparoscopic BPD/DS and SG to compare 30-day outcomes. Of the 741 patients who underwent BPD/DS or SG, two cohorts of 167 patients each were matched for age, sex, and BMI. Length of stay (LOS) was longer in the BPD/DS cohort (2.5 ± 0.9 days versus 2.1 ± 0.7 days, $p < .001$). There were no significant differences between the groups in relation to 30-day postoperative rates of leak (0.3% versus 0.6%, $p > 0.99$), bleed (0% versus .3%, $p > .99$), reoperation (1.2% versus .6%, $p > .99$), or readmission (3% versus 1.2%, $p = .45$). There

were no mortalities. After matching for age, sex, and BMI, the authors found no significant differences between BPD/DS and SG with regard to 30-day postoperative rates of leak, bleed, reoperation, readmission, or mortality.

Risstad et al. (2017) conducted a randomized clinical trial with 60 patients with body mass index 50-60 kg/m²) to investigate bile acid profiles up to five years after RYGB and BPD/DS. Total bile acid concentrations increased substantially over five years after both RYGB and BPD/DS, with greater increases in total and primary bile acids after BPD/DS. Higher levels of total bile acids at five years were associated with lower body mass index, greater weight loss, and lower total cholesterol.

Xie et al. (2016) prospectively evaluated Apnea-Hypopnea Index (AHI) and Functional Outcomes of Sleep Questionnaires Scores (FOSQ) pre- and post-operatively in patients undergoing bariatric surgery. A total of 167 subjects were studied. The median age was 46 (14-75) years and BMI 49 (36-69) kg/m². Ninety-two (55.0%) patients were diagnosed with OSA preoperatively. Fifty (54.0%) required positive airway pressure (PAP) therapy. The mean reduction in BMI post bariatric surgery was 12.2 ±4.52 kg/m² at 6.56 ±2.70 months. Eighty (87.9%) reported improved sleep quality reflected in improved scores in all domains of the FOSQ (p < 0.001, paired t-test). Improvement in FOSQ scores remained significant (p < 0.05) in those with and without OSA. Thirty-nine (90.7%) patients discontinued PAP due to resolution of daytime sleepiness. In conclusion, the authors identified that weight loss following bariatric surgery has a positive impact on sleep in patients with and without OSAS. The findings are, however, limited by lack of comparison group without bariatric surgery.

Giordano (2015) conducted retrospective comparative study of consecutive super-obese patients. Patients either underwent RYGB (n = 102) or LAGB (n = 79). Early complications and weight loss outcomes were comparable between the two groups in the short term. However, weight loss and excess weight loss percent at six and 12 months of follow-up was significantly higher in patients who underwent RYGB than LAGB.

Magallares et al. (2015) conducted a meta-analysis of 21 studies evaluating the mental and physical health-related quality of life (HR-QOL) measures with the Short Form-36 (SF-36) before and after bariatric surgery. Study authors reported that obese patients scored less in the mental health component of SF-36 prior to bariatric surgery (n = 2,680) compared with after surgery (n = 2,251). Similar results were observed in the physical health component of SF-36. Study authors concluded that obese patients experienced strong improvement in mental and physical QOL measures following surgery. The findings are limited by lack of comparison group.

A 2014 Cochrane Systematic Review of RCTs by Colquitt et al. found that surgery results in greater improvement in weight loss outcomes and weight associated comorbidities compared with non-surgical interventions, regardless of the type of procedures used. They noted the overall quality of evidence in this analysis to be moderate. When compared with each other, certain procedures resulted in greater weight loss and improvements in comorbidities than others. Outcomes were similar between RYGB and SG, and both of these procedures had better outcomes than AGB. However, in one RCT, the LRYGB procedure resulted in greater duration of hospitalization in two RCTs (4/3.1 versus 2/1.5 days) and a greater number of late major complications (26.1% versus 11.6%). For people with very high BMI, biliopancreatic diversion with duodenal switch resulted in greater weight loss than RYGB. Duodenojejunal bypass with sleeve gastrectomy and LRYGB had similar outcomes; however, this was based on one small trial. Isolated SG led to better weight-loss outcomes than AGB after three years follow-up. This was based on one trial only. Weight-related outcomes were similar between laparoscopic gastric imbrication and LSG in one trial. Across all studies adverse event rates and reoperation rates were generally poorly reported. The authors also found that most trials followed participants for only one or two years, therefore the long-term effects of surgery remain unclear. In addition, open RYGB, LRYGB and LSG led to losses of weight and/or BMI but there was no consistent picture as to which procedure was better or worse in the seven included trials. (The following publications previously cited in this policy are included in this systematic review: Dixon 2008, Mingrone 2012).

Biliopancreatic Diversion/Biliopancreatic Diversion With Duodenal Switch (BPD/DS)

Khalaj et al. (2020) conducted a cohort study comparing gastric bypass (GB) to sleeve gastrectomy (SG) and the effectiveness and safety of these two procedures. The authors evaluated 2,202 patients that underwent laparoscopic SG and 1,085 patients who underwent laparoscopic GB. The SG procedure was performed over a 36-F bougie and reinforced with an omental pouch; the GB procedure was performed as either RYGB or one anastomosis (OAGB). Evaluation of weight loss included body mass index change, percent of total weight loss, and percentage of excess weight loss. Type 2 diabetes mellitus (T2DM), hypertension (HTN), and dyslipidemia, as obesity-associated comorbidities, were assessed in all patients. There were no major complications identified which was recognized by a return to the operating room, prolonged hospital stays beyond seven days, or the need for re-admission. Quality of life (QoL) was assessed using the Iranian version of the Short-Form Health Survey which measured physical, social, and mental aspects of health. Patient follow up for both types of procedures occurred at 6, 12, and 24 months after surgery. The authors found no significant

differences between the two surgical groups; patients that underwent SG had a lower FPG and HbA1C when compared to the GB group. BMI was not significantly different between the two groups. Excess weight loss (EWL)% was 61.9 ± 15.7 , 74.8 ± 19.1 , and 75.0 ± 21.9 in the SG group and 62.7 ± 15.3 , 77.5 ± 18.4 , and 80.1 ± 20.8 in the GB group at 6-, 12-, and 24-month follow-ups, respectively. All patient comorbidities and QoL improved. The authors concluded that bariatric surgery is effective and safe for treatment of obesity; while both procedures are effective for weight loss, remission of obesity-associated comorbidities, and QoL, SG is associated with fewer complications and nutritional deficiencies.

Strain et al. (2017) reported nine-year outcomes of BPD/DS. Initially 284 patients received a BPD/DS; 275 patients (69.8 % women) age 42.7 years, BMI 53.4 kg/m² qualified for baseline analysis. Two hundred seventy-five patients were available in year 1; 275 patients in year 3; 273 patients in year 5; 259 patients in year 7; and 228 patients in year 9. Gender distribution was not different. BMI was 30.1 at 1 year and 32.0 at 9 years. Body fat was reduced to 26 % after 2 years. Nutritional problems developed in 29.8% of patients over the course of observation. There were significant positive changes in quality of life between baseline and year 1 for most patients. Data showed that after surgery, the resolution of comorbidities continued for the 9-year follow-up period. Weight loss during the first year was well maintained, resolving comorbidities, and improving quality of life. According to the authors, rates of surgical complications resemble other bariatric procedures; however long-term nutritional deficiencies are of concern. The findings are limited by lack of comparison group.

Bariatric Artery Embolization (BAE)

There is insufficient evidence for bariatric artery embolization and its outcomes for weight loss; additional robust RCTs are warranted for safety and efficacy along with long-term follow up.

Reddy et al. (2020) conducted a single-center, sham controlled, masked RCT to evaluate the efficacy of transcatheter bariatric embolization (TBE) for weight reduction in obesity. Participants were randomized to either sham procedure (n = 20) or TBE targeting the left gastric artery using embolic beads (n = 20). The primary efficacy endpoint was the difference in TBWL between the two groups at 6 months. All patients entered a lifestyle counseling program and follow-up was completed by physicians that were masked to allocated therapy. At 6 months, the TBWL for TBE in the intention to treat (ITT) population was 7.4 kg compared to 3.0 kg for sham procedure. The change in BMI at 6 months for ITT was -2.6 in TBE versus -1.1 in sham. The TBE ITT population did maintain the weight loss at 12 months. Patients within the sham group were unblinded at 6 months and permitted to crossover to TBE and then only initial group was followed for 12 months. Limitations included small sample size, single center, no control group after 6 months, and possibility that the efficacy of TBE was related to subject participation in weight management counseling as it is unknown if TBE alone would have an impact on obesity without lifestyle counseling. Additionally, four subjects withdrew consent after randomization and another three prior to the 6-month visit. Furthermore, the clinical significance of the effect, its long-term sustainability, and safety are unclear.

Hafezi-Nejad et al. (2019) conducted a systematic review and meta-analysis of case series investigating the safety and efficacy of left gastric artery (LGA) embolization as a bariatric procedure. Meta-regression was performed to assess associations of age, sex, body mass index, and ghrelin and leptin levels with weight change after LGA embolization were selected. Six case series published between January 2014 and April 2019, comprising 47 patients investigating the safety and/or efficacy of LGA embolization for weight loss were included in the meta-analysis. The results showed a mean weight loss of 8.68 kg (19.14 lbs.) after 12 months of follow-up, approximately 8% of baseline total body weight which is superior to weight loss from diet and exercise, and comparable to other more invasive interventions. Transient superficial mucosal ulcers were common after LGA embolization, and one case of major complications (severe pancreatitis, splenic infarct, and gastric perforation) was identified. There were considerable variations in patient age, sex distribution, and baseline characteristics among the studies. Significant variation was observed in the duration of follow-up, which ranged from three months to 20-24 months. Limitations of this study include variations in the indications for LGA embolization, study designs, embolization techniques, follow-up plans, dietary assessments, patient comorbidities, and availability of control subjects. The authors concluded that LGA embolization is an investigative method and not yet proven to be effective management for obesity. Larger studies are needed to expand these findings and determine other correlations of weight loss after LGA embolization. (Publications by Bai 2018, Syed 2016, and Weiss 2017, which were previously cited in this policy, are included in this systematic review).

Weiss et al. (2019) evaluated the safety and efficacy of bariatric artery embolization for up to twelve months following surgery in 20 severely obese patients [five of which are identified below in the Weiss et al. (2017) case series]. The primary endpoint was weight loss with additional end points assessed. Bariatric embolization was performed successfully in all participants. Participants experienced mean excess weight loss of 8.2% at one month, 11.5% at three months, 12.8% at six months and 11.5% at twelve months. The mean total weight loss was 7.6kg at twelve months. As a result of loss to follow-up, 18 participants remained at three months, 16 at six months, and 15 at twelve months. No major adverse

events (AE) were identified and only eleven minor AE occurred in eight participants. The authors found bariatric embolization is well tolerated and promotes clinically relevant weight loss in adults with severe obesity. Limitations included lack of comparison group, small sample size, insufficient data due to lack of continuous follow up for several participants, required weight management compliance before the embolization procedure on the first five participants only and a large portion of participants were African American thus overrepresenting that population.

Gastric Electrical Stimulator (GES)

While gastric electrical stimulation may provide benefit for obesity, additional well designed RCTs with long-term follow-up are warranted to demonstrate safety and efficacy.

In this 2020 first-in-human (early feasibility) multicenter, phase 1, open prospective cohort study, [Paulus et al. (2020)] the authors assessed the safety of the Exilis™ gastric electrical stimulation. They also sought to investigate whether the settings can be adjusted for comfortable chronic use in Class II or III obese patients. Meal intake and gastric emptying and motility were also evaluated. In this study, 20 obese patients were implanted with the Exilis system and amplitude was individually set during four amplitude titration visits. Subjects underwent two blinded baseline test days (GES ON vs. OFF), after which long-term, monthly follow-up continued for up to 52 weeks. The results suggested that this device is safe and caused no patient discomfort. At baseline food intake and satiety were not significantly different when the device was on or off, and significant weight loss occurred at week 26, with EWL of 14% at 52 weeks. The authors conclude that the data were comparable with studies of subjects on diet and/or exercise alone, but disappointing when compared to minimally invasive procedures, such as gastric banding or endoscopic gastroplasty. Furthermore, the authors did not observe changes in plasma glucose and insulin levels which other bariatric procedures are known to improve. The authors concluded that considerably more basic research is required before clinical use. Limitations included small sample size, lack of control group, and lack of long-term outcomes.

In a 12-month prospective multicenter study, Morales-Conde et al. (2018) monitored all participants (n = 47) up to 24 months after laparoscopic implantation of a closed-loop GES system (CLGES). Weight loss, safety, quality of life (QOL), and cardiac risk factors were analyzed. Weight regain was limited in the 35 (74%) participants remaining enrolled at 24 months. Mean %TBWL changed by only 1.5% between 12 and 24 months, reported at 14.8% (95% CI 12.3 to 17.3) and 13.3% (95% CI 10.7 to 15.8), respectively. The only serious device-/procedure-related AEs were two elective system replacements due to lead failure in the first 12 months, while improvements in QOL and cardiovascular risk factors were stable through 24 months. The authors conclude that during the 24-month follow-up, CLGES was shown to limit weight regain with strong safety outcomes, including no serious AEs in the second year. They hypothesize that CL GES and objective sensor-based behavior data combined to produce behavior change, and in their opinion supports GES as a safe obesity treatment with potential for long-term health benefits. Larger well-designed randomized controlled trials are needed to further evaluate the safety and efficacy of GES therapy in the treatment of obesity.

In a post-implant analysis, Alarcón Del Agua I, et al. (2017) evaluated possible preoperative predictors for obtaining clinically meaningful weight loss with GES. Ninety-seven obese participants in a prospective multicenter study conducted in nine European centers were implanted laparoscopically with the abiliti® CL GES system. The mean 12-month %EWL with CLGES was 35.1 ±19.7%, with a success rate of 52% and a failure rate of 19%. Significant predictors of success were BMI < 40 kg/m² and age ≥ 50 years, increasing probability of success by 22 and 29%, respectively. A low F1-cognitive-restraint score was a significant predictor of failure (p = 0.004). The best predictive model for success included F1-cognitive-restraint, F2-disinhibition, BMI < 40, and age ≥ 50 (p = 0.002). The authors concluded that age, preoperative BMI, and F1-cognitive-restraint and F2-disinhibition scores from a preoperative questionnaire are predictive of weight loss outcomes with closed-loop GES and may be used for patient selection.

In a systematic review, Cha et al. (2014) evaluated the current state regarding implantable gastric stimulators. Thirty-one studies consisting of a total of 33 different trials were included in the systematic review for data analysis. Weight loss was achieved in most studies, especially during the first 12 months, but only very few studies had a follow-up period longer than one year. Among those that had a longer follow-up period, many were from the Transcend(®) (Implantable Gastric Stimulation) device group and maintained significant weight loss. Other significant results included changes in appetite/satiety, gastric emptying rate, blood pressure and neurohormone levels or biochemical markers such as ghrelin or HbA1c, respectively. The authors conclude that although gastric electrical stimulation holds great promise, stronger evidence is required through more studies with a standardized way of carrying out trials and reporting outcomes, to determine the long-term effect of gastric electrical stimulation on obesity. (Publications by Shikora 2009, Sarr 2012, and Camilleri 2008, which were previously cited in this policy, are included in this systematic review).

Intragastric Balloon (IGB)

There is mixed evidence regarding the long-term efficacy and safety for intragastric balloons and their use with obesity; additional well designed RCTs and long-term data are warranted.

Based on a clinical evidence assessment by ECRI (2022), the evidence for the Spatz3® IGB is inconclusive. Assessment of two RCTs, three nonrandomized comparison studies, two case series, and two chart reviews assessing weight loss and adverse events for Spatz3® in adults with obesity revealed short-term clinically significant weight loss but whether these results were long-term remains to be seen. Limitations included small sample sizes, retrospective design of studies, lack of randomization, masking, and controls along with single-center focus. Large robust studies with long-term results are warranted and several ongoing clinical trials may address this in the future.

Zou et al. (2021) performed a systematic review and meta-analysis to evaluate the efficacy of the intragastric balloon (IGB) as an obesity management tool for metabolic dysfunction-associated fatty liver disease (MAFLD). Thirteen observational studies and one RCT met the inclusion criteria (624 participants in total). The results showed that over time, IGB therapy significantly improved the serum markers homeostasis model assessment of insulin resistance (HOMA-IR), alanine aminotransferase (ALT), aspartate aminotransferase (AST), and gamma-glutamyl transpeptidase (GGT) levels from baseline to follow-up. The authors concluded that IGB has the potential to become a multidisciplinary management tool of MAFLD, however IGB is a temporary measure, and if the patient cannot maintain an active lifestyle after the first balloon is removed, relapse of MAFLD is expected. Limitations include lack of comparison group; further RCTs are needed.

Hayes (2018, updated 2022) low-quality evidence suggests that IGBs have mixed results with regard to weight loss over the short term when used as an adjunct to diet and exercise. These devices are consistently associated with high AEs and all studies analyzed lacked long term follow up on maintaining weight loss and safety concerns.

A 2021 ECRI clinical evidence assessment on the Orbera® Intragastric Balloon System concluded that the evidence is inconclusive with mixed results, and shows the use of Orbera results in short-term, clinically significant weight loss in most patients; however, most patients regain weight, and by one year, the sustained weight loss has unclear clinical significance. Additional randomized studies are needed to determine whether Orbera use can reduce bariatric surgery risks for patients who are not surgery candidates and/or use the device to lose weight to become eligible for surgery. Additional studies that directly compare Orbera with other IGBs would also be useful.

In a multicenter, open-label industry-sponsored RCT, Abu Dayyeh et al. (2021, included in ECRI 2022 report above) investigated the safety and efficacy of the Spatz IGB in adults with obesity. 288 patients were randomly assigned to receive either the IGB plus dietary and exercise counselling or dietary and exercise counselling alone for 32 weeks. Inclusion criteria were patients aged 22-65 years, BMI of 30 kg/m² or greater for past two years, history of unsuccessful non-surgical weight loss methods and willingness to participate in the required dietary restrictions. The IGB was implanted via esophagogastroduodenoscopy (EGD) under conscious or monitored anesthesia sedation; depending on the patient's height an initial volume of 400 ml, 450 ml, 500 ml, or 550 ml was utilized. During the 32 weeks, all patients followed a 1000-1200 kcal/day diet and exercise plan. After 32 weeks, the IGB was removed, and patients were followed for another 24 weeks. Primary outcomes consisted of %TBWL and clinical responder rate, which was achieved by a decrease of at least 5% total bodyweight loss at 32 weeks. Mean %TBWL at 32 weeks was 15.0% (95% CI 13.9-16.1) in the IGB group versus 3.3% (2.0-4.6) in the control group (p < 0.0001). The authors found the adjustable IGB combined with lifestyle modification enabled significant weight loss over a period of 6 months with an observed acceptable safety profile. Limitations included no masking or sham intervention, and an approximately 20% loss to follow-up at 32 weeks. Future studies should assess the long-term safety of the device.

ECRI (2020) Health Technology Assessment focused on the safety and efficacy of the Elipse™ and Obalon®, two ingestible IGBs. The evidence was inconclusive, citing RCTs would be beneficial to determine whether any differences exist in weight loss and serious AE risks. Available clinical guideline recommendations on IGBs are mixed and none pertain to ingestible IGBs. Thus, major evidence gaps remain and additional comparative studies of ingestible and conventional IGB are needed.

Moore et al. (2019, included in the ECRI report above) performed a retrospective analysis of patients that underwent the Obalon Balloon System (OBS), a swallowable, gas-filled intragastric balloon system for weight loss. A web-based registry was accessed for the data on 1,343 patients with a starting BMI ≥ 25 kg/m². Nonserious and serious adverse events were reported in 14.2% and .15% of patients, respectively. Weight loss in the indicated use (BMI 30-40 kg/m²) was 9.7 ±6.1 kg and 10.0 ±6.1% TBWL. Weight loss in other BMI categories was 8.2 ±5.6 kg or 10.3 ±7.0% total body weight loss for BMI 25 to 29.9 kg/m²; and 11.6 ±7.8 kg or percent total body weight loss 9.3 ±6.0 for BMI > 40 kg/m². The authors concluded

that the OBS safe and effective at stimulating weight loss and provides practitioners with another tool to treat obese patients who have failed other weight loss programs. Limitations included lack of comparison group, the possible bias of a manufacture-sponsored study, variation with loss and behavior modification data collection, and lack of data collection for co-morbidities and metabolic data resulting in inability of data analysis for these areas.

Coffin et al. (2017, included in the Hayes 2021 report above) published findings from their multicenter randomized controlled trial, in which they compared six months of IGB or standard medical care (low-calorie diet, with bimonthly dietician evaluations) as bridge therapies to laparoscopic gastric bypass in super-obese patients (> 45 kg/m²). The surgery was performed at six months, shortly after removal of the IGB, and assessments were undertaken through 12 months. While the BMIs between groups were comparable at baseline, IGBs significantly reduced BMI by six months compared with standard care, with median BMI of 47.9 kg/m² for IGB patients and 50.7 kg/m² for control patients (p < 0.001). However, while the implanted IGB was effective on the short term, having the IGB before surgery did not impact postsurgical outcomes after 12 months (approximately six months post-surgery), the groups' BMIs were not significantly different at this time point (median BMI: IGB, 38.1 kg/m² versus standard care, 37.6 kg/m²; p = 0.56). The authors concluded that IGB insertion before LGBP induced weight loss but did not improve the perioperative outcomes or affect postoperative weight loss. Limitations of the study included short duration of the IGB intervention, poor recruitment rate, a higher than expected use of ICU facilities, and the poor weight loss in the IGB group.

Nunes et al. (2017) conducted a retrospective review of 2,002 patients who underwent an IGB procedure to determine its effectiveness with different degrees of obesity. A total of 946 patients were lost to follow-up. Overall, 40 (3.78%) had device removal due to intolerance, and 1,016 patients completed the 6-month treatment. The mean weight loss was 18.9%, excess weight loss 60.1% and a BMI reduction of 6.76 points. Six months after removal of the balloon 842 patients had continued follow-up (82.8%). At this time, weight loss was 19.84%, excess weight loss was 59.49%, and BMI reduction of 7.06 points. In all groups there was statistical difference between the times T0 and T1 and between T1 and T2 (p < 0.001). There was no statistical difference between T2 and T3, in any group. The authors concluded that IGB provided sustained weight loss in patients who remained in dietary follow-up for one year. The study is limited by lack of comparison group and high lost-to-follow up rate. Longer term outcomes with well-designed randomized clinical controlled trials are needed to further evaluate the IGB.

Saber et al. (2017) conducted a systematic review and meta-analysis to evaluate the efficacy and safety of intragastric balloon (IGB) treatment. A total of 20 RCTs involving 1,195 participants were identified. Weight loss results before and after three months were analyzed separately. The weight loss results of patients with and without IGB treatment were compared. A significant effect size was calculated that favored fluid filled IGBs over air-filled IGBs. Flatulence, abdominal fullness, abdominal pain, abdominal discomfort, and gastric ulcer were significantly more prevalent among IGB patients than among non-IGB control patients. No mortality was reported from IGB treatment. In the authors' opinion, IGB treatment, in addition to lifestyle modification, is an effective short-term modality for weight loss. However, there is not sufficient evidence confirming its safety or long-term efficacy.

The REDUCE pivotal trial (Ponce et al., 2015, included in the Hayes 2020 report above, and Jung 2020 systematic review) was a prospective, randomized controlled pivotal trial of a dual intragastric balloon to evaluate the safety and effectiveness of a dual balloon system plus diet and exercise in the treatment of obesity compared to diet and exercise alone. Participants (n = 326) with body mass index (BMI) 30-40 kg/m² were randomized to endoscopic dual balloon system (DBS) treatment plus diet and exercise (DUO, n = 187) or sham endoscopy plus diet and exercise alone (DIET, n = 139). Co-primary endpoints were a between-group comparison of % EWL and DUO subject responder rate, both at 24 weeks. Thereafter DUO patients had the DBS retrieved followed by 24 additional weeks of counseling; DIET patients were offered DBS treatment. Mean BMI was 35.4. Both primary endpoints were met. DUO weight loss was over twice that of DIET. DUO patients had significantly greater %EWL at 24 weeks 25.1% intent-to-treat (ITT), 27.9% completed cases (CC, n = 167) compared with DIET patients (11.3% ITT, p = .004, 12.3% CC, n = 126). DUO patients significantly exceeded a 35% response rate (49.1% ITT, p < .001, 54.5% CC) for weight loss dichotomized at 25% EWL. Accommodative symptoms abated rapidly with support and medication. Balloon deflation occurred in 6% without migrations. Early retrieval for non-ulcer intolerance occurred in 9%. Gastric ulcers were observed; a minor device change led to significantly reduced ulcer size and frequency (10%). The authors concluded that the dual balloon system was significantly more effective than diet and exercise in causing weight loss with a low adverse event profile. Additional RCT with longer follow-up are needed.

Laparoscopic Greater Curvature Plication (LGCP)

While laparoscopic greater curvature plication may appear to be safe for weight loss, additional robust RCTs with comparison groups and long-term data are needed.

In a 2023 single center retrospective analysis Park and Kim presented the weight loss and revision surgery rate outcomes of 75 patients following laparoscopic gastric greater curvature plication (LGGCP) surgery. The results showed that 13 out of 75 patients underwent revision surgery. The main reason for revision was weight regain, however chronic intermittent GERD, dyspepsia and chronic relapsing melena were also reasons. The mean body weight and BMI at initial LGGCP surgery were 207 lbs. (± 24) and 35.6 (± 3.9 kg/m²) respectively. Mean nadir body weight after LGGCP was 149 lbs. (± 13), and BMI was 25.8 (± 2.8 kg/m²). At revision, mean body weight was 196 lbs. (± 25), and BMI was 33.9 (± 4.2 kg/m²). The results showed that after 5 years, there was weight gain close to pre-surgery levels. The authors concluded that LGGCP as a primary surgery results in high rate of weight gain and the need for revisional surgery.

Doležalova-Kormanova et al. (2017) reported outcomes in a cohort of LGCP patients at five-year follow-up. Patients with complete weight data through five-year follow-up was 86.9%, (212/244). The ANOVA database indicated a significant BMI reduction out to two years ($p < 0.001$), a plateau at three and four years, and a moderate but significant BMI increase at five years ($p < 0.01$). Excess BMI loss at one, two, three, four, and five years was as follows: 50.7 \pm 9.1%, 61.5 \pm 8.1%, 60.2 \pm 7.0%, 58.5 \pm 7.0%, and 56.8 \pm 6.3%. At five years, 79.2% (168/212) of patients were successful; 20.8% (44/212) experienced a suboptimal weight outcome; mean weight regain, 9.2%. Cluster analysis identified four distinct LGCP patient profiles. Diabetes improvement rate was 65.5%. There were 12 reoperations (4.9%): four emergency (1.6%) and eight (3.3%) elective. There was no mortality. The authors concluded that based on their original cohort and a 56.8% Excess BMI loss and low rate of complications, LGCP proved to be safe and effective. The findings are limited by lack of comparison group. Additional long-term outcomes are needed to evaluate LGCP in comparison to other bariatric procedures.

In an 18-month prospective, observational, open-label study, Bužga et al. (2017) reported outcomes of 127 patients; 84 underwent LSG and 43, LGCP. LSG and LGCP were then compared during long-term follow-ups in terms of glycemic control, hormone and lipid secretion, and changes in body composition. Significant weight-loss and an improved body composition resulted from either procedure vs. baseline (i.e., pre-surgery), with levels of fasting glucose and glycated hemoglobin also showing statistically significant reductions (at 3 and 18 months for either surgery). Intergroup comparisons for glycemic parameters yielded no statistically significant differences. However, a dramatic reduction in ghrelin was detected following LSG, falling from pre-surgery levels of 140.7 to 69.6 ng/L by six months ($p < 0.001$). Subsequently, ghrelin levels increased, reaching 107.8 ng/L by month 12. Conversely, after LGCP, a statistically significant increase in ghrelin was seen, rising from 130.0 ng/L before surgery to 169.0 ng/L by month 12, followed by a slow decline. The authors concluded that although the data showed good metabolic outcomes following LGCP, this method was less effective than LSG, possibly due to its preservation of the entire stomach, including secretory regions.

Grubnik et al. (2016) compared two-year outcomes in a European prospective randomized controlled trial comparing LGCP versus LSG. A total of 54 patients with morbid obesity were allocated either to LGCP group ($n = 25$) or LSG group ($n = 27$). Main exclusion criteria were: ASA $>$ III, age $>$ 75 and BMI $>$ 65 kg/m² (2). There were 40 women and 12 men, and the mean age was 42.6 \pm 6.8 years (range 35-62). Data on the operation time, complications, hospital stay, BMI loss, % EWL, loss of appetite and improvement in comorbidities were collected during the follow-up examinations. One year after surgery, the mean %EWL was 59.5 \pm 15.4% in LSG group and 45.8 \pm 17% in LGCP group ($p > 0.05$). After two years, mean %EWL was 78.9 \pm 20 % in the LSG group and 42.4 \pm 18% in the LGCP group ($p < 0.01$). After three years, mean % EWL was 72.8 \pm 22 in the LSG group and only 20.5 \pm 23.9 in the LGCP group ($p < 0.01$). Loss of feeling of hunger after two years was 25% in LGCP group and 76.9% in the LSG group ($p < 0.05$). The comorbidities including diabetes, sleep apnea and hypertension were markedly improved in both groups after surgery. The authors concluded that the short-term outcomes demonstrated equal effectiveness of both procedures, but two-year follow-up showed that LGCP is not as effective as LSG as a restrictive procedure for weight loss.

Tang et al. (2015) conducted a meta-analysis to compare LGCP with LSG in terms of efficacy and safety. Eligible studies included one randomized controlled trial and three non-randomized controlled trials involving 299 patients. The meta-analysis demonstrated a significantly greater % EWL after LSG than LGCP at the follow-up time points of three months ($Z = 2.26$, $p = 0.02$), six months ($Z = 4.49$, $p < 0.00001$), and 12 months ($Z = 6.99$, $p < 0.00001$). The difference in the resolution of diabetes mellitus between these two approaches did not reach statistical significance ($p = 0.66$). According to the pooled data, LGCP was associated with more adverse events than LSG ($p = 0.01$). The operation time ($p = 0.54$) and postoperative hospital stay ($p = 0.44$) were comparable between the two groups. LGCP is inferior to LSG not only in terms of providing effective weight loss but also in terms of safety.

Mini-Gastric Bypass (MGB)/Laparoscopic Mini-Gastric Bypass (LMGB)/One-Anastomosis Gastric Bypass (OAGB)

Currently there is insufficient evidence regarding the long-term effectiveness and safety of one anastomosis gastric bypass (OAGB) for obesity and weight loss. Many studies on OAGB focus on short to medium term outcomes of 1-5

years. There is limited evidence regarding the long-term efficacy, weight maintenance, and complication rates beyond 5 years. Additional well designed RCTs are needed for assessing its sustained effectiveness and safety.

Robert et al. (2024) conducted a prospective, open-label, non-inferiority, randomized extension study aiming to report weight loss, metabolic, safety and efficacy of one anastomosis gastric bypass (OAGB) versus Roux-en-Y gastric bypass (RYGB) at the five-year follow-up. The original study, YOMEGA (NCT02139813) studied the outcomes at 24 months. Study criteria included individuals with a BMI of 40 kg/m² or more, or 35 kg/m² or more if at least one comorbidity such as hypertension, obstructive sleep apnea, dyslipidemia or arthritis is present. Ultimately, 253 participants were randomly assigned to OAGB (n = 129) or RYGB (n = 124). Five individuals did not undergo their assigned surgery and 14 were excluded from the post-protocol analysis. At the two-year follow-up point, the mean percentage excess BMI loss was -87.9% in the OAGB group and -85.8% in the RYGB group. After 5 years, the mean percentage excess BMI loss was -75.6% in the OAGB and -71.4% in the RYGB group. Remission of type 2 diabetes was similar in both groups. The most common adverse event was gastroesophageal reflux disease (GERD), occurring in 41% of the participants in the OAGB group versus 18% in the RYGB group. Although the mean percentage excess BMI weight loss was similar in both procedures, the study noted participants who were status post OAGB had a higher incidence of diarrhea, steatorrhea, and nutritional disorders. These adverse events suggest OAGB may result in a higher rate of malabsorptive effect. The higher rate of GERD after OAGB can be viewed as a limitation and raises the concern about the procedure's long-term outcomes. Therefore, OAGB and its long-term outcomes need to be further investigated.

In a 2023 systematic review and meta-analysis, Li et al. assessed the efficacy and safety outcomes of OAGB compared to the Roux-en-Y (RNY) procedure in eight randomized controlled trials that comprised a total of 931 patients. The mean preoperative BMI ranged from 42.6 to 53.5 kg/m². Due to inconsistent outcome measures being described in each study, the authors performed a meta-analysis using the post operative outcome measures of BMI, percent of excess weight loss (%EWL), or excess body mass index loss (EBMIL). The results showed that 6 months after surgery, BMI and %EWL did not show a statistically significant difference. Twelve months post-surgery, 4 articles showed OAGB resulted in better weight loss than RYGB for %EWL, and two articles showed OAGB had superior BMI reduction. Two articles reported 5-year outcomes and showed Five years no statistically significant differences in %EBMIL and BMI. Two articles reported intraoperative complications, for which there were no statistically significant differences between the two procedures. Three articles were included in the early postoperative complications OAGB showed fewer complications than RNY, 3 versus 8 serious complications, respectively. There was inconsistent reporting of obesity related illnesses across the studies, but articles that did report them included diabetes, hypertension, hyperlipidemia, and gastroesophageal reflux disease (GERD), and all showed a high rate of remission. The authors concluded that OAGB is not inferior to RNY in terms of weight loss and remission of comorbidities during the first 2 years post operatively but may have a higher incidence of malnutrition. Additional large sample and long-term randomized controlled trials are needed to verify these findings. (Eskandaros 2021, and Musella 2017 previously cited in this policy were included in this review)

Parmar et al. (2020) evaluated the role of one-anastomosis/mini-gastric bypass (OAGB-MGB) as a revisional/secondary procedure in patients who needed revisional bariatric surgery (RBS). A total of 17 studies were included in this systematic review with a total of 1,075 patients. The mean age was 43 years and 75% were female. The follow-up ranged from 6 to 60 months with a mean of 29 months. The following identifies the breakout of primary procedures performed: LAGB - 569 patients, SG – 397 patients, VBG – 105 patients, and lap gastric plication - five patients. The most common reason for RBS was poor response in 81%, followed by gastric band failure in almost 36% of patients. The mean BMI prior to RBS was 41.6 kg/m². Following the OAGB-MGB procedure, the mean %EWL was 50.8% at six months, 65.2% at one year, 68.5% at 24 months and 71.6% at five years. The author's conclusion suggests that OAGB-MGB is a safe and an effective choice for revisional surgery, however randomized studies and large prospective studies with long term follow-up are needed to validate these findings. Limitations included lack of comparison group or RCTs in analysis along with race and ethnicity differences which may have impacted the patient's eating habits, education, compliance, and expectations.

In a comparative effectiveness review from Hayes (2019, updated 2023) for primary bariatric surgery, the mini-gastric bypass/one-anastomosis gastric bypass (MGB-OAGB) was compared to RYGB and LSG separately. Data from two systematic reviews and 4 RCTs suggest an overall increase in percentage of weight loss with the MGB-OAGB procedure when compared to RYGB and LSG. The evidence also suggested MGB-OAGB may have a positive impact on resolution of T2D and HTN. However, additional long-term follow-up is warranted for further research on long-term follow-up, complications, adverse effects, and impact on nutrition.

In a prospective case series of 150 morbidly obese patients who underwent laparoscopic OAGB, lipid profiles were evaluated preoperatively and at different intervals during a two-year follow-up. The authors (Carbajo et al., 2017) reported a mean weight loss of 48.85 kg ±15.64 and mean % EWL of 71.87 ±13.41. kg. Total cholesterol and low-density lipoprotein (LDL) levels significantly decreased, and high-density lipoprotein (HDL) levels significantly increased which the

authors believe translate into theoretical relevant cardiovascular risk benefits. The findings are limited by lack of comparison group. Long-term randomized studies are needed to fully evaluate the impact of this procedure.

Lessing et al. (2017) conducted a retrospective analysis of all patients (n = 407) who underwent OAGB, reporting an average excess weight loss one year following surgery as 88.9 ±27.3 and 72.8 ±43.5% in patients that underwent primary and revision OAGB, respectively. Study limitations include lack of comparison group and single center data.

Wang et al. (2017) conducted a systematic review and meta-analysis to compare the safety and efficacy between laparoscopic mini-gastric bypass (MGB) and laparoscopic SG. Thirteen studies met the inclusion criteria of comparative studies between MGB and SG; patients were adults, with age ranging from 20 to 70 years old; at least one of the following endpoints was included: operation time, mortality, overall early complications, specific early complications, overall late complications, specific late complications, hospital stay, revision rate, remission rate of comorbidities, one-year %EWL or five-year % EWL. The authors observed that patients receiving mini-gastric bypass had more advantageous indexes than patients receiving sleeve gastrectomy, such as higher one-year EWL% (excess weight loss), higher five-year EWL%, higher T2D remission rate, higher hypertension remission rate, higher OSA remission rate, lower osteoarthritis remission rate, lower leakage rate, lower overall late complications rate, higher ulcer rate, lower GERD rate, shorter hospital stay and lower revision rate. No significant statistical difference was observed on overall early complications rate, bleed rate, vomiting rate, anemia rate, and operation time between MGB and SG. In their opinion, due to the biased data, small sample size and short follow-up time, the results of this review may be unreliable. RCTs with larger samples sizes are needed to compare the effectiveness and safety between MGB and SG. (Publications by Kansou 2016 and Plamper 2017, which were previously cited in this policy, are included in this systematic review).

Single-Anastomosis Duodenal-Ileal Switch (SADIS/SADI/SADS)

There is insufficient evidence regarding the safety and efficacy of the single-anastomosis duodenal switch (SADS) for obesity; additional robust RCTs with comparison groups along with long-term results are needed. Several clinical trials are in progress for the single-anastomosis duodenal switch; information can be found at <https://www.clinicaltrials.gov>.

Esparham et al. (2023) conducted a systematic review evaluating the efficacy and safety of SADI-S in mid- and long-term follow-up. The review included 10 studies with a total of 1707 individuals reporting outcomes with ≥ 3 years of follow-up. The majority of the articles were retrospective and due to the heterogeneity of included studies in regard to surgical technique and reported variables, the authors were unable to perform a meta-analysis. The percentage of excess weight loss (EWL) was 70.9%-88.7% at 6 years and 80.4% at 10 years. The more common late complications were malabsorption (6.3%) and GERD (3.6%). The remission rates of hypertension, diabetes, GERD, obstructive sleep apnea, and dyslipidemia were 62.9%, 81.3%, 53.2%, 60.9%, and 69.7%, respectively. While the authors conclude that SADI-S/SADS is a safe and effective surgical technique with durable weight loss and a high rate of comorbidity resolution, they also note that it is important to consider the potential risk and complications associated with this hypoabsorptive procedure and the need for long-term follow-up.

In a 2021 retrospective cohort study, Iranmanesh et al. compared short- and medium-term outcomes between the standard double-anastomosis duodenal switch (DADS), and single-anastomosis duodenal switch (SADS). Data of 107 patients was collected in the Ontario Bariatric Registry from a Canadian bariatric center of excellence between 2010 and 2019, with the primary outcome measurement weight loss at one- and two-years post-surgery. Short-term secondary outcomes included operative times, intra- and early postoperative complications, LOS, and 30-day readmissions. Medium-term secondary outcomes included late postoperative complications as well as nutritional deficiencies and persistent diarrhea at one- and two-years post-surgery. Of the 107 patients, 25 received SADS surgery and 82 received (DADS). Follow-up data was available for 59 patients at one year, and 47 after two years. The results showed similar %TWL at one year (23.6 versus 26.2) and two years (24.8 versus 30.2,) after surgery. Short- and medium-term outcomes were similar between groups. This study is limited by a small number of patients receiving the SADS procedure and large rate of lost-to-follow-up. Additional high-quality studies with longer follow up are necessary to validate these retrospective findings.

Pereira et al. (2021) conducted a prospective, observational cohort study of 112 patients receiving SADS or BPD/DS. Primary endpoints were BMI and TWL, and secondary endpoints included remission of obesity related disorders (T2D, hypertension, and dyslipidemia), nutritional deficiencies and post-operative complications. Eighty-three patients received SADS and 29 BPD/DS. There were no statistically significant differences between groups' demographic characteristics or clinical features, except for baseline weight and BMI, which were significantly higher in the BPD/DS group. Follow up times for SADS and BPD/DS ranged from an average of 40 months to 23 months, respectively. The results showed no significant differences in BMI and percent excess BMI loss (%EBMIL) between the groups, although the percentages of total weight loss observed from 12, 24, and 36 months were significantly higher after BPD/DS. Obesity related comorbidities resolved numerically better in the BPD/DS group than the SADS group, but it was not statistically significant.

Nutritional status was not consistently significant between the two procedures, and no differences were observed in surgical complications. Operative time and hospital stay was shorter for the SADS group. The authors concluded SADS is a simpler technique and shows similar results to BPD/DS. They acknowledged several limitations, including that there was a considerable numerical imbalance between the two groups, and the number of patients with a follow-up was small. Large scale randomized controlled clinical trials with long-term data are needed to confirm these results.

In a Medtronic funded study, Cottam et al. (2020) evaluated weight loss and one-year nutritional outcomes of the SADS procedure. 120 patients at six different sites were enrolled; participant inclusion criteria included BMI of 35-40 kg/m² with one obesity related comorbidity or a BMI of 40-60 kg/m² with no related comorbidity. Weight loss, comorbidities, quality of life, and AEs were followed post-procedure for 12 months. The authors found SADS to be an effective weight loss operation and the ability to reduce comorbid conditions, particularly diabetes. Limitations included lack of comparative cohort, patient loss to follow up and lack of long-term results for efficacy.

In a retrospective cohort study, Surve et al. (2017) compared biliopancreatic diversion with duodenal switch (BPD-DS) with single anastomosis duodenal switch (SIPS-stomach intestinal pylorus sparing surgery) at a single institution with two-year follow-up. One-hundred eighty-two patients received either a BPD-DS (n = 62) or SIPS (n = 120) procedure. BPD-DS and SIPS had weight loss at three months that were not statistically significantly different but %EWL was more with BPD-DS than SIPS at six, nine, 12, 18, and 24 months. Patient lost a mean BMI of 23.3 (follow-up: 69%) and 20.3 kg/m² (follow-up: 71%) at two years from the BPD-DS and SIPS surgery, respectively. However, patients who had undergone SIPS procedure had significantly shorter operative time, shorter length of stay, fewer perioperative and postoperative complications than BPD-DS (p < .001). There was no statistical difference between two groups for postoperative nutritional data such as vitamins D, B1, B12, serum calcium, fasting blood glucose, glycosylated hemoglobin (HbA1C), insulin, serum albumin, serum total protein, and lipid panel. The authors noted that as the BPD-DS procedures were done prior to SIPS, learning curve and experience may account for the post-operative complications. RCTs with larger patient populations and longer follow-up periods are needed to evaluate the SIPS procedure.

Cottam et al. (2016) conducted a retrospective matched cohort analysis to compare RYGB with SADS with 18-month follow-up. One-hundred eight patients received either a RYGB (n = 54) or SADS (n = 54). Regression analysis was used to compare weight loss outcomes as measured by BMI and weight loss percentages. The results failed to show statistically significant differences between the two procedures on weight loss at 18 months (39.6 vs. 41% weight loss, respectively). However, there were significantly more nausea complaints (26 vs. 5), diagnostic endoscopies (EGD) (21 vs. 3) and ulcers (6 vs. 0) with the RYGB than the SADS. The two-year outcomes for this same patient cohort had similar results (Cottam et al., 2017). RCTs with larger patient populations and longer follow-up periods are needed to validate these findings.

Stomach Aspiration Therapy

Currently there is insufficient evidence regarding the safety and efficacy of stomach aspiration therefore additional robust RCTs with comparison groups are needed along with long-term results.

A 2021 ECRI clinical evidence assessment on AspireAssist Gastric Aspiration Port (Aspire Bariatrics, Inc.) noted that evidence is somewhat favorable for AspireAssist when adding to lifestyle modification. It was noted to improve weight loss at 1 year which was maintained at up to 4 years, however, these findings are based on low-quality evidence from 2 systematic reviews and 1 single-arm extension of an RCT. It is unknown if AspireAssist therapy contributes to abnormal eating behaviors as only one single-arm extension of RCT reported too few events. Evidence limitations included risk of bias in most studies included in the systematic reviews due to small study size, lack of control group, or both. Additional larger RCTs are needed to confirm findings, especially in the long term, as well as to compare AspireAssist with other minimally invasive treatments.

Jirapinyo et al. (2020) conducted a systematic review and meta-analysis of five studies with a total of 590 patients to assess the outcomes of aspiration therapy (AT) (AspireAssist®) on obesity related comorbidities at one year follow up. Comorbidities included hypertension, hyperlipidemia, T2D, and NAFLD. Secondary outcomes were the amount of weight loss up to four years post operatively, and pooled serious adverse events (SAEs). The results showed after one year hypertension, hyperlipidemia, HbA1C, and NAFLD significantly improved. Weight loss at one year was 17% TWL (296 patients), two years 18.3% (174 patients), three years 18.6% (88 patients), and four years 18.6% (27 patients). The pooled SAE rate was 4.1% and included buried bumper, peritonitis severe abdominal pain, abdominal pain secondary to pre-pyloric ulcer and device malfunction requiring A tube replacement. Two studies reported a rate of persistent fistula following A-tube removal. The authors concluded that at one year AT resulted in significant improvement in metabolic function parameters and four years, patients maintained their significant weight loss of 18.6% of their baseline weight, meeting the definition of successful weight loss maintenance, and may improve access to treatment in obese patients with

concomitant comorbidities. The authors acknowledge the limitations of this study. The number of studies is small (to account for this, conference abstracts that met the a priori inclusion criteria were included in the analysis), and most of them were retrospective and observational in nature. Larger, high-quality studies with longer follow-up are required to validate these findings. (Publications by Sullivan 2013, Thompson 2017, and Nyström 2018, which were previously cited in this policy, are included in this systematic review).

In the post study of the PATHWAY Trial, Thompson et al. (2019) provide 4-year outcomes of the AT patients from the initial trial. Fifty-eight participants were enrolled in the follow-up study; of these 55 had achieved at least 10% TWL at the end of the first year. Of the 58 patients who enrolled in the follow-up study, 15, 21, and seven patients elected to have the A tube removed between years one and two, two and three, and three and four, respectively, thus withdrawing from the study but no loss to follow-up. The 43 patients who withdrew from the study between years two and four, 25 (58.1%) achieved at least 10% TWL. The mean %EWL of AT participants at years one, two, three, and four was 37.1 ± 27.6 ($n/n = 81/110$), 40.8 ± 25.3 ($n/n = 42/55$), 44.7 ± 29.7 ($n/n = 22/55$), and 50.8 ± 31.9 ($n = 15/55$), respectively. The clinical success rate for patients participating in the follow-up study was 40/58 (69%) at four years from A-tube placement. The authors concluded the AT is a safe and effective intervention for people with class II and III obesity and can achieve weight loss along with improvement of quality of life. Limitations of this study are the relatively small number of participants by the fourth year, participant commitment and the absence of weight loss data after A-tube removal. Additionally, the findings are limited by the design that only allowed continued follow-up of participants maintained at least 10% TWL from baseline at each year end and lack of comparison group for the long-term.

Norén and Forsell (2016) reported one- and two-year outcomes from their prospective observational study of 25 obese subjects to evaluate weight reduction and safety of AT with AspireAssist™. Twenty of the original 25 subjects completed the initial one-year treatment. These 20 subjects lost mean 54% of their excess weight. At two years, 15 subjects had lost mean 61% of their excess weight. This weight loss surpassed expectations and is nearly at the level of gastric bypass procedure and other major abdominal surgery for obesity. The subjects reported improved quality of life during treatment. There was neither mortality nor any event more severe than grade III-a according to Clavien-Dindo grading system. Limitation of this study is the combination of AT and cognitive behavioral therapy (CBT) without any control group. Long-term patency is still unknown.

Transoral Endoscopic Surgery [Including Transpyloric Shuttle® (TPS) Device]

The evidence for transoral endoscopic surgery for bariatric surgery is limited; additional studies including RCTs, long-term data including the safety and efficacy of the procedure are warranted.

In a brief from ECRI (2019), the evidence for the Transpyloric Shuttle® (TPS) device is inconclusive. The evidence is limited, indicating longer-term follow-up data is warranted. The RCT reviewed appeared to have a low risk of bias but results from a single trial were not conclusive and need independent confirmation in another controlled trial. The case series had a very high risk of bias due to small sample size, lack of a control group and randomization, and blinding. Both the RCT and case series report relatively short follow-up.

In a prospective, multicenter, single-arm, feasibility trial, Sandler et al. (2018) evaluate 32 obese subjects with a trans-oral endoscopic gastrointestinal bypass device. The device is a cuff attached to the distal esophagus by transmural anchors and connected to a 120-cm sleeve diverting undigested nutrients to the jejunum. Baseline data collected included body weight, vital signs, AEs, medications, HbA1c, fasting glucose, and lipids in addition to follow-up visits. The device status was endoscopically assessed every six months. At 12 months, the 32 subjects had lost an average of 44.8% of excess body weight, 17.6% of total body weight, 20.8 kg, and 7.5 BMI points. The authors concluded this study demonstrated the feasibility, safety, and efficacy of a fully trans-oral gastrointestinal bypass implant and that this endoscopic device may provide a valuable addition to the available treatment for the management of morbid obesity. However, this study is limited by lack of comparison group, small sample size and short-term follow-up.

Marinos et al. (2014) conducted a prospective, open-label, nonrandomized, single-center investigational clinical trial performed to evaluate the safety and efficacy of the transpyloric shuttle (TPS) device. The study enrolled twenty patients meeting the criteria in two cohorts with treatment periods of three and six months. Patients were required to be ≥ 18 and ≤ 55 years of age with a BMI between 30 and 50 kg/m². Before device placement, patients were provided with nutritional guidelines for a low-calorie diet and no additional dietary counseling was given after the initial consultation. Patients were placed under general anesthesia and the devices were deployed and retrieved with no complications. All 20 patients enrolled in the study had lost weight at the time of device removal. Both the three- and six-month patients had statistically significant improvements to the overall IWQOL-Lite score that exceeded the 7.7- to 12-point threshold to define a clinical change. All but two patients completed the planned treatment period; both patients had the device removed due to

complaints of epigastric pain. Limitations of the study were small participant size and short treatment duration. The authors concluded the TPS is a promising technology that has potential to benefit obese patients seeking to lose weight.

Eid et al. (2014) conducted a prospective, single-center, randomized, single-blinded study from July 2009 through February 2011, to investigate the safety and effectiveness of endoscopic gastric plication with the StomaphyX device vs. a sham procedure for revisional surgery in RYGB patients to reduce regained weight. Enrollment was closed prematurely because preliminary results indicated failure to achieve the primary efficacy end point in at least 50% of StomaphyX-treated patients. One-year follow-up was completed by 45 patients treated with StomaphyX and 29 patients in the sham treatment group. Primary efficacy outcome was achieved by 22.2% (10) with StomaphyX vs. 3.4% (1) with the sham procedure ($p < 0.01$). Patients undergoing StomaphyX treatment experienced significantly greater reduction in weight and BMI at three, six, and 12 months ($p \leq 0.05$). There was one causally related adverse event with StomaphyX, that required laparoscopic exploration and repair.

Endoscopic Sleeve Gastroplasty (OverStitch)

There is insufficient quality evidence regarding the safety and efficacy of endoscopic sleeve gastroplasty for obesity. Future studies including RCTs are needed to assess the safety and efficacy of this procedure along with long-term results.

In a 2024 evolving evidence review, Hayes evaluated the endoscopic sleeve gastroplasty (ESG) using the Apollo ESG System (Apollo Endosurgery inc.) for obesity. A review of the literature found minimal/weak support for using the Apollo ESG system for ESG to treat obesity based on six clinical studies, systematic reviews that included between 7 and 35 studies, and five practice guidelines. It was noted that there are ongoing clinical trials, however, it is unclear if these trials will provide useful information on the place of Apollo ESG System in the bariatric surgery landscape.

Weitzner et al. (2023) performed a systematic review to evaluate various endoscopic bariatric procedures using only randomized controlled trials (RCTs) and observational studies. The authors evaluated weight loss compared with conservative management, lifestyle modification and bariatric surgery. A total of thirty-seven studies which included a total of 15,639 individuals were included in the review. The primary outcomes included the percentage of total body weight loss (%TBWL), percentage of excess body weight loss (%EBWL) and adverse events. Secondary outcomes included data related to quality of life, and differences in hemoglobin A1C (HgbA1c) levels. ESG had less %TBWL, 4.7 – 14.4% compared to 18.8 – 26.5% after LSG at 6 months and 4.5-18.6% as compared to 28.4-29.3%, respectively, at the one-year follow-up. Additionally, the study did not demonstrate that endoscopic therapies resulted in significant differences in HgbA1c reduction compared to lifestyle modification. Despite endoscopic therapies resulting in greater weight loss compared to lifestyle modification, it did not result in greater weight loss when compared to bariatric surgery. The clinical utility of endoscopic bariatric procedures has not been convincingly addressed over regulator approved bariatric surgeries such as LSG. Ultimately, more robust data from RCTs or case-controlled studies are necessary.

In a 2023 retrospective study, Gudur et al. analyzed over 600, 000 patients in the Metabolic and Bariatric Surgery Accreditation and Quality Improvement Program database and compared short term (30 days) adverse events (AEs), readmissions, reoperations, and reinterventions in patients that underwent endoscopic sleeve gastroplasty (ESG) compared to sleeve gastrectomy (SG). A total of 6054 patients underwent ESG, and 597463 SG. The results showed that there was no significant difference in major AEs, but patients undergoing ESG had more readmissions, reoperations, and reinterventions. An additional analysis showed that chronic steroid use, renal insufficiency, and anticoagulation therapy contributed the most to the AEs in both groups. Race did not impact AEs after ESG, with an increased risk of AEs identified for Black patients after SG. This retrospective study is limited by a very short follow up period. The authors concluded that further prospective long-term evaluations of ESG versus SG with regards to safety and efficacy are needed.

Current evidence in an evolving technology report from Hayes (2022) identified four comparative studies and two systematic reviews which revealed minimal support for endoscopic sleeve gastroplasty (ESG) with the OverStitch device. Even though the OverStitch device is associated with clinically significant weight loss and fewer AEs, studies did not suggest the weight loss was more beneficial than an LSG.

Abu Dayyeh et al. (2022) conducted a randomised clinical trial to explore the safety and efficacy of endoscopic sleeve gastroplasty (ESG) with lifestyle modifications compared to lifestyle modification alone for the treatment of Class 1 and 2 obesity. Inclusion criteria was aged 21-65 with a BMI of 30 to less than 40 with a history of failure with non-surgical weight loss methods, and who agreed to comply with lifelong dietary restrictions required by this procedure. The primary outcome on efficacy was %EWL at 52 weeks. Secondary efficacy outcomes included proportion of patients with 25% or more EWL, % of total weight loss, and the proportion of patients with 5% or more and 10% or more of total weight loss. The effect of ESG on obesity related comorbidities and safety were also assessed. Seventy-seven participants were randomized to the

ESG plus moderate-intensity lifestyle modifications (ESG group), and 110 to the moderate-intensity lifestyle modifications alone (control group). During the first year, 12 follow-up visits were completed at weeks 1 and 4, and then every 4 weeks until the 52-week visit. The results showed ESG with lifestyle modifications, compared with lifestyle modifications alone, resulted in significant improvements in terms of weight loss, and metabolic comorbidities with no GERD incidence as seen with other bariatric surgeries. Adverse events included gastrointestinal symptoms such as pain, heartburn, nausea and vomiting which is not unexpected when acclimating post procedure. Three participants had a Clavien-Dindo grade 3 device or procedure related adverse event requiring intervention and included abscess, GI bleeding and one case of malnutrition requiring reversal of the ESG. The authors concluded that as a minimally invasive alternative to surgical sleeve gastrectomy, ESG is a safe and effective option for individuals that prefer a non-surgical option. This study is limited by the impact of the COVID-19 pandemic on study follow-up and participant retention, as well as a small number of participants.

Marincola et al. (2021) conducted a systematic review with meta-analysis to evaluate LSG versus ESG. The authors selected a total of sixteen studies for a total of 2188 participants. One randomized study and seven observational studies on LSG were selected and eight observational studies on ESG were included. The authors reviewed studies which included obese individuals with a baseline Body Mass Index (BMI) between 30 – 40 kg/m² and a minimum of one year follow-up. The mean BMI was 34.34 and 34.72 kg/m² for LSG and ESG, respectively. The mean percentage end weight loss was 80.32% for the LSG group and 62.20% for the ESG group. The mean adverse events rate was 0.19%. The study results indicated a moderate superiority of LSG versus ESG. Although the recent emergence of bariatric endoscopic techniques is a less invasive approach for the treatment of obesity, scientific evidence is still limited related to their outcomes. ESG was created as a more cost-effective endoscopic alternative to LSG, but there are very few comparative studies available. Therefore, there remains a need for a proper meta-analysis that combines data from the two different techniques. The studies revealed notable biases including the type of design, the loss to follow-up rate was not always reported in some studies and one study did not follow its own selection criteria and reported an average BMI lower than the minimum value set. The quality of the available studies is poor and additionally, valid studies to base guidelines on are necessary.

Singh (2020) conducted a systematic review and found eight studies addressing the OverStitch™ device which included a total of 1,859 patients. Studies were all observational and included single center and multicenter experiences. Primary outcomes measured were %TWL, %EWL, and SAE. The authors found the pooled mean %TWL at 6, 12, and 24 months was 14.86, 16.43, and 20.01. Similarly, % EWL at six, 12, and 24 months was 55.75, 61.84, and 60.40. The incidence of SAE was 2.26%, and no mortality was reported. Gastrointestinal bleeding was the most common documented SAE and was usually managed conservatively with packed red blood cell transfusion. Based on the analysis, the authors concluded that ESG is a promising technique with effective weight loss outcomes. Limitations included lack of controlled studies, lack of standardization definition for SAE and lack of long-term follow up data. (Publication by Lopez-Hava 2017, which was previously cited in this policy, was included in this systematic review).

Hedjoudje et al. (2020) conducted a systematic review and meta-analysis from eight studies which included 1,772 patients that underwent ESG. Primary outcome measurements included relative weight loss, decrease in BMI and relative estimated weight loss. Serious adverse events were reported in all studies with an occurrence of 2.2% and included 18 patients with pain or nausea that required hospitalization, nine patients that experienced upper GI bleeding, eight patients with perigastric leak or collection, one patient experienced pneumoperitoneum and one patient had a pulmonary embolism. The authors found the data suggested ESG gave way to significant sustained weight loss and safety. Patients had a BMI decrease of 5.6 kg/m², mean TBWL was 15.1% and relative EWL of 57.7%. These results appear to be sustained through 18-24 months of follow-up. Limitations included lack of control group, large loss to follow-up, lack of reporting for mild adverse events and lack of long-term outcomes; future studies are warranted.

Vagus Nerve Blocking

Currently there is insufficient quality evidence supporting the long-term effectiveness of vagus nerve blocking for obesity treatment; additional robust studies including randomization are warranted.

Apovian et al. (2017) reported the two-year outcomes from the ReCharge study among participants initially randomized to the active intervention. At 24 months, 123 (76%) vBloc participants remained in the trial. Participants who presented at 24 months (n = 103) had a mean EWL of 21% (8% TWL); 58% of participants had ≥ 5% TWL and 34% had ≥ 10% TWL. Among the subset of participants with abnormal preoperative values, significant improvements were observed in mean LDL (-16 mg/dL) and HDL cholesterol (+ 4 mg/dL), triglycerides (-46 mg/dL), HbA1c (-0.3%), and systolic (-11 mmHg) and diastolic blood pressures (-10 mmHg). QOL measures were significantly improved. Heartburn/dyspepsia and implant site pain were the most frequently reported AEs. The primary related serious AE rate was 4.3%. The findings are limited by lack of comparison group.

Morton et al. (2016) reported 12-month outcomes from the ReCharge study. Fifty-three participants were randomized to vBloc and 31 to sham. Qualifying obesity-related comorbidities included dyslipidemia (73%), hypertension (58%), sleep apnea (33%), and T2D (8%). The vBloc group achieved a %EWL of 33% (11% t%TWL) compared to 19% EWL (6% TWL) with sham at 12 months (treatment difference 14 percentage points, 95% CI, 7-22; $p < 0.0001$). Common AEs of vBloc through 12 months were heartburn/dyspepsia and implant site pain; the majority of events were reported as mild or moderate. The authors concluded that vBloc therapy resulted in significantly greater weight loss than the sham control among participants with moderate obesity and comorbidities, and with a well-tolerated safety profile. Longer-term outcomes are needed to demonstrate the continued durability of this procedure.

Shikora et al. (2016) reported two-year outcomes from the VBLOC DM² study, a prospective, case series of 28 subjects with T2DM and BMI between 30 and 40 kg/m² who underwent a VBLOC procedure. At 24 months, the mean percentage of EWL was 22% (95% CI, 15 to 28, $p < 0.0001$) or 7.0% TWL (95% CI, 5.0 to 9.0, $p < 0.0001$). Hemoglobin A1c decreased by 0.6 percentage points (95% CI, 0.2 to 1.0, $p = 0.0026$) on average from 7.8% at baseline. Fasting plasma glucose declined by 15 mg/dL (95% CI, 0 to 29, $p = 0.0564$) on average from 151 mg/dL at baseline. Among subjects who were hypertensive at baseline, systolic blood pressure declined 10 mmHg (95% CI, 2 to 19, $p = 0.02$), diastolic blood pressure declined by 6 mmHg (95% CI, 0 to 12, $p = 0.0423$), and mean arterial pressure declined 7 mmHg (95% CI, 2 to 13, $p = 0.014$). Waist circumference was significantly reduced by 7 cm (95% CI, 4 to 10, $p < 0.0001$) from a baseline of 120 cm. The most common AEs were mild or moderate heartburn, implant site pain, and constipation. The authors concluded that improvements in obesity and glycemic control were largely sustained after two years of treatment with VBLOC therapy with a well-tolerated risk profile. The findings are limited by lack of comparison group. Randomized controlled studies with larger patient populations are needed to validate these findings.

The ReCharge pivotal study, sponsored by the manufacturer, (Ikramuddin et al., 2014), was a prospective, randomized, double-blind, sham-controlled, multi-center trial to evaluate the safety and effectiveness of the Maestro system in treating obesity. The trial enrolled subjects who had a BMI 40-45 kg/m² or a BMI 35-39.9 kg/m² with at least one obesity-related co-morbid condition, and who had failed a more conservative weight reduction alternative. A total of 239 subjects were enrolled at 10 investigational sites; 162 subjects were randomized to the device group, and 77 were randomized to the sham control group. Subjects randomized to the sham control group underwent a surgical procedure consisting of anesthesia, implantation of a non-functional neuroregulator, and the same number of incisions an investigator would use during the laparoscopic placement of the leads. The study authors noted that the trial met its primary safety endpoint and helped more than half of patients lose at least 20% of their excess weight. The use of vagal nerve block therapy compared with a sham control device did not meet either of the prespecified coprimary efficacy objectives which were to determine whether the vagal nerve block was superior in mean percentage excess weight loss to sham by a 10-point margin with at least 55% of patients in the vagal block group achieving a 20% loss and 45% achieving a 25% loss.

Gastrointestinal Liner (EndoBarrier®)

Currently there is insufficient evidence regarding the effectiveness and safety of gastrointestinal liners for obesity and weight loss; additional well designed RCTs are needed along with long-term effects, and safety and efficacy results. Several clinical trials are in progress for the Endobarrier® device; information can be found at <https://www.clinicaltrials.gov>.

Ruban et al. (2022) conducted an RCT to study the clinical efficacy and safety of the duodenal-jejunal bypass liner (DJBL). Participants aged 18 to 65 years, with a BMI of 30 to 50 kg/m² and confirmed diagnosis of T2D for at least 1 year with inadequate glycemic control and on glucose-lowering medications were included in the trial. 170 patients were originally selected but due to several participants dropping out, 55 and 58 patients (DJBL and control arms, respectively) were included in the primary analysis at one year and 58 and 51 patients were included at year two. All participants received dietary and physical activity counselling. The primary outcome was to achieve an HbA1c reduction of 20% at 12 months post intervention. Secondary outcomes included lowered blood pressure, and a reduction in total body weight loss and the number of medications taken. The authors found that while the addition of the DJBL resulted in superior weight loss and improvement in cardiovascular risk factors, it did not make a significant impact on the patients' HbA1c. The findings are limited by the open-label design of the study and large loss to follow up that could have introduced biases.

Quezada et al. (2018) conducted a single-arm, open-label, case series to evaluate the safety and efficacy of endoscopically placed DJBL over a three-year period. Of 80 patients enrolled in the study, (age: 35 ±10 years; 69% female; weight: 109 ±17 kg; BMI: 42 ±5.4 kg/m²), 72 AEs were observed in 55 patients (68%). Nine subjects required a prolonged hospital stay and three subjects required major interventions. At 52 weeks (71 patients), 104 weeks (40 patients), and 156 weeks (11 patients), the mean %EWL were 44 ±16, 40 ±22, and 39 ±20, respectively ($p < 0.001$). This study shows significant and sustained weight loss after three years of treatment with the new DJBL. However, the high frequency and severity of AEs preclude the use of this prototype for periods longer than one year.

Forner et al. (2017) evaluated the outcomes of 114 obese patients treated with a DJBL. Mean total body weight change from baseline was 12.0 kg (SD 8.5 kg, $p < 0.001$). Over an average of 51 weeks, the mean %TWL was 10.5% (SD 7.3%). Mean HbA1c was not significantly improved, but of 10 patients on insulin, four ceased insulin and four reduced insulin dosages. There was a significant decrease in hemoglobin and total cholesterol and a significant increase in serum alkaline phosphatase. Seventy-four percent of patients experienced at least one AE, some of them serious including six device obstructions, five gastrointestinal hemorrhages, two liver abscesses, and one acute pancreatitis. Seventy-four percent of patients experienced weight gain after removal with a mean 4.5 ± 6.1 kg ($p < 0.0001$) within the first six months after explanation. The authors conclude that the DJBL provides significant but highly variable weight loss, and variable glycemic control. Most patients experienced an adverse event and most regained significant weight after device removal. In addition, the authors observed that major adverse events can occur, including the potentially life-threatening complications of hepatic abscess and gastrointestinal hemorrhage. The findings are limited by lack of comparison group. Further studies are needed to determine the long-term safety and efficacy of this procedure.

In a retrospective review, Betzel et al. (2017) evaluated the efficacy and safety profile of the DJBL. Inclusion criteria for treatment with a DJBL were age 18-70 years, BMI 28-45 kg/m² (2), and T2D with a HbA1c > 48 mmol/mol. Primary outcomes were changes in HbA1c and body weight. Secondary outcomes included changes in blood pressure, lipids, and anti-diabetic medication. Predictive factors for success of treatment with the DJBL were determined. The authors reported that 185 out of 198 patients successfully underwent a DJBL implantation procedure, with an intended implantation time of 12 months. In these 185 patients, body weight decreased by 12.8 ± 8.0 kg (total body weight loss of $11.9 \pm 6.9\%$, $p < 0.001$), HbA1c decreased from 67 to 61 mmol/mol ($p < 0.001$) despite a reduction in anti-diabetic medication, and blood pressure and serum lipid levels all decreased. In total, 57 (31%) DJBLs were explanted early after a median duration of 33 weeks. AEs occurred in 17% of patients. C-peptide ≥ 1.0 nmol/L and body weight ≥ 107 kg at screening were independent predictive factors for success. The authors concluded that treatment with the DJBL in patients with T2D and obesity resulted in improvement in glucose control, a reduction in anti-diabetic medication, and significant weight loss. The largest changes are observed within the first 3-6 months. Initial C-peptide levels and body weight may help to select patients with the greatest chance of success. The findings are limited by lack of comparison group.

Vilarrasa et al. (2017) evaluated the efficacy and safety of Endobarrier® in grade 1 obese patients with T2D and poor metabolic control and the role of gastro-intestinal hormone changes on the metabolic outcomes. Twenty-one patients aged 54.1 ± 9.5 years, diabetes duration 14.8 ± 8.5 years, BMI 33.4 ± 1.9 kg/m², and HbA1c 9.1 ± 1.3 %, under insulin therapy, were implanted with Endobarrier®. Fasting concentrations of PYY, ghrelin and glucagon, and AUC for GLP-1 after a standard meal test were determined prior to and at months 1 and 12 after implantation. They found that the Endobarrier® in in this subset of patients is associated with significant weight decrease and moderate reduction in HbA1c at month 12. Longer term outcome data is needed, and the findings are limited by lack of comparison group.

In a systematic review and meta-analysis, Rohde et al. (2016) evaluated the efficacy and safety of the DJBS. Five RCTs (235 subjects) and 10 observational studies (211 subjects) were included. The risk of bias was evaluated as high in all studies. The mean BMI ranged from 30 to 49.2 kg/m² (2) and 10-100% of the subjects had T2D. Meta-analysis showed that the DJBS was associated with significant mean differences in body weight and excess weight loss of -5.1 kg [95% confidence interval (CI) -7.3, -3.0; four trials; $n = 151$; $I^2 = 37\%$] and 12.6% (95% CI 9.0, 16.2; four trials; $n = 166$; $I^2 = 24\%$), respectively, compared with diet modification. The mean differences in glycated hemoglobin (-0.9%; 95% CI -1.8, 0.0) and fasting plasma glucose (-3.7 mM; 95% CI -8.2, 0.8) among subjects with T2D did not reach statistical significance. Adverse events consisted mainly of abdominal pain, nausea, and vomiting. No deaths occurred. Future high-quality long-term RCTs are needed to further assess efficacy and safety of the DJBS for obesity.

Clinical Practice Guidelines

American Diabetes Association (ADA)

The American Diabetes Association (ADA) *Standards of Medical Care in Diabetes – 2024* states that metabolic surgery should be considered as a weight and glycemic management approach for individuals with diabetes with a BMI ≥ 30.0 kg/m² (or ≥ 27.5 kg/m² in Asian American individuals) who are otherwise good candidates for surgery. They recommend that long-term lifestyle support and routine monitoring of micronutrient and nutritional status be provided to individuals after surgery.

The joint statement by international diabetes organizations on metabolic surgery in the treatment algorithm for type 2 diabetes (American Diabetes Association, International Diabetes Foundation, Diabetes UK, Chinese Diabetes Society, and Diabetes India) made the following recommendations:

- Metabolic surgery is recommended as an option to treat T2D in patients with the following conditions:
 - Class III obesity (BMI ≥ 40 kg/m²), regardless of the level of glycemic control or complexity of glucose-lowering regimens.

- Class II obesity (BMI 35.0 – 39.9 kg/m²) with inadequately controlled hyperglycemia despite lifestyle and optimal medical therapy.
- Metabolic surgery should also be considered and an option to treat T2D in patients with class I obesity and inadequately controlled hyperglycemia despite optimal medical treatment by either oral or injectable medications.
- All BMI thresholds used in these recommendations should be reconsidered depending on the ancestry of the patient. For example, for patients of Asian descent, the BMI values above should be reduced by 2.5 kg/m².

The organizations note that additional studies are needed to further demonstrate long-term benefits (Rubino et al., 2016).

American College of Gastroenterology (ACG)

In an ACG Clinical Guideline for the Diagnosis and Management of Gastroesophageal Reflux Disease (Katz, et al. 2022), the following recommendations are made:

- For refractory GERD, recommend optimization of PPI therapy as the first step in management of refractory GERD (Moderate quality of evidence/strong strength of evidence)
- For GERD management, recommend maintenance PPI therapy indefinitely or antireflux surgery for patients with LA grade C or D esophagitis (Moderate quality of evidence/strong strength of evidence)

American Gastroenterological Association (AGA)

In 2021, the AGA conducted a technical review on intragastric balloons (IGB) for the management of morbid obesity (Muniraj et al., 2021).

The review suggests that IGB therapy with lifestyle modification is an effective weight-loss intervention and seems to result in improvements in metabolic parameters and medical comorbidities. Several evidence gaps were addressed in this review and include long-term efficacy of IGB therapy compared with SOC beyond one year, variables such as the filling medium (fluid vs. gas) the potential efficacy of an ongoing dietary intervention, pharmacotherapy, or the need for sequential balloon placement for sustained weight loss, and the role of exercise in weight-loss sustainability. Although the risk of serious adverse events appears to be relatively low, early removal due to device intolerance seems to be relatively common. The AGA makes the following recommendations:

- In individuals with obesity seeking a weight-loss intervention who have failed a trial of conventional weight-loss strategies, suggest the use of IGB therapy with lifestyle modification over lifestyle modification alone. (Conditional recommendation, moderate certainty)
- In individuals with obesity undergoing IGB therapy, recommend moderate- to high-intensity concomitant lifestyle modification interventions to maintain and augment weight loss. (Strong recommendation, moderate certainty)
- In individuals undergoing IGB therapy, recommend prophylaxis with proton pump inhibitors. (Strong recommendation, moderate certainty)
- In individuals undergoing IGB therapy, suggest using the intraoperative anesthetic regimens associated with the lowest incidence of nausea along with perioperative antiemetics; suggest a scheduled antiemetic regimen for 2 weeks after IGB placement. (Conditional recommendation, low certainty)
- In individuals undergoing IGB therapy, suggest against perioperative laboratory screening for nutritional deficiencies. (Conditional recommendation, low certainty)
- Suggest daily supplementation with 1-2 adult dose multivitamins after IGB placement. (Conditional recommendation, very low certainty)
- After IGB removal, suggest subsequent weight loss or maintenance interventions that include dietary interventions, pharmacotherapy, repeat IGB or bariatric surgery. The choice of weight loss or maintenance method after IGB is determined based on patient's context and comorbidities following a shared decision-making approach. (Conditional recommendation, low certainty)

American Society for Gastrointestinal Endoscopy (ASGE)

The ASGE Technology Committee conducted a systematic review and meta-analysis to evaluate whether endoscopic technologies have met appropriate thresholds outlined by ASGE by the Preservation and Incorporation of Valuable endoscopic Innovations (PIVI) document (Abu Dayyeh et al., 2015a). The study authors evaluated Orbera intragastric balloon (IGB) (Apollo Endosurgery) and the EndoBarrier duodenal-jejunal bypass sleeve (DJBS) (GI Dynamics). Results of the meta-analysis (17 studies, n = 1,683) indicate that the Orbera IGB satisfies the PIVI thresholds for therapy for primary and non-primary bridge obesity. The percentage of EWL (% EWL) associated with the Orbera IGB at 12 months was 25.44% (95% CI, 21.45 to 29.41%) with a mean difference over controls of 26.9% (%EWL) (95% CI, 15.66% to 38.24%; p ≤ 0.01) in a total of three RCTs. The pooled %TWL after use of Orbera IGW was 13% at six months (95% CI, 12.37% to 13.95%) and 11.27% (95% CI, 8.17% to 14.36%), both which exceed the PIVI threshold of 5% TBWL for nonprimary bridge obesity therapy.

In its position statement on EBTs in clinical practice, the ASGE states that EBTs that have been approved by the FDA and meet thresholds of efficacy and safety as defined in the ASGE/ASMBS Preservation and Incorporation of Valuable Endoscopic Innovations should be included in the obesity treatment algorithm as adjunctive therapies to a lifestyle intervention program as outlined in the 2013 American Heart Association(AHA)/American College of Cardiology(ACC)/The Obesity Society (TOS) guidelines for the management of overweight and obesity in adults. ASGE advises that endoscopists performing EBT have a mechanism to enroll patients in long-term follow-up care for weight loss maintenance (Sullivan et al., 2015).

American Association of Clinical Endocrinologists (AACE)/Obesity Society/American Society for Metabolic and Bariatric Surgery (ASMBS)

In a clinical practice guideline for the perioperative nutritional, metabolic, and nonsurgical support of the bariatric surgery patient, the AACE, the Obesity Society, and the ASMBS (Mechanick, et al., 2019) cite the following:

- Patients with a BMI ≥ 40 kg/m² without coexisting medical problems and for whom bariatric surgery would not be associated with excessive risk should be eligible.
- Patients with a BMI ≥ 35 kg/m² and one or more severe obesity-related complications remediable by weight loss, including type 2 diabetes (T2D), high risk for T2D (insulin resistance, prediabetes, and/or metabolic syndrome), poorly controlled hypertension, nonalcoholic fatty liver disease/nonalcoholic steatohepatitis, obstructive sleep apnea, osteoarthritis of the knee or hip, and urinary stress incontinence, should be considered for a bariatric procedure. Patients with the following comorbidities and BMI ≥ 35 kg/m² may also be considered for a bariatric procedure, though the strength of evidence is more variable: obesity-hypoventilation syndrome and Pickwickian syndrome after a careful evaluation of operative risk; idiopathic intracranial hypertension; gastroesophageal reflux disease; severe venous stasis disease; impaired mobility due to obesity; and considerably impaired quality of life.
- Patients with BMI of 30–34.9 kg/m² and T2D with inadequate glycemic control despite optimal lifestyle and medical therapy should be considered for a bariatric procedure; current evidence is insufficient to support recommending a bariatric procedure in the absence of obesity.
- The body mass index criterion for bariatric procedures should be adjusted for ethnicity (e.g., 18.5 to 22.9 kg/m² is normal range, 23 to 24.9 kg/m² overweight, and ≥ 25 kg/m² obesity for Asians).
- Interventions should first include a multidisciplinary approach, including dietary change, physical activity, behavioral modification with frequent follow up; and then if appropriate, pharmacologic therapy and/or surgical revision.
- Selection of a bariatric procedure should be based on the individualized goals of therapy [e.g., weight loss and/or metabolic (glycemic) control], available local-regional expertise (surgeon and institution), patient preferences, and personalized risk stratification.

In addition, they recommend that all patients seeking bariatric surgery have a comprehensive preoperative evaluation. This assessment is to include an obesity-focused history, physical examination, and pertinent laboratory and diagnostic testing. A detailed weight history should be documented, including a description of the onset and duration of obesity, the severity, and recent trends in weight. Causative factors to note include a family history of obesity, use of weight-gaining medications, and dietary and physical activity patterns.

A brief summary of personal weight loss attempts, commercial plans, and physician-supervised programs should be reviewed and documented, along with the greatest duration of weight loss and maintenance. This information is useful in substantiating that the patient has made reasonable attempts to control weight before considering obesity surgery. The guidelines state that preoperative weight loss should be considered for patients in whom reduced liver volume can improve the technical aspects of surgery.

American Association of Clinical Endocrinologists (AACE)/American College of Endocrinology (ACE)

The AACE and the ACE developed comprehensive clinical practice guidelines for the medical care of patients with obesity (Garvey, et al., 2016) based on diligent review of clinical evidence with “transparent incorporation of subjective factors.” The final recommendations recognize that obesity is a complex, adiposity-based chronic disease, where management targets both weight-related complications and adiposity to improve overall health and quality of life. The detailed evidence-based recommendations allow for nuanced clinical decision-making that addresses real-world medical care of patients with obesity, including screening, diagnosis, evaluation, selection of therapy, treatment goals, and individualization of care. The goal is to facilitate high-quality care of patients with obesity and provide a rational, scientific approach to management that optimizes health outcomes and safety. Included in their clinical guideline are the following recommendations pertaining to BMI:

- Patients with a BMI of ≥ 40 kg/m² without coexisting medical problems and for whom the procedure would not be associated with excessive risk should be eligible for bariatric surgery.

- Patients with a BMI of ≥ 35 kg/m² and one or more severe obesity-related complications, including T2DM, hypertension, obstructive sleep apnea, obesity-hypoventilation syndrome, Pickwickian syndrome, nonalcoholic fatty liver disease or nonalcoholic steatohepatitis, pseudotumor cerebri, gastroesophageal reflux disease, asthma, venous stasis disease, severe urinary incontinence, debilitating arthritis, or considerably impaired quality of life may also be considered for a bariatric surgery procedure.
- Patients with a BMI of 30-34.9 kg/m² with diabetes or metabolic syndrome may also be considered for a bariatric procedure, although current evidence is limited by the number of patients studied and lack of long-term data demonstrating net benefit.
- Independent of BMI criteria, there is insufficient evidence for recommending a bariatric surgical procedure specifically for glycemic control alone, lipid lowering alone, or CVD risk reduction alone.

American Heart Association (AHA)/American College of Cardiology (ACC)/Obesity Society

The AHA/ACC and the Obesity Society published an updated 2013 Practice Guideline and Management of Overweight and Obesity in Adults (Jensen et al., 2014). The updated guidelines reflect such consensus and offer update regarding treatment for patients who are overweight or obese. While the focus remains on sustained weight loss and decreased waist circumference, the authors also recommend use of bariatric surgery for patients with a BMI ≥ 40 , or BMI ≥ 35 with comorbidities.

In a scientific statement on severe obesity in children and adolescents the American Heart Association (Kelly et al., 2013), summarized that RYGB has been associated with improvement or resolution of numerous comorbid conditions, including OSAS, T2DM, features of metabolic syndrome, pseudotumor cerebri, and psychosocial functioning. Controlled, prospective adult studies demonstrate a marked effect of bariatric surgery on mortality, comorbidity reversal, and prevention of comorbidity over ensuing decades; these beneficial effects of bariatric surgery help to inform clinical decision making for severely obese adolescents when no other treatments have demonstrated long-term effectiveness.

American Society for Metabolic and Bariatric Surgery (ASMBS) **Presurgical Evaluations**

The ASMBS published recommendations for the presurgical psychosocial evaluation of bariatric surgery patients (Sogg et al., 2016). They recommend that bariatric behavioral health clinicians with specialized knowledge and experience be involved in the evaluation and care of patients both before and after surgery. Given the importance of long-term follow up after weight loss surgery (WLS), the preoperative psychosocial assessment provides a valuable opportunity for patients to establish a trusted connection to a behavioral health provider as an additional resource and integral participant in their postoperative care. The need to ensure that postoperative psychosocial care is available has been noted in established practice guidelines and evidence suggests that such care is associated with better outcomes after surgery.

In a 2016 position statement on preoperative supervised weight loss requirements, the ASMBS noted that there is no data from any randomized controlled trial, large prospective study, or meta-analysis to support the practice of mandated preoperative weight loss. Further, there is no Level I data in the surgical literature, or consensus in the medical literature (based on over 40 published RCTs) that has clearly identified any one dietary regimen, duration or type of weight loss program that is optimal for patients with clinically severe obesity. Finally, they recommend that patients seeking surgical treatment for clinically severe obesity should be evaluated based on their initial BMI and co-morbid conditions.

Nutritional Impact of Bariatric Surgery

In an updated guideline on the integrated health nutritional guidelines for surgical weight loss, the ASMBS (Parrott et al., 2017) states that optimizing postoperative patient outcomes and nutritional status begins preoperatively. Patients should be educated before and after WLS on the expected nutrient deficiencies associated with alterations in physiology. Although surgery can exacerbate preexisting nutrient deficiencies, preoperative screening for vitamin deficiencies has not been the norm in the majority of WLS practices. Screening is important because it is common for patients who present for WLS to have at least one vitamin or mineral deficiency preoperatively.

Data continue to suggest that the prevalence of micronutrient deficiencies is increasing, while monitoring of patients at follow-up is decreasing. The ASMBS recommends that their guideline be considered a reasonable approach to patient nutritional care based on the most recent research, scientific evidence, resources, and information available. It is the responsibility of the registered dietitian nutritionist and WLS program to determine individual variations as they relate to patient nutritional care.

Indications for Surgery

In a joint update, the ASMBS and the International Federation for the Surgery of Obesity and Metabolic Disorders (IFSO) released revised guidelines on indications for metabolic and bariatric surgery (MBS) (Eisenberg et al., 2023). Updates to the guidelines include:

- MBS is recommended for individuals with a BMI ≥ 35 kg/m², regardless of presence, absence, or severity of comorbidities.
- MBS should be considered for individuals with metabolic disease and BMI of 30-34.9 kg/m².
- BMI thresholds should be adjusted in the Asian population such that a BMI ≥ 25 kg/m² suggests clinical obesity, and individuals with BMI ≥ 27.5 kg/m² should be offered MBS.
- Long-term results of MBS consistently demonstrate safety and efficacy.
- Appropriately selected children and adolescents should be considered for MBS.

Specific Bariatric Procedures

The ASMBS (2016, updated 2019) has approved, and supports the use of the following bariatric procedures and associated devices:

- Roux-en-Y Gastric Bypass
- BPD/Duodenal Switch
- Intra-gastric Balloon
- Sleeve Gastrectomy
- Adjustable Gastric Banding
- Single Anastomosis Duodeno-ileostomy with Sleeve

In a position statement (Ghiassi et al., 2024), the ASMBS indicated that the adoption of OAGB outside of the United States has resulted in numerous publications that report on the early, mid-, and long-term results. OAGB results in effective weight loss at 5 years and beyond, as well as metabolic effects that are comparable to RYGB or SG. Evidence has also shown that OAGB is effective as a revision option after restrictive operations such as LAGB, VBG, and SG. The authors note that since the majority of the peer-reviewed evidence on OAGB is retrospective with a few RCTs, well-designed RCTs to compare OAGB with other established bariatric procedures are encouraged.

In an updated statement (Kallies and Rogers, 2020) on the single-anastomosis duodenal switch (SADS), the ASMBS has concluded that single-anastomosis duodenoileal bypass with sleeve gastrectomy (SADI-S) provides for similar outcomes to those for the classic biliopancreatic diversion with duodenal switch (BPD-DS) procedure and therefore should be recognized. The society conclusion is that the current available peer-reviewed literature does not suggest outcomes will differ substantially from those seen with classic DS procedure. While the ASMBS endorses SADI-S as an appropriate bariatric surgical procedure, the society indicates publication of long-term safety and efficacy outcomes is still needed and is strongly encouraged; concerns remain about intestinal adaptation, nutritional issues, and long-term weight loss/regain following this procedure.

A 2017 ASMBS updated position statement on sleeve gastrectomy (SG) as a bariatric procedure (Ali et al., 2017) summarized that:

- Substantial long-term outcome data published in the peer-reviewed literature including studies comparing outcomes of various surgical procedures, confirm that sleeve gastrectomy (SG) provides significant and durable weight loss, improvements in medical co-morbidities, improved quality of life, and low complication and mortality rates for obesity treatment.
- SG is now the most commonly performed procedure in the United States (~53.8% of all bariatric procedures), followed by Roux-en-Y gastric bypass (RYGB; 23.1% of all procedures) (Chaar et al., 2018).
- In terms of initial early weight loss and improvement of most weight-related co-morbid conditions, SG and RYGB appear similar.
- SG is an acceptable option for a primary bariatric procedure or as a first-stage procedure in high-risk patients as part of a planned, staged approach.
- The effect of SG on GERD is less clear because GERD improvement is less predictable, and GERD may worsen or develop de novo. Preoperative counseling specific to GERD-related outcomes is recommended for all patients undergoing SG.
- Based on safety and efficacy data, there is a trend toward SG as the procedure of choice for adolescents, although both RYGB and SG are routinely performed in teen weight loss surgery programs.
- As with any bariatric procedure, long-term weight regain can occur after SG and may require one or more of a variety of re-interventions.

The ASMBS Clinical Issues Committee position statement on intragastric balloon therapy endorsed by SAGES (2016) includes the following summary and recommendations:

- Level 1 data regarding the clinical utility, efficacy, and safety of intragastric balloon therapy for obesity are derived from randomized clinical studies.
- Implantation of intragastric balloons can result in notable weight loss during treatment.
- Although utilization of intragastric balloons results in notable weight loss, separating the effect of the balloon alone from those of supervised diet and lifestyle changes may be challenging. Of note, recent FDA pivotal trials demonstrated a benefit to balloon use compared with diet alone in their study populations. In general, any obesity treatment, including intragastric balloon therapy, would benefit from a multidisciplinary team that is skilled and experienced in providing in-person medical, nutritional, psychological, and exercise counseling.
- The safety profiles for intragastric balloons indicate a safe intervention, with serious complications being rare. Early postoperative tolerance challenges can be significant but can be controlled with pharmacotherapy in the majority of patients, thereby minimizing voluntary balloon removals. These early symptoms should be discussed with the patient before the procedure.
- Although therapy with prolonged balloon in situ time and the use of sequential treatments with multiple balloons have been studied, awareness and adherence to absolute and relative contraindications of use and timely removal optimize device safety.
- Based on current evidence, balloon therapy is FDA approved as an endoscopic, temporary (maximum six months) tool for the management of obesity. Further review will evaluate the impact of diet, lifestyle changes, and pharmacotherapy during and after balloon removal.
- The ability to perform appropriate follow-up is essential when intragastric balloons are used for weight loss to enhance their safety and avoid complications related to spontaneous deflation and bowel obstruction.

The ASMBS (Moore and Rosenthal, 2018) released an addendum to their intragastric balloon therapy position statement in response to the FDA's warnings on complications not identified during initial clinical trials, and worldwide mortalities associated with intragastric balloons. They recommend that:

- As with all procedures, it is important that patients give informed consent and are aware of potential adverse events. Laypeople may need to be counseled to correct a misperception that endoluminal treatments are nonsurgical and thus risk-free.
- When less powerful treatments are chosen, behavioral modification increases in importance and there is risk of weight regain after the device is retrieved. The ASMBS routinely advocates for multidisciplinary care and support of the weight loss patient, and this recommendation is even more crucial for intragastric balloon recipients.

The ASMBS, in their 2015 position statement on vagal blocking therapy for obesity (Papasavas et al., 2015), conclude that the quantity of the data available at this time (six published studies; approximately 600 implanted devices) and the length of follow-up indicate adequate safety and efficacy in the short term. More prospective studies with longer follow-up are required to establish the clinically significant efficacy and patient tolerance of this device.

Bariatric Surgery in Adolescents

The updated ASMBS pediatric metabolic and bariatric surgery guidelines (Pratt et al., 2018) state that the disease of obesity has become recognized as a metabolic disease controlled by genetic factors, with clear evidence that the physiologic control of weight is through neuroendocrine pathways that regulate body mass by affecting satiety, hunger, and metabolism. The recognition that weight is largely not under volitional control leads to a strong need to offer effective, sustainable, proven therapies to children with obesity.

The summary of major changes in the guideline includes:

- Patient selection criteria of a BMI \geq 120% of the 95th percentile with a co-morbidity or a BMI \geq 140% of the 95th percentile should be used when determining weight cut offs for adolescents to undergo metabolic and bariatric surgery (MBS). In their opinion, Tanner stage and linear growth should not be used to determine readiness for MBS.
- Preoperative attempts at diet and exercise: there are no data that the number of weight loss attempts correlates with success after MBS. Compliance with a multi-disciplinary preoperative program may improve out-comes after MBS but prior attempts at weight loss should be removed as a barrier to definitive treatment for obesity.
- Requiring adolescents with a BMI $>$ 40 to have a co-morbidity (as in the old guidelines) puts children at a significant disadvantage to attaining a healthy weight. Earlier surgical intervention (at a BMI $>$ 45 kg/m²) can allow adolescents to reach a normal weight and avoid lifelong medication therapy and end organ damage from co-morbidities.
- Certain co-morbidities should be considered in adolescents, specifically the psychosocial burden of obesity, the orthopedic diseases specific to children, GERD, and cardiac risk factors. Given the poor outcomes of medical therapies for type 2 diabetes in children, these co-morbidities may be considered an indication for MBS in younger adolescents or those with lower obesity percentiles.

- Nonalcoholic fatty liver disease (NAFLD) and steatohepatitis (NASH): NAFLD may be present in at least 59% of adolescent patients referred for MBS. Given complete resolution of NASH in approximately 85% of patients who undergo VSG or RYGB, NAFLD should be considered a strong indication for MBS in adolescents with severe obesity.
- OSA has been shown to cause significantly decreased health-related quality of life (HRQoL) with increased risk of morbidity and mortality in adolescents. MBS in adolescents results in significant improvement or resolution of OSA. Thus, OSA should be considered a strong indication for MBS.
- Adolescents who suffer from severe obesity and have failed medical management of idiopathic intracranial hypertension should be considered for MBS.
- Adolescents with severe obesity have significant risk factors for cardiovascular disease (CVD), including hyperlipidemia, elevated inflammatory markers, hypertension, and insulin resistance. MBS significantly improves these risk factors, and therefore would be expected to decrease morbidity and mortality from CVD long term.
- Multidisciplinary teams should stabilize and treat preexisting eating disorders, assure stable social support, assess, and assist with nutrition and activity knowledge, and consider the addition of medications when appropriate.
- The Metabolic and Bariatric Surgery Accreditation and Quality Improvement Program (MBSAQIP) guidelines should be followed when building an adolescent MBS program. It is the responsibility of the adolescent MBS program to have a transition plan in place for adolescents to transition to an adult MBS program for lifelong care.

The ASMBS Pediatric Committee (Michalsky et al., 2012) best practice guidelines state that the associated risk/benefit analysis of bariatric surgery in adolescents should also include the consideration of the potential long-term health risks of untreated or inadequately treated obesity for the individual candidate. In addition, patients with a greater BMI and more serious medical illness are at increased risk of complications after bariatric surgery. Providing access to bariatric surgery earlier in life when the disease burden and severity is lower might decrease the operative risk, morbidity, and mortality. Additionally, earlier surgical intervention alters the natural course of many obesity-related co-morbidities that otherwise would put the patient at risk of long-term complications and early mortality.

Impact of Obesity and Obesity Treatment on Fertility and Fertility Therapy

In a position statement endorsed by the American College of Obstetricians and Gynecologists (ACOG) and the Obesity Society (Kominiarek et al., 2017), the ASMBS summarized that:

- Bariatric surgery is effective in achieving significant and sustained weight loss in morbidly obese women and has been shown in case-control studies to improve fertility.
- Pregnancy is not recommended during the rapid weight-loss phase after bariatric surgery; therefore, counseling and follow-up regarding contraception during this period is important.
- The specific impact of either medical weight-loss treatments or bariatric surgery on the responsiveness to subsequent treatments for infertility in both men and women is not clearly understood at this time.

Revisional Bariatric Surgery

In a systematic review of reoperative bariatric surgery, the ASMBS Revision Task Force (Brethauer et al., 2014) states that the indications and outcomes for reoperative bariatric surgery are procedure-specific, but the current evidence does support additional treatment for persistent obesity, co-morbid disease, and complications. Additional surgical therapy may benefit patients who present with insufficient weight loss, continued co-morbid disease, or weight gain after the index bariatric procedure. A thorough evaluation should be conducted by a multi-disciplinary program to determine the potential causes for their poor responses.

As the risks of reoperative bariatric surgery are higher than with the primary procedure, evidence suggests the need for careful patient selection. In addition, the specific type of reoperative procedure performed should be based on the patient's primary procedure, the patient's anatomy, the patient's weight and co-morbidities, and the experience of the surgeon.

An ASMBS Task Force (Sudan et al., 2015) on reoperative surgery provided the updated definitions for reoperative surgery as follows:

- Any operation after the first bariatric operation which qualified toward center of excellence volume requirements is considered a reoperation. Reoperations were further divided into corrective operations or conversions.
- An operation is considered corrective when complications or incomplete treatment effect of a previous bariatric operation was addressed but the initial operation was not changed.
- Conversions involve changing an index bariatric operation (first operation) to a different type of bariatric operation, and reversal restored original anatomy.

The Task Force also conducted a systematic review to evaluate morbidity, mortality, and weight loss outcomes after reoperative bariatric surgery. Data on reoperations was compared to that from patients who had initial bariatric operations but did not undergo reoperations. Reoperations were subdivided into corrective operations and conversions.

- Out of 449,753 bariatric operations, 28,720 (6.3%) underwent reoperations of which 19,970 (69.5%) were corrective and 8,750 (30.5%) were conversions.
- The mean % EBWL after conversion to a different bariatric operation was 39.3% and was 35.9% after a corrective operation. Although this % EBWL was lower than that after a primary operation (43.5%), it is still considered by the Task Force to be substantial and excellent weight loss. However, not all reoperations will result in further weight loss or resolution of comorbidity.
- Restorative operations necessitated by intolerable side effects or complications of the index procedure such as removal of the laparoscopic adjustable gastric band for band intolerance or dilated esophagus or reversing a duodenal switch or a gastric bypass for severe malabsorption, may in fact result in weight gain and return of comorbidities.
- Elderly patients (> 60 years of age) comprised 11% of the primary and 12% of the reoperative group of patients. The data suggests an overall improvement in the rates of morbidity and mortality after bariatric operations in recent years, even for higher risk populations.

The Task Force concluded that although most patients do not require reoperative surgery, among those who do, the complication rate is low, and outcomes are clinically comparable to primary procedures.

American Academy of Pediatrics (AAP)

In 2023, the AAP published the first edition of the clinical practice guideline for evaluation and management of children and adolescents with overweight and obesity. This document recommends metabolic and bariatric surgery for pediatric patients over the age of 12 for the following:

- Class II obesity, BMI \geq 35 or 120% of the 95th percentile for age and sex, whichever is lower with clinically significant disease, including but not limited to:
 - T2DM
 - Idiopathic intracranial hypertension
 - Non-alcoholic steatohepatitis
 - Blount's Disease
 - Slipped capital femoral epiphysis
 - GERD
 - OSA with an AHI > 5
 - Cardiovascular disease risks
 - Depressed health related QOL
- Class III obesity, BMI \geq 40 or 140% of the 95th percentile for age and sex, whichever is lower

Furthermore, the following is stated:

- The determination of eligibility for metabolic and bariatric surgery should rely heavily on a multicomponent and individualized approach between members of the metabolic and bariatric surgery team, the patient, and the patient's parents or guardians.
- A referral should be to a comprehensive metabolic and bariatric surgery center with experience and expertise in treatment of patients younger than 18 years.
- Evaluation for metabolic and bariatric surgery should include a holistic view of the patient and family, including individual needs (physical and psychosocial) and social risk factors.

American Society for Metabolic and Bariatric Surgery (ASMBS)/National Lipid Association (NLA)/Obesity Medicine Association (OMA)

The ASMBS, NLA and OMA published a 2-part joint scientific statement on lipids and bariatric procedures. Part 1 concluded that bariatric procedures reduce body fat and have favorable effects on adipocyte and adipose tissue function, which contributes to improvement in metabolic diseases such as dyslipidemia, high glucose levels, and high blood pressure. Among the mechanisms by which bariatric procedures may improve dyslipidemia includes favorable alterations in endocrine and inflammatory homeostasis. Bariatric procedures may also have favorable effects on bile acid metabolism and the intestinal microbiome, which may also improve dyslipidemia (Bays et al., 2016a).

Part 2 of this joint scientific statement summarized that the principles that apply to bariatric procedures and lipid levels include the following: (1) The greater the fat mass loss, the greater the improvement in lipid parameters such as triglycerides and especially LDL cholesterol; (2) bariatric procedures allow for a decrease in the use of drug treatment for

dyslipidemia; and (3) after bariatric procedures, HDL cholesterol may transiently decrease for the first three to six months after the procedure, which is usually followed by an increase in HDL cholesterol above the baseline value before the bariatric procedure. Finally, the authors observed that data are scarce regarding the effects of bariatric procedures on some of the lipid parameters such as non-HDL cholesterol, apolipoprotein B, and lipoprotein particle number and remnant lipoproteins (Bays et al., 2016b).

Endocrine Society

In its updated guideline for the assessment, prevention, and treatment of pediatric obesity (Styne et al., 2017) the Endocrine Society's recommendations include the following:

- Diagnose a child or adolescent > two years of age as overweight if the BMI is \geq 85th percentile but < 95th percentile for age and sex, as obese if the BMI is \geq 95th percentile, and as extremely obese if the BMI is \geq 120% of the 95th percentile or \geq 35 kg/m².
- Children or adolescents with a BMI of \geq 85th percentile should be evaluated for potential comorbidities.
- Insulin concentrations should not be utilized when evaluating children or adolescents for obesity.
- Bariatric surgery is suggested only under the following conditions:
 - The patient has attained Tanner 4 or 5 pubertal development and final or near-final adult height, the patient has a BMI of > 40 kg/m² or has a BMI of > 35 kg/m² and significant, extreme comorbidities.
 - T2DM, moderate to extreme sleep apnea, pseudotumor cerebri, debilitating orthopedic problems, and nonalcoholic steatohepatitis with advanced fibrosis.
 - Extreme obesity and comorbidities persist despite compliance with a formal program of lifestyle modification, with or without pharmacotherapy.
 - BMI of > 40 kg/m² with mild comorbidities (hypertension, dyslipidemia, moderate orthopedic problems, mild sleep apnea, nonalcoholic steatohepatitis, and extreme psychological distress that is secondary to their obesity).
 - Psychological evaluation confirms the stability and competence of the family unit [psychological distress due to impaired quality of life (QOL) from obesity may be present, but the patient does not have an underlying untreated psychiatric illness].
 - The patient demonstrates the ability to adhere to the principles of healthy dietary and activity habits.
 - There is access to an experienced surgeon in a pediatric bariatric surgery center of excellence that provides the necessary infrastructure for patient care, including a team capable of long-term follow-up of the metabolic and psychosocial needs of the patient and family.
- Bariatric surgery should not be performed in preadolescent children, pregnant or breast-feeding adolescents (and those planning to become pregnant within two years of surgery), and in any patient who has not mastered the principles of healthy dietary and activity habits and/or has an unresolved substance abuse, eating disorder, or untreated psychiatric disorder.

International Federation for the Surgery of Obesity and Metabolic Disorders (IFSO)

In a 2024 position statement (Ponce de Leon-Ballesteros et al.), the IFSO conducted a systematic review of evidence evaluating the SADI-S/SADS to guide clinical practice. While the IFSO endorsed the procedure in 2021 as safe and effective, they emphasized the need, at that time, for long-term multidisciplinary care and RCTs. For this review, there were a total of 93 articles included. Among the articles included, there were 1 RCT, 14 cohort studies, 25 case-control studies, 42 case series, and 11 case reports. The authors note that SADI-S/SADS demonstrated efficacy in weight loss and medium-to-long-term control of type 2 diabetes, as well as positive outcomes regarding hypertension and hyperlipidemia. The impact of SADI-S/SADS on other comorbidities remained inconclusive. The authors also note that there were frequent nutritional deficiencies identified included fat-soluble vitamins, anemia, and hypoalbuminemia. Limitations included considerable variation in the length of the common channel which may lead to differences in terms of weight loss outcomes as well as late complications and over-estimation of data due to the duplication of patients in the different studies. Despite significant efforts, there is a scarcity of high-quality evidence on SADI-S/SADS and the authors acknowledged that limited data was found beyond 5 years. The IFSO encouraged participation in national and international registries, publication of long-term follow-up studies, and RCTs to enhance the quality of evidence for SADI-S/SADS.

Society of American Gastrointestinal and Endoscopic Surgeons (SAGES)

A 2010 guideline by SAGES states that due to concerns for higher failure rates after fundoplication in the morbidly obese patient (BMI > 35 kg/m²) and the inability of fundoplication to address the underlying problem (obesity) and its associated co-morbidities, gastric bypass should be the procedure of choice when treating GERD in this patient group. The benefits in patients with BMI > 30 is less clear and needs further study (Stefanidis et al., 2010).

In its 2008 *Guidelines for Clinical Application of Laparoscopic Bariatric Surgery*, endorsed by the ASMBS, SAGES confirms that bariatric surgery is medically indicated for morbidly obese patients who fail to respond to dietary, behavioral, nutritional, and medical therapies, with clear evidence of efficacy and safety. BMI and age-based candidacy guidelines should not limit access for patients suffering with progressive or poorly controlled obesity-related comorbidities if the risk-versus-benefit analysis favors surgery. Laparoscopic RGB, AGB, and BPD have all been proven effective. They do not make a definitive recommendation for one procedure over another and note that at the present time, decisions are driven by patient and surgeon preferences, as well as considerations regarding the degree and timing of necessary outcomes versus tolerance of risk and lifestyle change.

Further, the 2008 guidelines state that there are no absolute contraindications to bariatric surgery. Relative contraindications to surgery may include severe heart failure, unstable coronary artery disease, end-stage lung disease, active cancer diagnosis/treatment, cirrhosis with portal hypertension, uncontrolled drug or alcohol dependency, and severely impaired intellectual capacity. Crohn's disease may be a relative contraindication to Roux-en-Y gastric bypass and biliopancreatic diversion.

Multidisciplinary Care Task Group

Greenberg et al. (2005) found a high incidence of depression, negative body image, eating disorders, and low quality of life (QoL) in patients with severe obesity and that perceived obesity-related health problems, motivation, and sense of coherence (SoC) predicted better weight loss. Although their investigation showed there are no predictive relationships between preoperative psychological evaluations and postoperative weight loss, the Behavioral and Psychological subgroup of the Multidisciplinary Care Task Group recommended that all bariatric surgery candidates be evaluated by a licensed mental health care provider experienced in the treatment of severely obese patients and working with a multidisciplinary team. Although research supports the association of psychological problems such as depression and personality disorder with less successful obesity surgery outcomes, rarely are the psychological problems cited as contraindications for surgery (Greenberg et al., 2005).

National Institute for Health and Care Excellence (NICE)

The National Institute for Health and Care Excellence (NICE) 2014 (updated 2023) guideline on obesity identification, assessment and management offers bariatric surgery as a treatment option for people with obesity when they have: a BMI of 40 kg/m² or more, or between 35 kg/m² and 39.9 kg/m² with a significant health condition (for example, type 2 diabetes or high blood pressure) that could be improved if they lost weight; have a multi-disciplinary team approach; and the individual agrees to necessary long-term follow-up after surgery. In addition, the NICE guideline notes that bariatric surgery should be expedited for individuals with a BMI of 35 kg/m² or more or 30 kg/m² to 34.9 kg/m² who have recent-onset (diagnosed within the past 10 years) type 2 diabetes and is receiving or will receive assessment in a specialist weight management service. Additionally, the guideline suggests consideration for individuals of South Asian, Chinese, other Asian, Middle Eastern, Black African or African-Caribbean family background using a lower BMI threshold (reduced by 2.5 kg/m²) to account for the fact that these groups are prone to central adiposity and their cardiometabolic risk occurs at a lower BMI. Further, surgical intervention is not generally recommended in children or young people, however it may be considered only in exceptional circumstances, and if they have achieved or nearly achieved physiological maturity.

A 2015 NICE interventional procedure guidance on managing type 2 diabetes states that current evidence on the safety and efficacy of implantation of a duodenal–jejunal bypass liner for managing type 2 diabetes is limited in quality and quantity. Therefore, the procedure should only be used in the context of research. Further research should give details of patient selection, including information about use of the procedure in patients with different levels of BMI. The research should provide information on complications; reasons for early removal of the device; medication used for treating type 2 diabetes, both when the device is in place and after its removal; and control of type 2 diabetes after device removal. In 2018, the following statement was added to this guidance: The device used in this procedure (EndoBarrier) no longer has a current CE mark. The CE mark is necessary for medical devices to be marketed in the European Union. A non-CE marked device can only be used in the context of clinical investigations with MHRA and research ethical approval.

Interventional procedures guidance [IPG569] from NICE (2016) states that the current evidence on the safety of single-anastomosis duodeno-ileal bypass with sleeve gastrectomy (SADI-S) for treating morbid obesity shows that there are well-recognized complications. The evidence on efficacy is limited in both quality and quantity. Therefore, this procedure should only be used with special arrangements for clinical governance, consent and audit or research.

A 2020 NICE interventional procedure guidance on swallowable gastric balloon for weight loss states that the evidence on efficacy is inadequate, and this procedure should only be done in a research setting.

A 2024 NICE interventional procedure guidance on ESG for obesity states that evidence on safety shows this procedure is safe in the short and long term. Evidence of efficacy shows that, when combined with lifestyle changes, people with a BMI over 30 kg/m² who have the procedure lose weight.

American Academy of Sleep Medicine (AASM)

The American Academy of Sleep Medicine commissioned a task force of experts in sleep medicine, otolaryngology, and bariatric surgery to develop recommendations based on a systematic review of the literature (Kent, 2021). The following are recommendations intended as a guide for clinicians who treat overweight adults with OSA:

- Recommend clinicians discuss referral to a sleep surgeon with adults with OSA and BMI < 40 who are intolerant or unaccepting of CPAP (STRONG).
- Recommend clinicians discuss referral to a bariatric surgeon with adults with OSA and obesity (class II/III, BMI ≥ 35) who are intolerant or unaccepting of PAP (STRONG).
- Suggest clinicians discuss referral to a sleep surgeon with adults with OSA, BMI < 40 and persistent inadequate PAP adherence due to pressure-related side effects (CONDITIONAL).
- Suggest clinicians recommend PAP as an initial therapy for adults with OSA and a major upper airway anatomic abnormality prior to consideration of referral for upper airway surgery (CONDITIONAL).

Department of Veterans Affairs (VA)/Department of Defense (DoD)

The 2020 guideline from the VA/DoD (Mayer et al., 2020) for the management of adult overweight or obesity makes the following suggestions or recommendations:

- In patients with a body mass index of ≥ 30 kg/m² and type 2 diabetes mellitus, suggest offering the option of metabolic/bariatric surgery, in conjunction with a comprehensive lifestyle intervention.
- In adult patients with a body mass index ≥ 40 kg/m² or those with body mass index ≥ 35 kg/m² with obesity-associated condition(s), suggest offering the option of metabolic/bariatric surgery, in conjunction with a comprehensive lifestyle intervention, for long-term weight loss/maintenance and/or to improve obesity-associated condition(s).
- In patients with obesity (body mass index ≥ 30 kg/m²) who prioritize short-term (up to six months) weight loss, suggest offering intragastric balloons in conjunction with a comprehensive lifestyle intervention.
- There is insufficient evidence to recommend for or against metabolic/bariatric surgery to patients over age 65.
- There is insufficient evidence to recommend for or against percutaneous gastrostomy devices for weight loss in patients with obesity.
- There is insufficient evidence to recommend for or against intragastric balloons for long-term weight loss to support chronic weight management or maintenance.

Thoracic Society

In a clinical practice guideline from the Thoracic Society (Hudgel, 2018), the following recommendations are made for patients who are overweight and suffer from OSA:

- Reduced-calorie diet; and
- Exercise or increased physical activity; and
- Behavioral guidance.

In addition, it was stated that pharmacological therapy and bariatric surgery are appropriate for selected patients who require further assistance with weight loss.

U.S. Food and Drug Administration (FDA)

This section is to be used for informational purposes only. FDA approval alone is not a basis for coverage.

Bariatric surgical procedures are not subject to FDA regulation. FDA approval information for several devices related to bariatric surgery is described below.

The FDA approved the ORBERA™ IntraGastric Balloon System (Apollo Endosurgery, Inc.) on August 5, 2015. The ORBERA System is indicated for use as an adjunct to weight reduction in obese adults with BMI ≥ 30 and ≤ 40 kg/m². It is to be used in conjunction with a long-term supervised diet and behavior modification program designed to increase the likelihood of significant long-term weight loss and weight loss maintenance. It is indicated for adults who have failed conservative weight reduction strategies, such as supervised diet, exercise, and behavior modification program. ORBERA has a maximum placement period of six months. For more information, refer to:

- <https://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfpma/pma.cfm?id=p140008>

- <https://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfpma/pma.cfm?id=P140008S016>
(Accessed October 2, 2024)

Gastric banding involves the use of an adjustable or nonadjustable gastric band, which is subject to FDA marketing approval. In 2001, the BioEnterics® LAP-BAND System was approved by FDA for marketing under the premarket approval process. According to the FDA labeling, this is approved for surgical treatment for severely obese adults for whom more conservative treatments (e.g., diet, exercise, behavioral modification) have failed. The LAP-BAND System is indicated for use in weight reduction for severely obese patients with a Body Mass Index (BMI) of at least 40 or a BMI of at least 35 with one or more severe co-morbid conditions, or those who are 100 lbs. or more over their estimated ideal weight according to the 1983 Metropolitan Life Insurance Tables (use the midpoint for medium frame). It is indicated for use only in severely obese adult patients who have failed more conservative weight-reduction alternatives, such as supervised diet, exercise, and behavior modification programs.

In February 2011, the FDA approved the Lap-Band Adjustable Gastric Banding System, by Allergan, for weight reduction in obese patients, with a Body Mass Index (BMI) of at least 40 kg/m² or less obese patients who have at least a body mass index (BMI) of 30 kg/m² and one or more additional obesity-related co-morbid condition, such as diabetes or hypertension. Additional information is available at: http://www.accessdata.fda.gov/cdrh_docs/pdf/p000008s017a.pdf. (Accessed October 2, 2024)

Adjustable gastric bands are contraindicated in patients younger than 18 years of age.

Surgical stapling devices are used in all bariatric surgical procedures except gastric banding. These devices have been approved by FDA for use in various general surgical procedures. One device is the Endo Gia Universal Auto Suture, which inserts six parallel rows of staples into tissue. Other surgical staplers are manufactured by Ethicon Endo-Surgery. Additional information, product code GDW and GAG, is available at: <http://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfRL/listing.cfm>. (Accessed October 2, 2024)

The OverStitch™ Endoscopic Suturing System was granted 510(k) marketing approval on June 27, 2018. According to the FDA, it is intended for endoscopic placement of suture(s) and approximation of soft tissue within the gastrointestinal tract. The device can utilize either a single- or dual-channel endoscope. Additional information is available at: https://www.accessdata.fda.gov/cdrh_docs/pdf18/K181141.pdf. (Accessed October 2, 2024)

The TransPyloric Shuttle/TransPyloric Shuttle Delivery Device was granted Premarket Approval on April 18, 2019 and is indicated for weight reduction in adult patients with obesity with a BMI of 35.0-40.0 kg/m² or a BMI of 30.0 to 34.9 kg/m² with one or more obesity related comorbid conditions and intended to be used in conjunction with a diet and behavior modification program. https://www.accessdata.fda.gov/cdrh_docs/pdf18/P180024a.pdf. (Accessed October 2, 2024)

In August of 2018, the FDA granted GI Dynamics Inc., Boston, MA an Investigational Device Exemption for the EndoBarrier® gastrointestinal liner. Additional information is available at: <https://www.fda.gov/medical-devices/how-study-and-market-your-device/investigational-device-exemption-ide>. (Accessed October 2, 2024)

References

Abu Dayyeh BK, Bazerbachi F, Vargas EJ, et al.; MERIT Study Group; Galvao Neto M, Zundel N, Wilson EB. Endoscopic sleeve gastropasty for treatment of class 1 and 2 obesity (MERIT): a prospective, multicentre, randomised trial. *Lancet*. 2022 Aug 6;400(10350):441-451.

Abu Dayyeh BK, Edmundowicz SA, Jonnalagadda S, et al. Endoscopic bariatric therapies. *Gastrointest Endosc*. 2015a; 81(5).

Abu Dayyeh BK, Kumar N, Edmundowicz SA, et al. ASGE Bariatric Endoscopy Task Force systematic review and meta-analysis assessing the ASGE PIVI thresholds for adopting endoscopic bariatric therapies. *Gastrointest Endosc*. 2015b; 82(3):425-438.

Abu Dayyeh BK, Maselli DB, Rapaka B, et al. Adjustable intragastric balloon for treatment of obesity: a multicentre, open-label, randomised clinical trial. *Lancet*. 2021 Nov 27;398(10315):1965-1973.

Alarcón Del Agua I, Socas-Macias M, Busetto L, et al. Post-implant analysis of epidemiologic and eating behavior data related to weight loss effectiveness in obese patients treated with gastric electrical stimulation. *Obes Surg*. 2017 Jun;27(6):1573-1580.

Ali M, El Char M, Ghiassi S, et al. ASMBS guidelines/statements. American Society for Metabolic and Bariatric Surgery updated position statement on sleeve gastrectomy as a bariatric procedure. *Surg Obes Relat Dis*. 2017 Oct;13(10):1652-1657.

American Academy of Pediatrics (AAP). Clinical practice guideline for the evaluation and treatment of children and adolescents with obesity. February 2023. Available at: <https://publications.aap.org/pediatrics/article/151/2/e2022060640/190443/Clinical-Practice-Guideline-for-the-Evaluation-and?autologincheck=redirected>. Accessed October 2, 2024.

American Diabetes Association Professional Practice Committee. Obesity and weight management for the prevention and treatment of type 2 diabetes: Standards of care in diabetes – 2024. *Diabetes Care* 2024 Jan 1; 47(Suppl 1):S145-S157.

American Diabetes Association (ADA). Standards of medical care in diabetes – 2017. *Diabetes Care* 2017;40(Suppl.1):S1–S2. American Society for Metabolic and Bariatric Surgery (ASMBS). Position statement. Preoperative supervised weight loss requirements. April 2016.

American Society for Metabolic and Bariatric Surgery (ASMBS). Approved procedures and devices. March 2019. Available at: <https://asmbs.org/resources/endorsed-procedures-and-devices?/resources/approved-procedures>. Accessed October 2, 2024.

Apovian CM, Shah SN, Wolfe BM, et al. Two-year outcomes of vagal nerve blocking (vBloc) for the treatment of obesity in the ReCharge Trial. *Obes Surg*. 2017 Jan;27(1):169-176.

Bai ZB, Qin YL, Deng G, et al. Bariatric embolization of the left gastric arteries for the treatment of obesity: 9-month data in 5 patients. *Obes Surg*. 2018 Apr;28(4):907-915.

Bays HE, Jones PH, Jacobson TA, et al. Lipids and bariatric procedures part 1 of 2: scientific statement from the National Lipid Association, American Society for Metabolic and Bariatric Surgery, and Obesity Medicine Association: full report. *J Clin Lipidol*. 2016a Jan-Feb;10(1):33-57.

Bays HE, Kothari SN, Azagury DE, et al. ASMBS guidelines/statements. Lipids and bariatric procedures part 2 of 2: scientific statement from the American Society for Metabolic and Bariatric Surgery (ASMBS), the National Lipid Association (NLA), and Obesity Medicine Association (OMA). *Surg Obes Relat Dis*. 2016b Mar-Apr;12(3):468-495.

Betzel B, Homan J, Aarts EO, et al. Weight reduction and improvement in diabetes by the duodenal-jejunal bypass liner: a 198 patient cohort study. *Surg Endosc*. 2017 Jul;31(7):2881-2891.

Bužga M, Švagera Z, Tomášková H, et al. Metabolic effects of sleeve gastrectomy and laparoscopic greater curvature plication: an 18-month prospective, observational, open-label study. *Obes Surg*. 2017 Dec;27(12):3258-3266.

California Technology Assessment Forum. Institute for Clinical and Economic Review 2015. Controversies in obesity management. A technology assessment. June 2015.

Camilleri M, Toouli J, Herrera MF, et al. Intra-abdominal vagal blocking (VBLOC therapy): Clinical results with a new implantable medical device. *Surgery*. 2008;143(6):723-731.

Carbajo MA, Fong-Hirales A, Luque-de-León E, et al. Weight loss and improvement of lipid profiles in morbidly obese patients after laparoscopic one-anastomosis gastric bypass: 2-year follow-up. *Surg Endosc*. 2017 Jan;31(1):416-421.

Centers for Disease Control and Prevention (CDC), National Center for Health Statistics. Clinical growth charts. Updated December 2022. Available at <https://www.cdc.gov/growthcharts/extended-bmi.htm>. Accessed October 2, 2024.

Centers for Disease Control and Prevention (CDC). National Center for Health Statistics. Obesity and Overweight. <https://www.cdc.gov/nchs/fastats/obesity-overweight.htm>. Accessed October 2, 2024.

Centers for Disease Control and Prevention (CDC). What is BMI. August 2021. Available at: https://www.cdc.gov/healthyweight/assessing/bmi/adult_bmi/index.html. Accessed October 2, 2024.

Cha R, Marescaux J, Diana M. Updates on gastric electrical stimulation to treat obesity: Systematic review and future perspectives. *World J Gastrointest Endosc*. 2014 Sep 16;6(9):419-31.

Chaar ME, Lundberg P, Stoltzfus J. Thirty-day outcomes of sleeve gastrectomy versus Roux-en-Y gastric bypass: first report based on Metabolic and Bariatric Surgery Accreditation and Quality Improvement Program database. *Surg Obes Relat Dis*. 2018 May;14(5):545-551.

Chang SH, Stoll CR, Song J, et al. The effectiveness and risks of bariatric surgery: an updated systematic review and meta-analysis, 2003-2012. *JAMA Surg*. 2014 Mar;149(3):275-87.

Christou NV, Sampalis JS, Liberman M, et al. Surgery decreases long-term mortality, morbidity, and health care use in morbidly obese patients. *Ann Surg*. 2004 Sep;240(3):416-23.

Coffin B, Maunoury V, Pattou F, et al. Impact of intragastric balloon before laparoscopic gastric bypass on patients with super obesity: a randomized multicenter study. *Obes Surg.* 2017;27(4):902-909.

Colquitt JL, Pickett K, Loveman E, et al. Surgery for weight loss in adults. *Cochrane Database Syst Rev.* 2014 Aug 8;(8):CD003641.

Cottam A, Cottam D, Medlin W, et al. A matched cohort analysis of single anastomosis loop duodenal switch versus Roux-en-Y gastric bypass with 18-month follow-up. *Surg Endosc.* 2016 Sep;30(9):3958-64.

Cottam A, Cottam D, Portenier D, et al. A matched cohort analysis of stomach intestinal pylorus saving (SIPS) surgery versus biliopancreatic diversion with duodenal switch with two-year follow-up. *Obes Surg.* 2017;27(2):454-461.

Cottam D, Roslin M, Enochs P, et al. Single Anastomosis Duodenal Switch: 1-Year Outcomes. *Obes Surg.* 2020 Apr;30(4):1506-1514.

Courcoulas A, Abu Dayyeh BK, Eaton L, et al. Intragastric balloon as an adjunct to lifestyle intervention: a randomized controlled trial. *Int J Obes.* 2017;41(3):427-433.

Desai NK, Wulkan ML, Inge TH. Update on adolescent bariatric surgery. *Endocrinol Metab Clin North Am.* 2016 Sep;45(3):667-76.

Dixon J, O'Brien P, Playfair J, et al. Adjustable gastric banding and conventional therapy for type 2 diabetes. *JAMA.* 2008;299(3):316-323.

Doležalova-Kormanova K, Buchwald JN, Skochova D, et al. Five-year outcomes: laparoscopic greater curvature plication for treatment of morbid obesity. *Obes Surg.* 2017 Nov;27(11):2818-2828.

ECRI Institute. Clinical Evidence Assessment. AspireAssist Gastric Aspiration Port (Aspire Bariatrics, Inc.) for Treating Obesity. February 2021.

ECRI Institute. Clinical Evidence Assessment. Ingestible intragastric balloons for treating obesity. April 2020.

ECRI Institute. Clinical Evidence Assessment. Orbera Intragastric Balloon (Apollo Endosurgery, Inc.) for Treating Obesity. September 2015. Updated February 2021.

ECRI Institute. Clinical Evidence Assessment Rechargeable vagal blocking system (Maestro) for treating obesity. July 2016. Updated May 2017.

ECRI Institute. Clinical Evidence Assessment. Spatz3® Adjustable Gastric Balloon (Spatz FGIA, Inc.) for Treating Obesity. January 2022.

ECRI Institute. Custom Product Brief. TransPyloric Shuttle implant (BAROnova, Inc.) for Treating Obesity. May 2019.

ECRI Institute. Health Information Assessment. Ingestible Intragastric Balloons for Treating Obesity. April 2020.

Eid GM, McCloskey CA, Eagleton JK, et al. StomaphyX vs a sham procedure for revisional surgery to reduce regained weight in Roux-en-Y gastric bypass patients: a randomized clinical trial. *JAMA Surg.* 2014 Apr;149(4):372-9.

Eisenberg D, Shikora SA, Aarts E, et al. 2022 American Society of Metabolic and Bariatric Surgery (ASMBS) and International Federation for the Surgery of Obesity and Metabolic Disorders (IFSO) Indications for Metabolic and Bariatric Surgery. *Obes Surg.* 2023 Jan;33(1):3-14. Erratum in: *Obes Surg.* 2022 Nov 29.

English WJ, DeMaria EJ, Brethauer SA, et al. American Society for Metabolic and Bariatric Surgery estimation of metabolic and bariatric procedures performed in the United States in 2016. *Surg Obes Relat Dis.* 2018 Mar;14(3):259-263.

Epstein LJ, Kristo D, Strollo PJ, et al. Adult Sleep Apnea Taskforce of the American Academy of Sleep Medicine. Clinical guideline for the evaluation, management and long-term care of obstructive sleep apnea in adults. *Journal of Clinical Sleep Medicine.* March 2009. 5(3): 263-276.

Eskandaros MS, Abbass A, Zaid MH, et al. Laparoscopic one anastomosis gastric bypass versus laparoscopic Roux-en-Y gastric bypass effects on pre-existing mild-to-moderate gastroesophageal reflux disease in patients with obesity: a randomized controlled study. *Obes Surg.* 2021 Aug 18.

Fornier PM, Ramacciotti T, Farey JE, et al. Safety and effectiveness of an endoscopically placed duodenal-jejunal bypass device (EndoBarrier®): outcomes in 114 patients. *Obes Surg.* 2017 Dec;27(12):3306-3313.

Gallas S, Fetissova S. Ghrelin, appetite and gastric electrical stimulation. *Peptides.* Volume 32, Issue 11, November 2011, Pages 2283-2289.

Garvey WT, Mechanick JI, Brett EM, et al. American Association of Clinical Endocrinologists and American College of Endocrinology comprehensive clinical practice guidelines for medical care of patients with obesity. *Endocr Pract.* 2016 Jul;22 Suppl 3:1-203.

Gersin KS, Rothstein RI, Rosenthal RJ, et al. Open-label, sham-controlled trial of an endoscopic duodenojejunal bypass liner for preoperative weight loss in bariatric surgery candidates. *Gastrointest Endosc.* 2010;71(6):976-982.

Ghiassi S, Nimeri A, Aleassa EM, et al. American Society for Metabolic and Bariatric Surgery Clinical Issues Committee. American Society for Metabolic and Bariatric Surgery position statement on one-anastomosis gastric bypass. *Surg Obes Relat Dis.* 2024 Apr;20(4):319-335.

Giordano S. Laparoscopic Roux-en-Y gastric bypass versus laparoscopic adjustable gastric banding in the super-obese: peri-operative and early outcomes. *Scandinavian Journal of Surgery.* 2015; 104(1):5-9.

Greenberg I, Perna F, Kaplan M, et al. Behavioral and psychological factors in the assessment and treatment of obesity surgery patients. *Obesity Research.* 2005;13(2),244-149.

Grubnik VV, Ospanov OB, Namaeva KA, et al. Randomized controlled trial comparing laparoscopic greater curvature plication versus laparoscopic sleeve gastrectomy. *Surg Endosc.* 2016 Jun;30(6):2186-91.

Gudur AR, Geng CX, Kshatri S, et al. Comparison of endoscopic sleeve gastroplasty versus surgical sleeve gastrectomy: a Metabolic and Bariatric Surgery Accreditation and Quality Improvement Program database analysis. *Gastrointest Endosc.* 2023 Jan;97(1):11-21.

Hardin AP, Hackell JM, American Academy of Pediatrics Committee on Practice and Ambulatory Medicine. Age limit of pediatrics. *Pediatrics.* 2017 Sep;140(3). pii: e20172151.

Hayes, Inc., Clinical Research Response. AspireAssist (Aspire Bariatrics) Aspiration Therapy for Weight Loss in Obese Individuals. Lansdale PA: Hayes, Inc., April 2021.

Hayes, Inc. Comparative Effectiveness Review. Mini-Gastric Bypass-One Anastomosis Gastric Bypass for the Treatment of Obesity: A Review of Reviews. Lansdale PA: Hayes Inc., May 2019. Updated January 2023.

Hayes, Inc. Evolving Evidence Review. Apollo ESG System (Apollo Endosurgery Inc.) for Endoscopic Sleeve Gastroplasty. Lansdale PA: Hayes, Inc., September 2024.

Hayes, Inc. Evolving Evidence Review. OverStitch Endoscopic Suturing System (Apollo Endosurgery Inc.) for Endoscopic Sleeve Gastroplasty. Lansdale PA: Hayes, Inc., May 2022.

Hayes, Inc. Hayes Medical Technology Directory. Intra-gastric balloons for treatment of obesity. Lansdale, PA: Hayes, Inc.; March 2018. Updated May 2021.

Hayes, Inc. Health Technology Brief. Single-anastomosis duodenal switch for weight loss. Landsale PA: Hayes, Inc., February 2018. Updated March 2020. Archived March 2021.

Hafezi-Nejad N, Bailey CR, Gunn AJ, et al. Weight loss after left gastric artery embolization: a systematic review and meta-analysis. *J Vasc Interv Radiol.* 2019 Oct;30(10):1593-1603.

Hedjoudje A, Abu Dayyeh BK, Cheskin LJ, et al. Efficacy and safety of endoscopic sleeve gastroplasty: a systematic review and meta-analysis. *Clin Gastroenterol Hepatol.* 2020 May;18(5):1043-1053.e4.

Hervieux E, Baud G, Dabbas M, et al. Comparative results of gastric banding in adolescents and young adults. *J Pediatr Surg.* 2016 Jul;51(7):1122-5.

Hudgel DW, Patel SR, Ahasic AM, et al; American Thoracic Society Assembly on Sleep and Respiratory Neurobiology. The role of weight management in the treatment of adult obstructive sleep apnea. An Official American Thoracic Society Clinical Practice Guideline. *Am J Respir Crit Care Med.* 2018 Sep 15;198(6):e70-e87.

Ikramuddin S, Blackstone RP, Brancatisano A, et al. Effect of reversible intermittent intra-abdominal vagal nerve blockade on morbid obesity: the ReCharge randomized clinical trial. *JAMA.* 2014; 312(9):915-922.

Inge TH, Laffel LM, Jenkins TM, et al. Comparison of surgical and medical therapy for type 2 diabetes in severely obese adolescents. *JAMA Pediatr.* 2018 May 1;172(5):452-460.

Iranmanesh P, Boudreau V, Barlow K, et al. Comparison of single- versus double-anastomosis duodenal switch: a single-center experience with 2-year follow-up. *Int J Obes (Lond).* 2021 Aug;45(8):1782-1789.

Jensen MD, Ryan DH, Apovian CM, et al; American College of Cardiology/American Heart Association Task Force on Practice Guidelines; Obesity Society. 2013 AHA/ACC/TOS guideline for the management of overweight and obesity in adults: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines and The Obesity Society. *Circulation.* 2014 Jun 24;129(25 Suppl 2):S102-38.

Jirapinyo P, de Moura DTH, Horton LC, et al. Effect of Aspiration Therapy on Obesity-Related Comorbidities: Systematic Review and Meta-Analysis. *Clin Endosc.* 2020 Nov;53(6):686-697.

Jung SH, Yoon JH, Choi HS, et al; Korean Research Group for Endoscopic Management of Metabolic Disorder and Obesity. Comparative efficacy of bariatric endoscopic procedures in the treatment of morbid obesity: a systematic review and network meta-analysis. *Endoscopy*. 2020 Nov;52(11):940-954.

Kallies K, Rogers AM; American Society for Metabolic and Bariatric Surgery Clinical Issues Committee. American Society for Metabolic and Bariatric Surgery updated statement on single-anastomosis duodenal switch. *Surg Obes Relat Dis*. 2020 Jul;16(7):825-830.

Kansou G, Lechaux D, Delarue J, et al. Laparoscopic sleeve gastrectomy versus laparoscopic mini gastric bypass: One year outcomes. *Int J Surg*. 2016 Jul 22;33(Pt A):18-22.

Kapeluto JE, Tchernof A, Masckauchan D, et al. Ten-year remission rates in insulin-treated type 2 diabetes after biliopancreatic diversion with duodenal switch. *Surg Obes Relat Dis*. 2020 Nov;16(11):1701-1712.

Katz PO, Dunbar KB, Schnoll-Sussman FH, et al. ACG Clinical Guideline for the Diagnosis and Management of Gastroesophageal Reflux Disease. *Am J Gastroenterol*. 2022 Jan 1;117(1):27-56.

Kelly AS, Barlow SE, Rao G, et al. Severe obesity in children and adolescents: identification, associated health risks, and treatment approaches: a scientific statement from the American Heart Association. *Circulation*. 2013 Oct 8;128(15):1689-712.

Khalaj A, Tasdighi E, Hosseinpanah F, et al. Two-year outcomes of sleeve gastrectomy versus gastric bypass: first report based on Tehran obesity treatment study (TOTS). *BMC Surg*. 2020 Jul 20;20(1):160.

Kim J. American Society for Metabolic and Bariatric Surgery statement on single-anastomosis duodenal switch. *Surg Obes Relat Dis*. 2016;12(5):944-945.

King K, Sudan R, Bardaro S, et al. Assessment and management of gastroesophageal reflux disease following bariatric surgery. *Surg Obes Relat Dis*. 2021 Nov;17(11):1919-1925.

Kominiarek MA, Jungheim ES, Hoeger KM, et al. American Society for Metabolic and Bariatric Surgery position statement on the impact of obesity and obesity treatment on fertility and fertility therapy. Endorsed by the American College of Obstetricians and Gynecologists and the Obesity Society. *Surg Obes Relat Dis*. 2017 May;13(5):750-757.

Lager CJ, Esfandiari NH, Subauste AR, et al. Roux-En-Y gastric bypass vs. sleeve gastrectomy: balancing the risks of surgery with the benefits of weight loss. *Obes Surg*. 2017 Jan;27(1):154-161.

Lannoo M, Dillemans B. Laparoscopy for primary and secondary bariatric procedures. *Best Pract Res Clin Gastroenterol*. 2014;28(1):159-173.

Li X, Hu X, Fu C, et al. Efficacy and safety of one anastomosis gastric bypass versus Roux-en-Y gastric bypass for obesity: a meta-analysis and systematic review. *Obes Surg*. 2023 Feb;33(2):611-622.

Lopez-Nava G, Sharaiha RZ, Vargas EJ, et al. Endoscopic sleeve gastroplasty for obesity: a multicenter study of 248 patients with 24 months follow-up. *Obes Surg*. 2017 Oct;27(10):2649-2655.

Lundell LR, Dent J, Bennett JR, et al. Endoscopic assessment of oesophagitis: clinical and functional correlates and further validation of the Los Angeles classification. *Gut*. 1999 Aug;45(2):172-80.

Maciejewski ML, Arterburn DE, Van Scoyoc L, et al. Bariatric surgery and long-term durability of weight loss. *JAMA Surg*. 2016 Nov 1;151(11):1046-1055.

Magallares A. Mental and physical health-related quality of life in obese patients before and after bariatric surgery: a meta-analysis. *Psychology, Health & Medicine*. 2015; 20(2):165-176.

Manco M, Mosca A, De Peppo F, et al. The benefit of sleeve gastrectomy in obese adolescents on nonalcoholic steatohepatitis and hepatic fibrosis. *J Pediatr*. 2017 Jan;180:31-37.e2.

Marinos G, Eliades C, Raman Muthusamy V, et al. Weight loss and improved quality of life with a nonsurgical endoscopic treatment for obesity: clinical results from a 3- and 6-month study. *Surg Obes Relat Dis*. 2014 Sep-Oct;10(5):929-34.

Mayer SB, Graybill S, Raffa SD, et al. Synopsis of the 2020 U.S. VA/DoD Clinical Practice Guideline for the Management of Adult Overweight and Obesity. *Mil Med*. 2021 Aug 28;186(9-10):884-896.

Mechanick JI, Apovian C, Brethauer S, et al. Clinical practice guidelines for the perioperative nutritional, metabolic, and nonsurgical support of patients undergoing bariatric procedures—2019 update: cosponsored by American Association of Clinical Endocrinologists/ American College of Endocrinology, the Obesity Society, and American Society for Metabolic & Bariatric Surgery, Obesity Medicine Association, and American Society of Anesthesiologists – Executive Summary. *Endocr Pract*. 2019 25(12):1346-1359.

Metabolic and Bariatric Surgery Accreditation and Quality Improvement Program (MBSAQIP). Available at: <https://www.facs.org/quality-programs/mbsaqip>. Accessed October 2, 2024.

Michalsky M, Reichard K, Inge T, et al; American Society for Metabolic and Bariatric Surgery. ASMBS pediatric committee best practice guidelines. *Surg Obes Relat Dis*. 2012 Jan-Feb;8(1):1-7.

Mingrone G, Panunzi S, Gaetano A, et al. Bariatric surgery versus conventional medical therapy for type 2 diabetes. *N Engl J Med*. 2012;366(17):1577-1585.

Mirkin K, Alli VV, Rogers AM. Revisional Bariatric Surgery. *Surg Clin North Am*. 2021 Apr;101(2):213-222.

Moore R, Rosenthal P. ASMBS proposed addendum to position statement on intragastric balloon therapy. January 2018.

Moore RL, Seger MV, Garber SM, et al. Clinical safety and effectiveness of a swallowable gas-filled intragastric balloon system for weight loss: consecutively treated patients in the initial year of U.S. commercialization. *Surg Obes Relat Dis*. 2019 Mar;15(3):417-423.

Morales-Conde S, Alarcón Del Agua I, Busetto, et al. Implanted closed-loop gastric electrical stimulation (cgles) system with sensor-based feedback safely limits weight regain at 24 months. *Obes Surg*. 2018 Jun;28(6):1766-1774.

Morton JM, Shah SN, Wolfe BM, et al. Effect of vagal nerve blockade on moderate obesity with an obesity-related comorbid condition: the ReCharge Study. *Obes Surg*. 2016 May;26(5):983-9.

Muniraj T, Day LW, Teigen LM, et al. AGA Clinical Practice Guidelines on Intragastric Balloons in the Management of Obesity. *Gastroenterology*. 2021 Apr;160(5):1799-1808.

Musella M, Susa A, Manno E, et al. Complications following the mini/one anastomosis gastric bypass (MGB/OAGB): a Multi-Institutional survey on 2678 patients with a mid-term (5 years) follow-up. *Obes Surg*. 2017 Nov;27(11):2956-2967.

National Heart, Lung and Blood Institute (NHLBI). Classification of overweight and obesity by BMI, waist circumference, and associated disease risks. 2016.

National Heart, Lung and Blood Institute (NHLBI). Managing Overweight and Obesity in Adults. Systematic Evidence Review from the Obesity Panel, 2013.

National Heart, Lung and Blood Institute (NHLBI). Think tank on enhancing obesity research at the National Heart, Lung and Blood Institute. August 2004.

National Heart Lung and Blood Institute (NHLBI). The Practical Guide Identification, Evaluation and Treatment of Overweight and Obesity in Adults. October 2020.

National Institute for Health and Care Excellence (NICE). Interventional procedures guidance [IPG518]. Implantation of a duodenal-jejunal bypass liner for managing type 2 diabetes. March 2015.

National Institute for Health and Care Excellence (NICE). Interventional procedures guidance [IPG783]. Endoscopic sleeve gastroplasty for obesity. February 2024.

National Institute for Health and Care Excellence (NICE). NICE guidelines [CG189]. Obesity: identification, assessment and management. November 2014. Updated January 2023.

National Institute for Health and Care Excellence (NICE). NICE guidelines [IPG569]. Single-anastomosis duodeno-ileal bypass with sleeve gastrectomy for treating morbid obesity. November 2016.

National Institute for Health and Care Excellence (NICE). NICE guidelines [IPG684]. Swallowable gastric balloon capsule for weight loss. November 2020.

Neto MG, Moon RC, de Quadros LG, et al. Safety and short-term effectiveness of endoscopic sleeve gastroplasty using overstitch: preliminary report from a multicenter study. *Surg Endosc*. 2020 Oct;34(10):4388-4394.

Neylan CJ, Dempsey DT, Tewksbury CM, et al. Endoscopic treatments of obesity: a comprehensive review. *Surg Obes Relat Dis*. 2016 Jun;12(5):1108-15.

Noel P, Nedelcu M, Eddbali I, et al. What are the long-term results 8 years after sleeve gastrectomy? *Surg Obes Relat Dis*. 2017 Jul;13(7):1110-1115.

Norén E, Forssell H. Aspiration therapy for obesity; a safe and effective treatment. *BMC Obes*. 2016 Dec 28;3:56.

Nunes GC, Pajecki D, de Melo ME, et al. Assessment of weight loss with the intragastric balloon in patients with different degrees of obesity. *Surg Laparosc Endosc Percutan Tech*. 2017 Aug;27(4):e83-e86.

Nyström M, Machytka E, Norén E, et al. Aspiration therapy as a tool to treat obesity: 1- to 4-year results in a 201-patient multi-center post-market European registry study. *Obes Surg*. 2018 Jul;28(7):1860-1868.

O'Brien PE, Hindle A, Brennan L, et al. Long-term outcomes after bariatric surgery: a systematic review and meta-analysis of weight loss at 10 or more years for all bariatric procedures and a single-centre review of 20-year outcomes after adjustable gastric banding. *Obes Surg*. 2019 Jan;29(1):3-14.

O'Brien PE, Sawyer SM, Laurie C, et al. Laparoscopic adjustable gastric banding in severely obese adolescents: A Randomized Trial. *JAMA*. 2010;303(6):519-526.

Olbers T, Beamish AJ, Gronowitz E, et al. Laparoscopic Roux-en-Y gastric bypass in adolescents with severe obesity (AMOS): a prospective, 5-year, Swedish nationwide study. *Lancet Diabetes Endocrinol* 2017. Online January 5, 2017.

Papasavas P, El Chaar M, Kothari SN. American Society for Metabolic and Bariatric Surgery Clinical Issues Committee. American Society for Metabolic and Bariatric Surgery position statement on vagal blocking therapy for obesity. *Surg Obes Relat Dis*. 2016 Mar-Apr;12(3):460-461.

Park JH, Kim SM. High-rate of long-term revision surgery due to weight regain after Laparoscopic Gastric Greater Curvature Plication (LGGCP). *Asian J Surg*. 2023 Feb;46(2):850-855.

Parmar CD, Gan J, Stier C, et al. One Anastomosis/Mini Gastric Bypass (OAGB-MGB) as revisional bariatric surgery after failed primary adjustable gastric band (LAGB) and sleeve gastrectomy (SG): A systematic review of 1075 patients. *Int J Surg*. 2020 Sep;81:32-38.

Parrott J, Frank L, Rabena R, et al. American Society for Metabolic and Bariatric Surgery integrated health nutritional guidelines for the surgical weight loss patient 2016 update: micronutrients. *Surg Obes Relat Dis*. 2017 May;13(5):727-741.

Pereira AM, Guimarães M, Pereira SS, et al. Single and dual anastomosis duodenal switch for obesity treatment: a single-center experience. *Surg Obes Relat Dis*. 2021 Jan;17(1):12-19.

Plamper A, Lingohr P, Nadal J, et al. Comparison of mini-gastric bypass with sleeve gastrectomy in a mainly super-obese patient group: first results. *Surg Endosc*. 2017 Mar;31(3):1156-1162.

Polega JR, Barreto TW, Kemmeter KD, et al. A matched cohort study of laparoscopic biliopancreatic diversion with duodenal switch and sleeve gastrectomy performed by one surgeon. *Surg Obes Relat Dis*. 2017 Mar;13(3):411-414.

Ponce de Leon-Ballesteros G, Romero-Velez G, Higa K, et al. Single anastomosis duodeno-ileostomy with sleeve gastrectomy/single anastomosis duodenal switch (SADI-S/SADS) IFSO position statement-Update 2023. *Obes Surg*. 2024;34(10):3639-3685.

Ponce J, Woodman G, Swain J, et al. The REDUCE pivotal trial: a prospective, randomized controlled pivotal trial of a dual intragastric balloon for the treatment of obesity. *Surg Obes Relat Dis*. 2015 Jul-Aug;11(4):874-81.

Pratt JSA, Browne A, Browne NT, et al. ASMBS pediatric metabolic and bariatric surgery guidelines, 2018. *Surg Obes Relat Dis*. 2018 Jul;14(7):882-901.

Qiu J, Lundberg PW, Javier Birriel T, et al. Revisional bariatric surgery for weight regain and refractory complications in a single MBSAQIP accredited center: what are we dealing with? *Obes Surg*. 2018 Sep;28(9):2789-2795.

Quezada N, Muñoz R, Morelli C, et al. Safety and efficacy of the endoscopic duodenal-jejunal bypass liner prototype in severe or morbidly obese subjects implanted for up to 3 years. *Surg Endosc*. 2018 Jan;32(1):260-267.

Reddy VY, Neuzil P, Musikantow D, et al. Transcatheter bariatric embolotherapy for weight reduction in obesity. *J Am Coll Cardiol*. 2020 Nov 17;76(20):2305-2317.

Risstad H, Kristinsson JA, Fagerland MW, et al. Bile acid profiles over 5 years after gastric bypass and duodenal switch: results from a randomized clinical trial. *Surg Obes Relat Dis*. 2017 Sep;13(9):1544-1553.

Rohde U, Hedbäck N, Gluud LL, et al. Effect of the EndoBarrier Gastrointestinal Liner on obesity and type 2 diabetes: a systematic review and meta-analysis. *Diabetes Obes Metab*. 2016 Mar;18(3):300-5.

Ruban A, Miras AD, Glaysher MA, et al. Duodenal-jejunal bypass liner for the management of Type 2 diabetes mellitus and obesity: A multicenter randomized controlled trial. *Ann Surg*. 2022 Mar 1;275(3):440-447.

Rubino F, Nathan DM, Eckel RH, et al. Metabolic surgery in the treatment algorithm for type 2 diabetes: a joint statement by international diabetes organizations. *Diabetes Care* 2016 Jun; 39(6): 861-877.

Saber AA, Shoar S, Almadani MW, et al. Efficacy of first-time intragastric balloon in weight loss: a systematic review and meta-analysis of randomized controlled trials. *Obes Surg*. 2017 Feb;27(2):277-287.

Salminen P, Helmiö M, Ovaska J, et al. Effect of laparoscopic sleeve gastrectomy vs laparoscopic Roux-en-Y gastric bypass on weight loss at 5 years among patients with morbid obesity: The SLEEVEPASS randomized clinical trial. *JAMA*. 2018 Jan 16;319(3):241-254.

Sanchez B, Mohr C, Morton J, et al. Comparison of totally robotic laparoscopic Roux-en-Y gastric bypass and traditional laparoscopic Roux-en-Y gastric bypass. 2005; 1(6):549-554.

Sandler BJ, Biertho L, Anvari M, et al. Totally endoscopic implant to effect a gastric bypass: 12-month safety and efficacy outcomes. *Surg Endosc*. 2018 Apr 20. [Epub ahead of print].

Sarr MG, Billington CJ, Brancatisano A, et al. The EMPOWER study: randomized, prospective, double-blind, multicenter trial of vagal blockade to induce weight loss in morbid obesity. *Obes Surg*. 2012 Nov;22(11):1771-1782.

Schauer PR, Bhatt DL, Kirwan JP, et al. Bariatric surgery versus intensive medical therapy for diabetes - 5-year outcomes. *N Engl J Med*. 2017 Feb 16;376(7):641-651.

Schouten R, Rijs CS, Bouvy ND, et al. A multicenter, randomized efficacy study of the EndoBarrier Gastrointestinal Liner for presurgical weight loss prior to bariatric surgery. *Ann Surg*. 2010 Feb;251(2):236-43.

Serrano OK, Zhang Y, Kintzer E, et al. Outcomes of bariatric surgery in the young: a single-institution experience caring for patients under 21 years old. 2016 Nov;30(11):5015-5022. Epub 2016 Mar 11.

Shah AS, Jenkins T, Gao Z, et al. Lipid changes 8 years post gastric bypass in adolescents with severe obesity (FABS-5 + study). *Int J Obes (Lond)*. 2017 Oct;41(10):1579-1584.

Sheikh L, Pearless LA, Booth MW. Laparoscopic silastic ring mini-gastric bypass (sr-mgbp): up to 11-year results from a single centre. *Obes Surg*. 2017 Sep;27(9):2229-2234.

Sheiner E, Levy A, Silverberg D, et al. Pregnancy after bariatric surgery is not associated with adverse perinatal outcome. *Am J Obstet Gynecol*. 2004 May;190(5):1335-1340.

Shikora SA, Bergenstal R, Bessler M, et al. Implantable gastric stimulation for the treatment of clinically severe obesity: results of the SHAPE trial. *Surg Obes Relat Dis*. 2009 Jan-Feb; 5(1):31-7.

Shikora SA, Toouli J, Herrera MF, et al. Intermittent vagal nerve block for improvements in obesity, cardiovascular risk factors, and glycemic control in patients with type 2 diabetes mellitus: 2-year results of the VBLOC DM² Study. *Obes Surg*. 2016 May;26(5):1021-8.

Singh S, Hourneaux de Moura DT, Khan A, et al. Safety and efficacy of endoscopic sleeve gastroplasty worldwide for treatment of obesity: a systematic review and meta-analysis. *Surg Obes Relat Dis*. 2020 Feb;16(2):340-351.

Society of American Gastrointestinal and Endoscopic Surgeons (SAGES). Guidelines for clinical application of laparoscopic bariatric surgery. March 2008.

Sogg S, Lauretti J, L West-Smith. Recommendations for the presurgical psychosocial evaluation of bariatric surgery patients *Surg Obes Relat Dis* 2016;12:731-749.

Stefanidis D, Hope WW, Kohn GP, et al. Practice/clinical guidelines by the Society of American Gastrointestinal and Endoscopic Surgeons (SAGES). Guidelines for surgical treatment of gastroesophageal reflux disease. February 2010.

Still CD, Benotti P, Wood C, et al. Outcomes of preoperative weight loss in high-risk patients undergoing gastric bypass surgery. *Arch Surg*. 2007;142(10):994-998.

Still CD, Sarwer D., Blankenship, J. The ASMBS Textbook of Bariatric Surgery, Volume 2: Integrated Health. Chapter 19. Pages 185-191.

StomaphyX - Gastric bypass revision. About weight loss surgery. April 30, 2010.

Styne DM, Arslanian SA, Connor EL, et al. Pediatric obesity-assessment, treatment, and prevention: an Endocrine Society clinical practice guideline. *J Clin Endocrinol Metab*. 2017 Mar 1;102(3):709-757.

Sudan R, Nguyen NT, Hutter MM, et al. Morbidity, mortality, and weight loss outcomes after reoperative bariatric surgery in the USA. *J Gastrointest Surg*. 2015 Jan;19(1):171-8.

Sullivan S, Kumar N, Edmundowicz SA, et al. ASGE position statement on endoscopic bariatric therapies in clinical practice. *Gastrointest Endosc*. 2015 Nov;82(5):767-72.

Sullivan S, Stein R, Jonnalagadda S, et al. Aspiration therapy leads to weight loss in obese subjects: a pilot study. *Gastroenterology*. 2013 Dec;145(6):1245-52.e1-5.

Surve A, Zaveri H, Cottam D, et al. A retrospective comparison of biliopancreatic diversion with duodenal switch with single anastomosis duodenal switch (SIPS-stomach intestinal pylorus sparing surgery) at a single institution with two year follow-up. *Surg Obes Relat Dis*. 2017 Mar;13(3):415-422.

Syed MI, Morar K, Shaikh A, et al. Gastric Artery Embolization Trial for the Lessening of Appetite Nonsurgically (GET LEAN): six-month preliminary data. *J Vasc Interv Radiol*. 2016 Oct;27(10):1502-8.

Tang Y, Tang S, Hu S. Comparative efficacy and safety of laparoscopic greater curvature plication and laparoscopic sleeve gastrectomy: a meta-analysis. *Obes Surg*. 2015 Nov;25(11):2169-75.

Thompson CC, Abu Dayyeh BK, Kushner R, et al. Percutaneous Gastrostomy Device for the Treatment of Class II and Class III Obesity: Results of a Randomized Controlled Trial. *Am J Gastroenterol*. 2017 Mar;112(3):447-457.

Thompson CC, Abu Dayyeh BK, Kushnir V, et al. Aspiration therapy for the treatment of obesity: 4-year results of a multicenter randomized controlled trial. *Surg Obes Relat Dis*. 2019 Aug;15(8):1348-1354.

Topart P, Becouarn G, Delarue J. Weight loss and nutritional outcomes 10 years after biliopancreatic diversion with duodenal switch. *Obes Surg*. 2017 Jul;27(7):1645-1650.

U.S. Preventive Services Task Force (USPSTF). Obesity in Adults: Screening and Management. June 2012.

Vilarrasa N, de Gordejuela AG, Casajoana A, et al. Endobarrier® in grade I obese patients with long-standing type 2 diabetes: role of gastrointestinal hormones in glucose metabolism. *Obes Surg*. 2017 Mar;27(3):569-577.

Wang FG, Yu ZP, Yan WM, et al. Comparison of safety and effectiveness between laparoscopic mini-gastric bypass and laparoscopic sleeve gastrectomy: A meta-analysis and systematic review. *Medicine (Baltimore)*. 2017 Dec;96(50):e8924.

Weiss CR, Abiola GO, Fischman AM, et al. Bariatric Embolization of Arteries for the Treatment of Obesity (BEAT Obesity) Trial: Results at 1 Year. *Radiology*. 2019 Jun;291(3):792-800.

Weitzner ZN, Phan J, Begashaw MM, et al. Endoscopic therapies for patients with obesity: a systematic review and meta-analysis. *Surg Endosc*. 2023;37(11):8166-8177.

Weiss CR, Akinwande O, Paudel K, et al. Clinical safety of bariatric arterial embolization: preliminary results of the BEAT Obesity Trial. *Radiology*. 2017 May;283(2):598-608.

Xie H, Doherty L, O'Boyle C. The positive impact of bariatric surgery on sleep. *Ir Med J*. 2016 Jan;109(1):328-30.

Zhao H, Jiao L. Comparative analysis for the effect of Roux-en-Y gastric bypass vs sleeve gastrectomy in patients with morbid obesity: Evidence from 11 randomized clinical trials (meta-analysis). *Int J Surg*. 2019 Dec;72:216-223.

Zou ZY, Zeng J, Ren TY, et al. Efficacy of intragastric balloons in the markers of metabolic dysfunction-associated fatty liver disease: results from meta-analyses. *J Clin Transl Hepatol*. 2021 Jun 28;9(3):353-363.

Policy History/Revision Information

Date	Summary of Changes
04/01/2026	<p>Related Policies</p> <ul style="list-style-type: none"> Updated reference link to reflect the current policy title for <i>Minimally Invasive Procedures for the Treatment of Upper Gastrointestinal Diseases (for Kentucky Only)</i>
05/01/2025	<p>Coverage Rationale</p> <ul style="list-style-type: none"> Removed language indicating: <ul style="list-style-type: none"> The following bariatric surgical procedures are proven and medically necessary for treating obesity: <ul style="list-style-type: none"> Adjustment of gastric band diameter Biliopancreatic diversion with duodenal switch Laparoscopic adjustable gastric band Laparoscopic adjustable gastric band (repair or revision) Laparoscopic adjustable gastric band removal Revisional procedure Roux-en-Y gastric bypass (RYGB) Sleeve gastrectomy Removal of adjustable gastric band and all related components which does not result in a revisional surgery is proven and medically necessary <p>Applicable Codes</p> <ul style="list-style-type: none"> Added CPT code 43999 <p>Supporting Information</p> <ul style="list-style-type: none"> Updated <i>Description of Services</i>, <i>Clinical Evidence</i>, and <i>References</i> sections to reflect the most current information Archived previous policy version CS007KY.10

Instructions for Use

This Medical Policy provides assistance in interpreting UnitedHealthcare standard benefit plans. When deciding coverage, the federal, state, or contractual requirements for benefit plan coverage must be referenced as the terms of the federal, state, or contractual requirements for benefit plan coverage may differ from the standard benefit plan. In the event of a

conflict, the federal, state, or contractual requirements for benefit plan coverage govern. Before using this policy, please check the federal, state, or contractual requirements for benefit plan coverage. UnitedHealthcare reserves the right to modify its Policies and Guidelines as necessary. This Medical Policy is provided for informational purposes. It does not constitute medical advice.

UnitedHealthcare uses InterQual® for the primary medical/surgical criteria, and the American Society of Addiction Medicine (ASAM) for substance use, in administering health benefits. If InterQual® does not have applicable criteria, UnitedHealthcare may also use UnitedHealthcare Medical Policies, Coverage Determination Guidelines, and/or Utilization Review Guidelines that have been approved by the Kentucky Department for Medicaid Services. The UnitedHealthcare Medical Policies, Coverage Determination Guidelines, and Utilization Review Guidelines are intended to be used in connection with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice.