

# Negative Pressure Wound Therapy (for Louisiana Only) **Retired April 1, 2026**

**Policy Number:** CS157LA.C

**Effective Date:** August 1, 2025 – March 31, 2026

[➔ Instructions for Use](#)

Table of Contents	Page
<a href="#">Application</a> .....	1
<a href="#">Coverage Rationale</a> .....	1
<a href="#">Definitions</a> .....	2
<a href="#">Applicable Codes</a> .....	3
<a href="#">Description of Services</a> .....	3
<a href="#">Clinical Evidence</a> .....	4
<a href="#">U.S. Food and Drug Administration</a> .....	12
<a href="#">References</a> .....	12
<a href="#">Policy History/Revision Information</a> .....	15
<a href="#">Instructions for Use</a> .....	15

## Application

This Medical Policy only applies to the state of Louisiana.

## Coverage Rationale

### Notes:

- The proven and medically necessary coverage statements in this policy apply to the use of negative pressure wound therapy (NPWT) in the outpatient setting.
- The unproven and not medically necessary coverage statements in this policy apply to all settings.

**NPWT, in an outpatient setting or upon discharge from an inpatient setting, is proven and medically necessary for treating individuals who have undergone a complete wound therapy program and meet indication-specific criteria as noted below.**

A complete wound therapy program, meeting the following criteria, must have been tried or considered and ruled out prior to initiation of NPWT:

- Documentation of evaluation, care, and wound measurements; and
- Application of dressings to maintain a moist wound environment; and
- Debridement of necrotic tissue, if present; and
- Evaluation of and provision for adequate nutritional status; and
- Documentation, by provider, of indication for NPWT; and
- Documentation, that open wound has not responded to conventional treatment after 30 days

### Indications

- Pressure ulcer ([Stage III or IV](#)) with documentation of the following:
  - [Complete wound therapy program](#), as outlined above; and
  - Appropriate turning and positioning; and
  - Use of a pressure-reducing support surface; and

- Moisture and incontinence management
- Neuropathic ulcer (e.g., diabetic ulcer) with documentation of the following:
  - [Complete wound therapy program](#), as outlined above; and
  - Comprehensive diabetic management program; and
  - Reduction in pressure on ulcer
- Venous insufficiency ulcer with documentation of the following:
  - [Complete wound therapy program](#), as outlined above; and
  - Compression bandages and/or garments have been used consistently, for at least 30 days; and
  - Leg elevation and ambulation
- Open surgical wound with documentation of the following:
  - Post-operative dehiscence (separation of a previously closed surgical incision) with documentation of a [complete wound therapy program](#), as outlined above; or
  - Open, non-healing amputation site in diabetics; or
  - Post-sternotomy infection (mediastinitis); or
  - Delayed healing or non-healing of skin graft is likely due to irregularly contoured or inadequate blood flow of the graft bed
- High-risk open fracture ([Gustilo Grade III](#))

**The following indications and devices are unproven and not medically necessary due to insufficient evidence of efficacy:**

- NPWT for treating **all** other indications including but not limited to:
  - Closed surgical incisions
  - Pilonidal disease
- Disposable/single-use NPWT systems
- NPWT systems with instillation of wound solutions

## Contraindications to NPWT

- Active bleeding or exposed vasculature in wound
- Anticoagulation therapy
- Eschar or necrotic tissue present in wound
- Exposed bone, nerves or organs in vicinity of wound
- Malignancy present in wound
- Uncontrolled soft tissue infection or untreated osteomyelitis within vicinity of wound
- Presence of an open fistula to body organs or cavities within vicinity of wound

NPWT should be discontinued when **any** of the following criteria are present:

- Documentation of weekly assessment of the wound's dimensions and characteristics by the provider indicate failure of progressive wound healing [i.e., wound is not diminishing in size (either surface area or depth) within 30 days]; or
- The depth of the wound is 1 mm or less; or
- Uniform granulation tissue has been obtained

## Definitions

**Gustilo Grade III Fracture:** An open fracture with extensive soft-tissue damage or an open segmental fracture.

- **IIIA:** Adequate soft-tissue coverage of a fractured bone despite extensive soft-tissue laceration or flaps, or high-energy trauma regardless of wound size.
- **IIIB:** Extensive soft-tissue injury loss with periosteal stripping and bone exposure; associated with massive contamination; often requires soft-tissue coverage (i.e., flap).
- **IIIC:** Arterial injury requiring repair (Gustilo and Anderson, 1976; Gustilo et al., 1984).

**National Pressure Injury Advisory Panel (NPIAP) Staging System (NPIAP, 2019):**

- **Stage III:** Characterized by full-thickness loss of skin, in which fat is visible in the ulcer and granulation tissue and epibole (rolled wound edges) are often present. Slough and/or eschar may be visible. The depth of tissue damage varies by anatomical location; areas of significant adiposity can develop deep wounds. Undermining and tunneling may occur. Fascia, muscle, tendon, ligament, cartilage and/or bone are not exposed. If slough or eschar obscures the extent of tissue loss this is an unstageable pressure injury.
- **Stage IV:** Characterized by full-thickness skin and tissue loss with exposed or directly palpable fascia, muscle, tendon, ligament, cartilage or bone in the ulcer. Slough and/or eschar may be visible. Epibole (rolled edges),

undermining and/or tunneling often occur. Depth varies by anatomical location. If slough or eschar obscures the extent of tissue loss this is an unstageable pressure injury.

- **Unstageable pressure injury:** Characterized by obscured full-thickness skin and tissue loss, in which the extent of tissue damage within the ulcer cannot be confirmed because it is obscured by slough or eschar. If slough or eschar is removed, a Stage III or Stage IV pressure injury will be revealed. Stable eschar (i.e., dry, adherent, intact without erythema or fluctuance) on the heel or ischemic limb should not be softened or removed.

## Applicable Codes

The following list(s) of procedure and/or diagnosis codes is provided for reference purposes only and may not be all inclusive. Listing of a code in this policy does not imply that the service described by the code is a covered or non-covered health service. Benefit coverage for health services is determined by federal, state, or contractual requirements and applicable laws that may require coverage for a specific service. The inclusion of a code does not imply any right to reimbursement or guarantee claim payment. Other Policies and Guidelines may apply.

**Coding Clarification:** Suction pumps and dressing codes (HCPCS codes K0743–K0746) apply to devices other than negative pressure wound therapy.

CPT Code	Description
*97605	Negative pressure wound therapy (e.g., vacuum assisted drainage collection), utilizing durable medical equipment (DME), including topical application(s), wound assessment, and instruction(s) for ongoing care, per session; total wound(s) surface area less than or equal to 50 square centimeters
*97606	Negative pressure wound therapy (e.g., vacuum assisted drainage collection), utilizing durable medical equipment (DME), including topical application(s), wound assessment, and instruction(s) for ongoing care, per session; total wound(s) surface area greater than 50 square centimeters
*97607	Negative pressure wound therapy, (e.g., vacuum assisted drainage collection), utilizing disposable, non-durable medical equipment including provision of exudate management collection system, topical application(s), wound assessment, and instructions for ongoing care, per session; total wound(s) surface area less than or equal to 50 square centimeters
*97608	Negative pressure wound therapy, (e.g., vacuum assisted drainage collection), utilizing disposable, non-durable medical equipment including provision of exudate management collection system, topical application(s), wound assessment, and instructions for ongoing care, per session; total wound(s) surface area greater than 50 square centimeters

*CPT® is a registered trademark of the American Medical Association*

HCPCS Code	Description
A6550	Wound care set, for negative pressure wound therapy electrical pump, includes all supplies and accessories
*A9272	Wound suction, disposable, includes dressing, all accessories and components, any type, each
E2402	Negative pressure wound therapy electrical pump, stationary or portable

Codes labeled with an asterisk (\*) are not on the State of Louisiana Medicaid Fee Schedule and therefore may not be covered by the State of Louisiana Medicaid Program.

## Description of Services

Negative pressure wound therapy (NPWT), also referred to as vacuum-assisted wound closure, is a therapeutic dressing system in which negative pressure is continuously or intermittently applied to the surface of a wound. The system includes dressings, a suction pump, tubing and a collection chamber. The wound and porous dressing are sealed with an occlusive dressing and connected to the drainage tubing connected to a suction pump that delivers subatmospheric pressure. NPWT is intended to assist wound healing by the removal of exudate or debris, reduction of bacterial contamination, increase in local blood flow, reduction of local edema, approximation of the wound edges, and the production of granulation tissue. NPWT is intended as an adjunct treatment for wounds that do not respond to conventional treatment such as debridement, pressure relief, and infection control.

## Clinical Evidence

Despite a lack of strong evidence to support its use, NPWT has gained wide acceptance for a variety of wounds.

Hurd et al. (2021) published consensus-based recommendations on the use of NPWT in acute and chronic wound management. The document presents a clinical decision-making tool for initiating NPWT and the optimal system to be utilized based on a number of factors.

Kirsner et al. (2019) conducted a multicenter, randomized, comparative-efficacy study in patients with venous leg ulcers or diabetic foot ulcers. The study compared the change in target ulcer dimensions (area, depth, and volume) using single-use NPWT versus traditional NPWT (t-NPWT) over a 12-week treatment period or up to confirmed healing. Randomized by wound type and size, 164 patients with non-infected diabetic foot ulcers and venous leg ulcers were included. The intention to treat population was composed of 161 patients (101 with venous leg ulcers, 60 with diabetic foot ulcers) and 115 patients completed follow-up (64 in the s-NPWT group and 51 in the t-NPWT group) (PP population). Primary endpoint analyses on wound area reduction demonstrated statistically significant reduction in favor of s-NPWT ( $p = 0.003$ ) for the PP population and for the ITT population ( $p < 0.001$ ). Changes in wound depth ( $p = 0.018$ ) and volume ( $p = 0.013$ ) were also better with s-NPWT. Faster wound closure was observed in the ITT population. Wound closure occurred in 45% of patients in the s-NPWT group vs. 22.2% of patients in the t-NPWT group ( $p = 0.002$ ). Median estimate of the time to wound closure was 77 days for s-NPWT. No estimate could be provided for t-NPWT. Device-related adverse events were more frequent in the t-NPWT group than in the s-NPWT group. The s-NPWT system met noninferiority and achieved statistical superiority versus t-NPWT in terms of wound progression toward healing over the treatment period. This study is limited by small numbers and short-term follow-up. Also, the study was designed to compare two types of NPWT systems, not to compare NPWT against standard of care or standard dressings.

Anghel and Kim (2016) conducted a comprehensive literature review of NPWT versus standard care for various wound types. A total of 26 publications were included. The authors tabulated and discussed the level of evidence, wound type studied, reported outcomes and impact and key findings. The authors concluded that NPWT has a role in managing chronic, complex and infected wounds. Randomized controlled trials (RCTs) validating superiority of NPWT in certain patient populations are cited. They also noted that more robust, randomized, prospective studies are needed to support its expanding use.

### Pressure Ulcers

An updated Cochrane systematic review first published 2015 evaluated the effectiveness of NPWT for treating adults with pressure ulcers in any care setting. Eight RCTs with a total of 327 individuals. Five studies compared NPWT with dressings. The evidence for this outcome was assessed as very low certainty (RR 1.25, 95% CI 0.64 to 2.44). One study compared NPWT with a series of gel treatments, but this study provided no usable data. Another study compared NPWT with moist wound healing which did not report primary outcome data. The last study compared NPWT combined with internet-plus home care with standard care and no primary outcome data were reported. All studies demonstrated low certainty of evidence. The authors concluded the efficacy, safety, and acceptability of NPWT in treating pressure ulcers compared to usual care are uncertain due to the lack of key data on complete wound healing, adverse events, and time to complete healing. Further high-quality research is still needed to help decision-makers judge the value of NPWT in the treatment of pressure ulcers. Limitations of study evidence include small sample sizes, poorly reported or unclear duration, limited data and inconclusive adverse effects (Shi et al., 2023).

Sahin et al. (2021) conducted an RCT on the effectiveness of NPWT compared to wet-to-dry dressing in pressure injuries (PIs). Thirty patients with stage 3 and 4 pressure injuries were divided into two groups: NPWT group and the wet-to-dry-dressing group. Following three rounds of treatment in all patients, data were collected using a three-dimensional wound measurement (3DWM) device, pressure ulcer scale for healing (PUSH) tool and patient identification form. The data revealed significant granulation tissue formation in the experimental group ( $p < .05$ ) with significant wound shrinkage ( $p < .05$ ) and a decrease in PUSH tool scores ( $p < .05$ ). Patient wounds were assessed with the PUSH Tool and 3DWM system which showed device measurements were correlated with PUSH tool findings ( $p < .05$ ) a significant correlation noted between device-measured granulation findings and PUSH tool score results of the experimental group's third measurements ( $p < .05$ ). The authors concluded NPWT is an effective treatment measure for pressure injuries. Limitations include small sample size and short-term follow-up. The authors recommend future studies with larger sample sizes that monitor treatment efficacy until the wound has completely healed.

Vig et al. (2011) published evidence-based recommendations for the use of NPWT in chronic wounds. Based on a systematic review of the literature, the international panel of experts recommended the following regarding pressure ulcers:

- NPWT may be used until surgical closure is possible/desirable.
- Alternatively, NPWT should be considered to achieve closure by secondary intention.
- NPWT should be used to reduce wound dimensions.
- NPWT should be used to improve the quality of the wound bed.

## Neuropathic Ulcers

Wu et al. (2023) conducted an RCT to compare the efficacy of NPWT and alginate dressings on wound bed preparation prior to split thickness skin graft (STSG) surgery for patients with chronic diabetic foot ulcers (DFUs). Patients were randomly assigned into two groups NPWT (with vacuum-assisted closure, n = 50) or the control group (with alginates dressings, n = 50). The results demonstrated the patients in the NPWT group had less time to STSG surgery than the control group. The patients in the NPWT group had prominently increased survival rates of skin graft, increased wound blood perfusion, and decreased (neutrophil extracellular traps) NET formation, and polarization of macrophages. The authors concluded NPWT is superior to conventional moist dressings in wound preparation prior to STSG surgery for patients with chronic DFUs. Limitations of the study include not a multi-center trial, small sample size and no long-term follow-up.

A systematic review and meta-analysis performed by Wang et al. (2022) compared the efficacy and safety of the NPWT with moist wound care (MWC) in the treatment of diabetic foot ulcers (DFUs). A total of 10 RCTs (619 patients in NPWT group and 625 in MWC group) were included in the review, and 8 trials were included for qualitative and quantitative syntheses. The clinical outcomes analyzed healing results, amputation or resection incidence and risk of adverse events. The data demonstrated significantly lower risk of non-closure of the wound [risk ratio (RR) = 0.74, 95% confidence interval {CI}: 0.63–0.87; p = .001], lower average wound area (standard mean difference = -0.80, 95% CI: -1.54 to -0.06; p = .034), more wound area decrease (standard mean difference = 0.81, 95% CI: 0.36–1.26; p = .001), an increase in the appearance rate of granulation tissue (RR = 1.61, 95% CI: 1.07–2.41; P=0.021), and lower risk of amputation or resection (RR = 0.70, 95% CI: 0.50–0.99; p = .045), were shown for the NPWT group when compared to MWC group. There was no statistically significant distinction found for the disappearance rate of wound discharge at 8 weeks, the rate of blood culture positivity, pain scoring, and the overall prevalence of adverse events between the groups (p = .05). The authors concluded that NPWT could accelerate the wound healing process and decrease the risk of post-treatment amputation or resection, without any additional frequency of adverse events when compared with MWC, in patients with DFUs. Limitations include the addition of RCTs with relatively low-quality and small sample sizes to the study. The authors recommend more high-quality RCTs are needed to identify the treatment efficacy of NPWT compared with MWC.

A Hayes report on the use of NPWT in the home setting as an adjunct treatment for chronic wounds in adults reported on three studies for chronic diabetic foot ulcers. The studies found benefit with NPWT for complete wound healing or wound closure. An additional study found time to wound closure was shorter for patients receiving NPWT (Hayes, 2016; updated 2021).

Rys et al. (2020) conducted a systematic review and meta-analysis of 16 observational studies (n = 1882 managed with NPWT) evaluating the efficacy and safety of NPWT in patients with diabetic foot ulcers. In the NPWT-treated patients, ulcers were larger (average size range 6.6-27.9 cm<sup>2</sup>), as compared with controls (≤ 3 cm<sup>2</sup>). The pooled results showed healing and major amputation in 51% and 5% of NPWT patients, respectively. A meta-analysis of four comparative studies revealed lower risk of major amputation in NPWT-treated patients. The pooled results for healing rate and risk of any amputation were inconclusive due to large heterogeneity between studies. These results support earlier RCT data that NPWT is an effective and safe adjunct therapy in the management of diabetic foot ulcers.

An updated Cochrane systematic review assessed the effects of NPWT compared with standard care or other adjuvant therapies in the healing of diabetic foot wounds. Eleven RCTs (n = 972) were included. The authors found low-certainty evidence to suggest that NPWT may be effective in healing postoperative foot wounds and ulcers of the foot in people with diabetes compared with wound dressing, in terms of the proportion of wound healed and time to healing. For the comparisons of different pressures of NPWT for treating foot ulcers in people with diabetes, it is uncertain whether there is a difference in the number of wounds closed or covered with surgery, and adverse events. None of the included studies provided evidence on time to closure or coverage surgery or health-related quality of life (Liu et al., 2018).

Liu et al. (2017) performed a systematic review and meta-analysis to assess the safety and efficacy of NPWT in the treatment of diabetic foot ulcers. A total of eleven RCTs (n = 1044) were included. Compared with standard dressing changes, NPWT had a higher rate of complete healing, shorter healing time, greater reduction in ulcer area and depth and fewer amputations.

Anghel and Kim (2016) conducted a comprehensive literature review of NPWT versus standard care for various wound types. Seven of the studies investigated complicated wounds in diabetic patients, either following amputations, significant surgical intervention or chronic stable ulcers. The consensus was that NPWT is safe, effective and reduces operative interventions for complicated wounds in diabetic patients.

Zhang et al. (2014) conducted a meta-analysis to evaluate the safety and effectiveness of NPWT for diabetic foot ulcers. Eight RCTs (n = 669) were included. Compared with non-NPWT treatments, NPWT resulted in a significantly higher proportion of healed ulcers, more reduction of ulcer area, fewer major amputations and shorter time to wound healing.

Vig et al. (2011) published evidence-based recommendations for the use of NPWT in chronic wounds. Based on a systematic review of the literature, the international panel of experts recommended the following regarding diabetic foot ulcers:

- NPWT must be considered as an advanced wound care therapy for postoperative grade 2 and 3 diabetic feet without ischemia.
- NPWT must be considered to achieve healing by secondary intention.
- Alternatively, NPWT should be stopped when wound has progressed suitably to be closed by surgical means.
- NPWT should be considered in an attempt to prevent amputation or re-amputation.

## Venous Insufficiency Ulcers

A Hayes report on the use of NPWT in the home setting as an adjunct treatment for chronic wounds in adults found one study demonstrating that venous ulcers were more likely to heal among patients who received NPWT than among those who did not (Hayes, 2016; updated 2021).

Vig et al. (2011) published evidence-based recommendations for the use of NPWT in chronic wounds. Based on a systematic review of the literature, the international panel of experts recommended the following regarding venous leg ulcers:

- If first-line therapy (compression) is not efficacious, NPWT should be considered to prepare the wound for surgical closure as part of a clinical pathway.

## Open Surgical Wounds

Moreno et al. (2025) conducted an RCT between March 2019 and March 2021 to assess the efficacy of NPWT in the prevention of surgical site occurrences in open abdominal surgery. Two hundred seventy-five participants were randomized to two groups, 147 in the NPWT group and 128 in the conventional dressing group (control group). The aim of the study was to evaluate the benefit of NPWT in the surgical site infection rates and other surgical site occurrences in both groups at a 30-day follow-up. The results demonstrated 31 (11.3%) surgical site infections, and 71 (25.8%) other surgical site occurrences were observed, being significantly lower in the NPWT group (odds ratio 0.31, 95% confidence interval 0.14-0.71; p = .005) and (odds ratio 0.51, 95% confidence interval 0.29 -0.90; p = .02), respectively. Absolute risk reduction was 13% for surgical site infection and 12% for other surgical site occurrences. Number needed to treat nine (95% confidence interval 5-29) for surgical site infection and number needed to treat of eight (95% confidence interval 5-51) for other surgical site occurrences. Median hospital stay was three days lower in the negative-pressure wound therapy group than in the control group (9 vs 12 days; p = .03). No severe adverse events attributable to the negative-pressure wound therapy dressing were reported. The authors concluded NPWT decreases the risk of surgical site occurrences and surgical site infection after laparotomy, so that its use should be considered in patients with risk factors. Limitations of the study include lack of blinding, detection bias and single-center sample.

The Cochrane Library (Cheng et al. 2022) performed a systematic review that compared NPWT with any other type of temporary abdominal closure in non-trauma patients with open abdomen in any care setting. A secondary comparison was performed for different types of NPWT systems. Based on the available trial data where only two studies met inclusion criteria, the authors concluded there is uncertainty whether NPWT has any benefit in terms of primary fascial closure of the abdomen, adverse events (fistulae formation), all-cause mortality or length of hospital stay. Given this uncertainty of evidence in this review clinicians, patients, and other stakeholders may need to take into account other considerations when making treatment decisions.

A Hayes evidence report summarized the publications related to the use of NPWT in the outpatient setting for treatment of chronic wounds. The report described the potential benefits of NPWT which included symptom management, reduced frequency of dressing changes, and faster healing times. However, there may be potential harm associated with NPWT such as pain, retention of foreign bodies from the dressing, bleeding, infection, death from infection and complications stemming from loss of electricity (Hayes, 2022).

Anghel and Kim (2016) conducted a comprehensive literature review of NPWT versus standard care for various wound types. Four studies evaluated the use of NPWT for split thickness skin graft retention, with 3 specifically investigating the use in acute injury or burn patients. All found that NPWT resulted in better outcomes than standard dressing. The use of NPWT after skin-grafted free muscle flaps resulted in reduced inflammatory response and edema formation.

A Cochrane systematic review concluded that there is some evidence that NPWT may reduce time to healing following a punch skin graft transplant (Dumville et al., 2015).

Azzopardi et al. (2013) systematically reviewed the evidence for the perioperative application of NPWT to split-thickness skin grafts. Thirty-eight studies were included. The authors reported two complementary trends explaining the mechanisms whereby grafts benefit from NPWT: active stimulation of epithelial mitosis and prevention of complications. NPWT also promotes microcirculatory flow and stimulates angiogenesis. This study concluded that NPWT increases quantity and quality of graft take compared to traditional bolster dressings. The advantages are increased in irregularly contoured, technically difficult wounds and suboptimal recipient wound beds.

Pan et al. (2013) performed a systematic review and meta-analysis to evaluate the efficacy of NPWT compared to conventional therapy in the treatment of post-sternotomy infections. Twelve cohort studies (n = 873) were included. The authors reported that wound closure was obtained more frequently in the NPWT group when compared to conventional therapy. NPWT was associated with a significant reduction in length of stay compared with standard of care.

## Open Fractures

Alves et al. (2024) conducted a systematic review and meta-analysis to assess the effectiveness of negative wound pressure therapy (NPWT) compared with conventional wound dressing (CWD), also described as standard wound coverage without sub-atmospheric pressure, in the management of Gustilo 3 lower limb fractures with a focus on overall rates, superficial infection, and deep infection rates. A systematic review of medical research databases was conducted in accordance with PRISMA guidelines. Studies comparing NPWT with CWD for Gustilo 3 fractures were included. Data extraction and quality assessment were performed. Treatment with CWD was associated with higher rates of overall infection [pooled risk ratio (RR): 0.33; 95% confidence interval {CI}: 0.14–0.51] and pooled risk difference (RD: 0.27; 95% CI: 0.15–0.38), superficial infection (pooled RR: 0.35; 95% CI: 0.04–0.66), and deep infection (pooled RR: 0.20; 95% CI: 0.02–0.38) compared with NPWT treatment. Overall infection rate remained higher in the CWD group after analyzing only open tibia fractures (pooled RR: 0.35; 95% CI: 0.21–0.48). Nonunion rate was higher in the CWD group (pooled RR: 0.30; 95% CI: 0.00–0.59). Flap failure rate was similar in both groups (pooled RR: 0.09; 95% CI: –0.05 to 0.23). The authors concluded that NPWT appears to be a reasonable option for wound management in Gustilo 3 lower limb fractures in terms of infection rates. This systematic review and meta-analysis have limitations. First, there is a paucity of high-quality randomized controlled trials (RCTs) specifically addressing Gustilo 3 lower limb fractures. Second, treatment protocols varied across studies, and most did not provide full details of their wound care protocol. Additionally, details of bone fixation and soft-tissue coverage were often limited and lacked specificity. Final reported outcomes were mainly based on subjective and nonquantifiable measures. These variations in study design and reporting standards highlight the complexity and heterogeneity of Gustilo 3 fractures. Furthermore, the present meta-analysis may not capture all the relevant clinical nuances and patient specific factors influencing treatment decisions. The findings of this study need to be validated by well-designed studies.

Over a five-year period Costa et al. (2024) conducted the Wound Healing in Surgery for Trauma (WHIST) trial reporting the outcomes of patients with complex fracture of the lower limb. The trial compared NPWT dressings with standard dressings applied at the end of the first operation for patients undergoing internal fixation of the fracture. Complex fractures included periarticular fractures and open fractures when the wound could be closed primarily at the end of the first debridement. A total of 1548 patients aged sixteen or older completed the initial follow-up, six months after injury. Patients reported their Disability Rating Index (DRI) and health-related quality of life, chronic pain scores and neuropathic pain scores annually along with any complications. The results demonstrated there was no evidence of a difference in patient-reported disability between the two groups at five years (NPWT group mean DRI 30.0 (SD 26.5), standard dressing group mean DRI 31.5 (SD 28.8), adjusted difference -0.86 (95% CI -4.14 to 2.40; p = 0.609). Also, there was no difference in complication rates. The study had limited follow-up, where only 66% of the 1548 patients provided data during the 5 years, thus the study did not have the same statistical power to detect subtle differences in outcomes. The authors concluded no evidence of a difference in disability ratings between NPWT compared with standard wound dressings in the five years following surgical treatment of a complex fracture of the lower limb. Patients in both groups reported high levels of persistent disability and reduced quality of life.

Haidari et al. (2021) performed a systematic review to assess the role of NPWT in the management of soft tissue defects in patients with fracture-related infection (FRI). Eight articles that focused on the infection recurrence which ranged from 2.8% to 34.9% were included in the review. Six studies reported on wound healing time, varying from one to seven weeks.

Four studies took repeated microbial swabs during subsequent NPWT dressing changes. One study reported newly detected pathogens in 23% of the included patients and three studies did not find new pathogens. The authors concluded there is no clear evidence to support the use of NPWT for FRI as definitive treatment. However, the authors did recommend early soft tissue coverage with a local or free flap, stating that NPWT may be safe for a few days as temporary soft tissue coverage until further definitive treatment could be performed. The authors note that due to the lack of uniformity in the studies, caution should be used when drawing conclusions, and further comparative studies are needed. Limitations include the limited number of high-quality studies on FRI treatment with NPWT, lack of uniformity between the studies, and small sample sizes. In the multicenter, randomized WOLFF trial, 460 patients with a severe open fracture of the lower limb were treated with NPWT (n = 226) or standard dressings without NPWT (n = 234). At 12 months, deep surgical site infection (SSI) rates, self-rated disability and quality of life were similar in both groups (Costa et al., 2018).

In a Cochrane systematic review, Ihezor-Ejiofor et al. (2018) evaluated the effectiveness of NPWT for treating open traumatic wounds. Seven RCTs (n = 1377) were included. Study sample sizes ranged from 40 to 586 participants. Four studies compared NPWT with standard care for open fracture wounds. The authors concluded that there is moderate-certainty evidence for no clear difference between NPWT and standard care on the proportion of wounds healed at six weeks for open fracture wounds. It is uncertain whether there is a difference in risk of wound infection, adverse events, time to closure or coverage surgery, pain or health-related quality of life between NPWT and standard care for any type of open traumatic wound.

Virani et al. (2016) conducted a prospective randomized trial to evaluate the role of NPWT on the incidence of deep infections/osteomyelitis after open tibial fractures. Ninety-three adults with open tibial fractures were randomized into two groups: NPWT and daily cleaning, dressing and debridement. After 23 weeks, the rate of infection was significantly lower (4.6%) in the NPWT group compared to the control group (22%). NPWT was also associated with less bacterial colonization (6.9% vs. 34%) of wounds compared to the control group. Five patients (25%) from the control group developed osteomyelitis. The authors concluded that NPWT is beneficial for preventing the incidence of both acute infections and osteomyelitis in open fractures. The time required for the wounds to be ready for closure or coverage was similar in both groups (8.3 days vs. 9.8 days).

Tansarli et al. (2014) performed a meta-analysis of four RCTs (n = 367) evaluating the incidence of SSIs in patients with open wounds following fracture stabilization. Infection rates in patients whose wounds were treated with vacuum-assisted closure (n = 196) were reduced by 53% when compared to nonvacuum closure (n = 171).

## Closed Surgical Incisions

There is insufficient clinical evidence demonstrating the safety and/or efficacy of NPWT systems, including disposable systems, for treating closed surgical incisions. Studies to date have been too small or at risk of bias to support routine use. Further results from prospective, high quality studies are needed to determine which patient population would benefit from the use of these devices.

Mantyh et al. (2024) conducted a systematic review and meta-analysis to assess the impact of incision negative pressure therapy (ciNPT) on postsurgical outcomes in patients undergoing open abdominal surgeries. The literature search identified 22 studies from nine countries that met inclusion criteria. The results showed significant reductions in relative risk (RR) of SSC (RR: 0.568, p = .003), surgical site infection (SSI) (RR: 0.512, p < .001), superficial SSI (RR: 0.373, p < .001), deep SSI (RR: 0.368, p = .033), and dehiscence (RR: 0.581, p = .042) were associated with ciNPT use. ciNPT use was also associated with a reduced risk of readmission and a 2.6-day reduction in hospital length of stay (p < .001). The authors concluded ciNPT was associated with an overall reduction in surgical site complications (SSCs), hospital LOS and readmissions. Although the findings of this meta-analysis indicate that ciNPT is beneficial and could potentially result in cost savings for patients undergoing abdominal surgery, additional research is needed to determine optimal use of ciNPT in this population and provide a robust assessment of the cost-effectiveness of the therapy. Limitations of the study include study design, potential for selection bias, varied ciNPT training and differences in outcome measures reporting.

In an ECRI clinical evidence assessment on NPWT for preventing surgical site infection after cesarean section in women with obesity, it was concluded that based on evidence reported in two systematic reviews (SRs) with meta-analysis, NPWT is safe and reduces surgical site infections when compared with conventional dressings. One reported lower rates of dehiscence, seroma, hematoma and bleeding, and the other showed similar complication rates when NPWT is compared to conventional dressings when all wound complications were grouped. One SR reported similar rates of reoperation and readmission.

A Hayes Evolving Evidence Review found minimal support for the use of PICO single-use NPWT systems for wound care in women with clinical obesity following a cesarean birth. A limited amount of clinical evidence and one clinical practice guideline indicate some support for the use of single-use NPWT in this population; however, study conclusions were inconsistent (Hayes, 2021d; updated 2024).

A Hayes report on the prophylactic use of NPWT following elective open abdominal surgeries concluded that the current body of overall low-quality evidence suggests that there may be a benefit to NPWT over standard sterile dressing driven by a lower rate of superficial infections; however, recent RCT evidence has not confirmed these findings and uncertainty remains. Significant heterogeneity exists between patient populations, underlying reason for abdominal surgery, and treatment characteristics within the included body of evidence making it difficult to discern which patients might most benefit from this form of prophylaxis (Hayes, 2021e; updated 2024).

An updated Cochrane review (Norman et al., 2020), assessed the effects of NPWT for preventing SSI in wounds healing through primary closure. This update added 15 new RCTs, resulting in a total of 44 RCTs (n = 7447). Studies evaluated NPWT in the context of a wide range of surgeries including orthopedic, obstetric, vascular and general procedures. Most studies had unclear or high risk of bias for at least one key domain. The review concluded that there were no clear difference in number of deaths or wound dehiscence between people treated with NPWT and standard dressings (low-certainty evidence). There were also no clear differences in secondary outcomes where all evidence was low or very low-certainty. In cesarean section in obese women and surgery for lower limb fracture, there was probably little difference in quality of life scores (moderate-certainty evidence). Most evidence on pain was very low-certainty, but there is probably no difference in pain between NPWT and standard dressings after surgery for lower limb fracture (moderate-certainty evidence). In a fourth update of this review (Norman et al. 2022) added 18 new RCTs to the study for a total of 62 RCTs of 13,340 participants. Once more, the studies evaluated a wide range of surgeries, including orthopedic, obstetric, vascular and general procedures. All studies compared NPWT with standard dressings. Most studies had unclear or high risk of bias for at least one key domain. The results concluded there is no difference in wound dehiscence (moderate-certainty evidence) for people with primary closure of their surgical wound treated prophylactically with NPWT following surgery compared with people treated with standard dressings; however, NPWT probably results in fewer surgical site infections (SSIs) than standard dressings. There may be reduced risk of death after surgery for those who are treated with NPWT compared with standard dressings (low-certainty evidence). Also, there may be more instances of skin blistering when comparing NPWT with standard dressing treatment (low-certainty evidence). There is no clear difference in secondary outcomes where most evidence is low or very low certainty. Decisions about NPWT use should consider surgical indication, setting and evidence for all outcomes.

Two separate RCTs (Tuuli et al., 2020; Hussamy et al., 2019) found that prophylactic NPWT did not significantly reduce the risk of postoperative wound morbidity when compared with standard wound dressing in obese women undergoing cesarean delivery. The Tuuli study (included in the 2022 Cochrane review above) was terminated after 1624 of 2850 participants were recruited when a planned interim analysis showed increased adverse events in the NPWT group and futility for the primary outcome of reduced SSIs.

In the WHIST multicenter RCT, Costa et al. (2020) assessed outcomes of incisional NPWT (n = 785) versus standard wound dressing (n = 763) on deep SSI after surgery for lower limb fractures associated with major trauma. At 30 days, deep SSI occurred in 5.84% (45 of 770 patients) of the incisional NPWT group and in 6.68% (50 of 749 patients) of the standard wound dressing group. At 90 days, there was no significant difference in the infection rates between groups. Additionally, there were no significant differences, at any time point, for the secondary outcomes of patient-reported disability, health-related quality of life, surgical scar assessment and chronic pain. The authors concluded that the results do not support the use of incisional NPWT for this indication, although the event rate at 30 days was lower than expected.

Singh et al. (2019) performed a meta-analysis of 30 studies evaluating single-use NPWT systems for treating closed wounds. RCTs and observational studies were assessed across specialties including cardiothoracic, lower extremity, colorectal/abdominal, obstetrics and vascular surgery. Results demonstrated that the Prevena system performed significantly better at reducing the incidence of SSIs in comparison to traditional and advanced wound dressings. Author noted limitations include heterogeneity of data and lack of high-quality studies for the review.

Strugala and Martin (2017) conducted a meta-analysis of 16 studies (10 RCTs and 6 observational studies) comparing prophylactic use of the PICO NPWT system with standard care. A total of 1863 patients were represented in the data. The study reported significant reduction in SSIs, wound dehiscence and hospital length of stay in patients treated with NPWT. Similar effects were seen irrespective of the kind of surgery (orthopedic, abdominal, colorectal or cesarean section). The inclusion of patients with incisions that would not be classified as “clean” is a noted limitation.

Scalise et al. (2016) performed a systematic review of studies evaluating NPWT for preventing complications of closed surgical incisions. Eighteen studies were included: 1 biomedical engineering study, 2 animal studies and 15 human studies (6 RCTs, 5 prospective cohorts, 7 retrospective analyses). Human studies investigated the outcomes of 1042 incisions on 1003 patients. The review noted a decrease in infections, hematomas and re-operation rates; however, results were inconsistent regarding wound dehiscence. Noting limited studies, the authors concluded that further study is needed to identify proper recommendations for NPWT in this patient population.

## **Pilonidal Disease**

There is insufficient clinical evidence demonstrating the safety and/or efficacy of NPWT systems, including disposable systems, for treating pilonidal disease. Further results from prospective, high quality studies are needed to determine which patient population would benefit from the use of these devices.

Ensor et al. (2024) conducted an RCT to investigate whether NPWT would reduce rates of surgical wound dehiscence (SWD) compared to conventional passive (CP) dressings for pilonidal sinus disease (PSD) excisions with off-midline primary closure. Secondary outcomes included patient quality of life and return time to normal activities. Fifty participants from four tertiary hospitals were randomized, 25 to NPWT and 25 to CP. The results demonstrated overall dehiscence rate was 42% (21/50); 12/25 (48%) for NPWT & 9/25 (36%) for CP,  $p = 0.6$ . Five deep ( $\geq 5$  mm) SWDs occurred in each group,  $p > 0.9$ . SWD was associated with increased excision dimensions in the NPWT group only,  $p = 0.03$ . Median duration to wound healing was equivalent in non-dehiscenced wounds, [CP 21.0 (14.0-29.5) versus NPWT 21.0 (16.0-24.0) days,  $p = 0.7$ ]. There were no differences in mean time to the following: return to school/work (NPWT  $26.1 \pm 18.2$  versus CP  $29.3 \pm 14.7$  d,  $p = 0.6$ ), sit normally (NPWT  $22.3 \pm 16.2$  versus CP  $20.1 \pm 9.4$  d,  $p = 0.7$ ), or return to physical activity (NPWT  $21.6 \pm 17.2$  versus CP  $40.3 \pm 2.4$  d,  $p = 0.2$ ). The authors concluded no difference in the rates of SWD. In addition, NPWT was not found to improve post-operative analgesia utilization, time to wound healing, patient satisfaction, or return to normal activities. Therefore, NPWT is not recommended to prevent SWD. Limitations of the study include small sample size, nonblinded design and potential attrition and performance bias.

A Hayes report on the use of NPWT after surgery for pilonidal disease concluded that the current body of overall very-low quality evidence does not allow for conclusions to be drawn regarding the benefits and potential associated risks of NPWT as a treatment adjunct over standard wound care methods alone. There is a need for additional, larger well-designed studies to evaluate this therapy more thoroughly and to determine which patients may benefit from NPWT after surgery for pilonidal disease (Hayes, 2020; updated 2023).

Danne et al. (2017) conducted a retrospective chart analysis of pilonidal sinus healing using NPWT versus alginate or gauze dressings. Thirty-two patients received NPWT and 30 received daily dressings. The median time to healing in the group receiving daily dressings was 10 weeks compared to 8 weeks in the group receiving NPWT. Among patients who healed, the difference in average time to healing was 5.2 weeks. However, the differences were not statistically significant. Study limitations include retrospective design and small patient numbers. Larger prospective, RCTs are needed to evaluate the efficacy of NPWT for treating pilonidal disease.

## **NPWT With Instillation of Wound Solutions**

De Pellegrin et al. (2023) conducted a systematic review and meta-analysis to compare negative pressure wound therapy with solution instillation and dwell time (NPWTi-d) with NPWT and standard of care for wound management in orthoplastic surgery. A comprehensive literature search using PubMed, Web of Science, and Cochrane databases was performed, including studies describing the outcomes of NPWTi-d for traumatic/Orthopedic injuries. A meta-analysis on the number of surgical debridements, as well as the rate of complete wound closure and complications was carried out, although for other outcomes, a descriptive statistic was applied. Risk of bias and quality of evidence were assessed using the Downs & Black's Checklist for Measuring Quality. Thirteen studies with a total number of 871 patients were included, in which NPWTi-d demonstrated higher primary wound closure and lower complication rates ( $p < .05$ ). No difference in the number of surgical procedures required for final wound healing was observed. Moreover, five out of six studies showed better results for NPWTi-d when the change of the bioburden and bacterial count of the wound were analyzed. A singular study investigating the length of the hospital stay of patients treated with NPWTi-d showed a reduction in the latter. The authors concluded that the present meta-analysis shows that NPWTi-d is superior to NPTW or conventional dressings in orthoplastic wound care management, in terms of complete wound closure rate and the reduced number of complications. Still, the limited quality of the studies analyzed shows that future randomized studies are needed to confirm the benefits and to identify the most appropriate recommendations for using NPWTi-d in orthoplastic surgery, as well as to investigate the cost-effectiveness of this wound-dressing system. This systematic review and meta-analysis have several limitations. Firstly, the studies using NPWTi-d applied the latter to different anatomical regions of the body, treating wounds of variable size and comparing different wound closure techniques. Secondly, many different wound care products were used throughout the studies with regard to the control group, which may also have had an influence on the results.

Several studies with a shorter follow-up time that did not report on the duration of treatment may have biased the results by underreporting the complication rate in the long run. Finally, the overall heterogeneity of the available studies and their limited quality made it difficult to properly investigate all outcomes related to these kinds of treatments. In view of this, a meta-analysis could only be performed on several of the outcomes. Future studies are needed to confirm the study findings, as well as to better document and quantify the potential benefits of NPWTi-d for wound care in the orthoplastic field.

Diehm et al. (2021) conducted a systematic review evaluating the use of NPWT with instillation and dwell time (NPWTi-d) for the treatment of acute and traumatic wounds. Ten articles (n = 109 acute and traumatic wounds) met inclusion criteria. No high-quality RCTs were identified. The majority of studies were retrospective cohort studies, followed by lesser-quality RCTs, comparative studies or prospective cohorts, and 2 retrospective comparative studies. While NPWTi-d showed promise to be effective in facilitating wound closure and reducing the time for wound closure, the authors found a relatively low level of evidence to support this effect. Large prospective, RCTs are necessary to determine the role of NPWTi-d in the clinical routine for this wound category.

Gabriel et al. (2021) performed a systematic review and meta-analysis of comparative studies evaluating the effects of NPWTi-d versus standard wound care in the treatment of multiple wound types. Thirteen studies (n = 720) were included in the analysis. NPWTi-d, when used in conjunction with good clinical practice (e.g., debridement, appropriate antibiotics), was found to be more beneficial than the comparator with respect to number of surgical debridements during therapy, time to readiness for final wound closure, number of patients with reduced bacterial bioburden, duration of therapy, and number of wounds closed, but similar with respect to hospital length of stay. However, author-noted study limitations, including low-level evidence and high patient and wound population heterogeneity across studies, suggested cautious interpretation of the results. Large prospective, RCTs are needed to confirm these results.

Kanapathy et al. (2020) conducted a systematic review and meta-analysis of studies evaluating the efficacy of NPWTi-d. Thirteen studies were included with a total of 624 wounds in 542 patients involving wounds of various etiology. These included surgical wounds (n = 186), trauma (n = 112), pressure ulcers (n = 73), neuropathic (n = 56), infection (n = 28), diabetic ulcers (n = 20), necrotizing fasciitis (n = 19), burns (n = 15), venous (n = 10) and vasculitis (n = 2). Normal saline was the most commonly used instillation solution. The pooled proportion of wounds that achieved complete healing was 93.65%. The authors concluded that although NPWTi-d has versatility to improve wound healing in a broad range of wounds, these conclusions are limited by the lack of high-quality level 1 evidence. The included studies were mostly small retrospective case series where NPWTi-d was performed on wounds of various etiologies and sizes along with different wound closure techniques. RCTs evaluating the efficacy of NPWTi-d against NPWT or standard dressings are needed.

## **Clinical Practice Guidelines**

### ***American Society of Colon and Rectal Surgeons (ASCRS)***

ASCRS practice parameters for the management of pilonidal disease do not specifically address NPWT as a treatment option (Johnson et al., 2019).

### ***International Working Group on the Diabetic Foot (IWGDF)***

IWGDF evidence-based guidelines on the prevention and management of diabetic foot disease (Rayman et al., 2020) make the following recommendations:

- Consider the use of NPWT to reduce wound size, in addition to best standard of care, in patients with diabetes and a post-operative (surgical) wound on the foot. (GRADE Strength of recommendation: Weak; Quality of evidence: Low).
- We suggest not using NPWT in preference to best standard of care in non-surgical diabetic foot ulcers. (GRADE Strength of recommendation: Weak; Quality of evidence: Low).

### ***National Institute for Health and Care Excellence (NICE)***

A NICE guideline concluded that the VAC Veraflo Therapy system (wound instillation with negative pressure therapy) shows promise for treating acute infected or chronic wounds that are not healing. However, there is not enough good-quality evidence to support the case for routine adoption. Further research is recommended to show clinically meaningful benefits for the device compared with NPWT alone (NICE, 2021a).

An amended NICE guideline suggests considering NPWT after cesarean birth for women with a body mass index (BMI) of 35 or more to reduce the risk of wound infections. The “consider” recommendation reflects that the evidence of benefit is less certain (NICE, 2021b).

A NICE guideline concluded that PICO negative pressure wound dressings should be considered as an option for closed surgical incisions in people who are at high risk of developing SSIs. They are associated with fewer SSIs and seromas

compared with standard wound dressings. The report called out the clinical and statistical heterogeneity of the studies as a limitation. It also noted a wide variation in the risk characteristics of the populations, the definition of SSIs, how long the dressing was in place and the length and frequency of follow up (NICE, 2019).

### ***National Pressure Injury Advisory Panel (NPIAP)***

NPIAP guidelines recommend considering NPWT as an early adjunct therapy for reducing the size and depth of Stage III and IV pressure injuries (NPIAP, 2019).

#### Strength of Evidence

- Level 1 studies of moderate or low quality providing direct evidence
- Level 2 studies of high or moderate quality providing direct evidence
- Most studies have consistent outcomes and inconsistencies can be explained

Strength of Recommendation – weak positive recommendation.

### ***Society for Vascular Surgery (SVS)***

SVS, in collaboration with the American Podiatric Medical Association and the Society for Vascular Medicine, makes the following recommendations on the management of diabetic foot ulcers (Hingorani et al., 2016):

- Standard of care for diabetic foot ulcers will lead to improvement in the majority of cases, and only in those cases without improvement should adjunctive modalities be used.
- For diabetic foot ulcers that fail to demonstrate improvement (> 50% wound area reduction) after a minimum of 4 weeks of standard wound therapy, the guidelines recommend adjunctive wound therapy options, including NPWT. Choice of adjuvant therapy is based on clinical findings. Re-evaluation of vascular status, infection control and off-loading are recommended to ensure optimization before initiation of adjunctive wound therapy (Grade 1B – strong recommendation based on moderate-quality evidence).
- The guidelines suggest the use of NPWT for chronic diabetic foot wounds that do not demonstrate expected healing progression with standard or advanced wound dressings after 4 to 8 weeks of therapy (Grade 2B – weak recommendation based on moderate-quality evidence).

### ***Wound Healing Society (WHS)***

WHS wound care guidelines make the following recommendations:

- Consider using NPWT for stage III or IV pressure ulcers that fail to progress in healing with conventional therapy. Current evidence indicates that NPWT may support pressure ulcer healing by increasing wound perfusion and formation of granulation tissue and by reducing bacterial load (Gould et al., 2016). Level I evidence – a meta-analysis of multiple RCTs or at least two RCTs supporting the intervention.
- NPWT may be useful prior to a skin graft/flap by helping promote the development of granulation tissue in the wound base, or postoperatively by preventing shearing and removing exudates. However, its reported experience in venous ulcers is limited (Marston et al., 2016). Level II - at least one RCT and at least two significant clinical series or expert opinion papers with literature reviews supporting the intervention.
- NPWT has been shown to increase the proportion of wounds that heal, and the rate of wound healing compared with standard wound care in diabetic lower extremity wounds (Lavery et al., 2016). Level I evidence – a meta-analysis of multiple RCTs or at least two RCTs supporting the intervention.

## **U.S. Food and Drug Administration (FDA)**

This section is to be used for informational purposes only. FDA approval alone is not a basis for coverage.

For information on NPWT systems, refer to the following website (use product code OMP):

<http://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfPMN/pmnm.cfm>. (Accessed January 2, 2025)

## **References**

Alves AS, Martineau J, Scampa M, et al. Negative pressure wound therapy versus conventional dressing in lower limb fractures: Systematic review and meta-analysis. *Plast Reconstr Surg Glob Open*. 2024 May 15;12(5):e5806.

Anghel EL, Kim PJ. Negative pressure wound therapy: A comprehensive review of the evidence. *Plast Reconstr Surg*. 2016 Sep;138(3 Suppl):129S-37S.

Azzopardi EA, Boyce DE, Dickson WA, et al. Application of topical negative pressure (vacuum-assisted closure) to split-thickness skin grafts: A structured evidence-based review. *Ann Plast Surg.* 2013 Jan;70(1):23-9.

Cheng Y, Wang K, Gong J, et al. Negative pressure wound therapy for managing the open abdomen in non-trauma patients. *Cochrane Database Syst Rev.* 2022 May 6;5(5):CD013710.

Costa ML, Achten J, Knight R, et al.; WHIST Trial Collaborators. Effect of incisional negative pressure wound therapy vs standard wound dressing on deep surgical site infection after surgery for lower limb fractures associated with major trauma: The WHIST randomized clinical trial. *JAMA.* 2020 Feb 11;323(6):519-526.

Costa ML, Achten J, Bruce J, et al.; UK WOLFF Collaboration. Effect of negative pressure wound therapy vs standard wound management on 12-month disability among adults with severe open fracture of the lower limb: The WOLFF randomized clinical trial. *JAMA.* 2018 Jun 12;319(22):2280-2288.

Danne J, Gwini S, McKenzie D, Danne P. A retrospective study of pilonidal sinus healing by secondary intention using negative pressure wound therapy versus alginate or gauze dressings. *Ostomy Wound Manage.* 2017 Mar;63(3):47-53.

De Pellegrin L, Feltri P, Filardo G, et al. Effects of negative pressure wound therapy with instillation and dwell time (NPWTi-d) versus NPWT or standard of care in orthoplastic surgery: A systematic review and meta-analysis. *Int Wound J.* 2023 Aug;20(6):2402-2413.

Diehm YF, Fischer S, Wirth GA, et al. Management of acute and traumatic wounds with negative-pressure wound therapy with instillation and dwell time. *Plast Reconstr Surg.* 2021 Jan 1;147(1S-1):43S-53S.

Dumville JC, Land L, Evans D, Peinemann F. Negative pressure wound therapy for treating leg ulcers. *Cochrane Database Syst Rev.* 2015 Jul 14;(7):CD011354.

ECRI. Negative pressure wound therapy for preventing surgical site infections after cesarean section in women with obesity. *Clinical Evidence Assessment.* 2024 Jan.

Ensor N, Martin S, Chang A, et al. Negative pressure dressing versus conventional passive dressing in pilonidal surgery: A randomized controlled trial. *J Surg Res.* 2024 Nov;303:313-321.

Gabriel A, Camardo M, O'Rorke E, et al. Effects of negative-pressure wound therapy with instillation versus standard of care in multiple wound types: Systematic literature review and meta-analysis. *Plast Reconstr Surg.* 2021 Jan 1;147(1S-1):68S-76S.

Gould L, Stuntz M, Giovannelli M, et al. Wound Healing Society 2015 update on guidelines for pressure ulcers. *Wound Repair Regen.* 2016 Jan-Feb;24(1):145-62.

Gustilo RB, Anderson JT. Prevention of infection in the treatment of one thousand and twenty-five open fractures of long bones: Retrospective and prospective analyses. *J Bone Joint Surg Am.* 1976 Jun;58(4):453-8.

Gustilo RB, Mendoza RM, Williams DN. Problems in the management of type III (severe) open fractures: a new classification of type III open fractures. *J Trauma.* 1984 Aug;24(8):742-6.

Haidari S, IJpma FFA, Metsemakers WJ, et al. The Role of Negative-Pressure Wound Therapy in Patients with Fracture-Related Infection: A Systematic Review and Critical Appraisal. *Biomed Res Int.* 2021 Oct 19;2021:7742227.

Hannan E, Harding T, Feizal H, et al. Negative pressure wound therapy following excision of pilonidal sinus disease: A retrospective review. *Colorectal Dis.* 2021 Nov;23(11):2961-2966.

Hayes, Inc. Clinical Research Response. Negative pressure wound therapy (NPWT) for clean, closed, orthopedic incisions of the ankle and lower leg. Lansdale, PA: Hayes, Inc.; July 2021a. Archived 2025.

Hayes, Inc. Clinical Research Response. Negative pressure wound therapy (NPWT) for primary arthroplasty of the hip or knee. Lansdale, PA: Hayes, Inc.; July 2021b. Archived 2025.

Hayes, Inc. Clinical Research Response. Negative pressure wound therapy (NPWT) for revision arthroplasty of the hip or knee. Lansdale, PA: Hayes, Inc.; July 2021c. Archived 2025.

Hayes, Inc. Evidence Analysis Research Brief. Negative pressure wound therapy for closed surgical incisions following total joint arthroplasty. Lansdale, PA: Hayes, Inc.; October 2022. Archived 2025.

Hayes, Inc. Evidence Analysis Research Brief. Outpatient negative pressure wound therapy for treatment of chronic wounds. Lansdale, PA: Hayes, Inc; March 2022.

Hayes, Inc. Evolving Evidence Review. PICO Single Use Negative Pressure Wound Therapy System (Smith & Nephew) for cesarean birth wound care. Lansdale, PA: Hayes, Inc.; April 2021d. Updated July 2024.

Hayes, Inc. Hayes Health Technology Assessment. Negative pressure wound therapy for chronic wounds: home use. Lansdale, PA: Hayes, Inc.; December 2016. Updated May 2021.

Hayes, Inc. Hayes Health Technology Assessment. Negative pressure wound therapy after surgery for pilonidal disease. Lansdale, PA: Hayes, Inc.; February 2020. Updated March 2023.

Hayes, Inc. Hayes Health Technology Assessment. Prophylactic negative pressure wound therapy in elective open abdominal surgeries. Lansdale, PA: Hayes, Inc.; February 2021e. Updated March 2024.

Hingorani A, LaMuraglia GM, Henke P, et al. The management of diabetic foot: A clinical practice guideline by the Society for Vascular Surgery in collaboration with the American Podiatric Medical Association and the Society for Vascular Medicine. *J Vasc Surg*. 2016 Feb;63(2 Suppl):3S-21S.

Hurd T, Kirsner RS, Sancho-Insenser JJ, et al. International Consensus Panel Recommendations for the optimization of traditional and single-use negative pressure wound therapy in the treatment of acute and chronic wounds. *Wounds*. 2021 Feb;33(suppl 2):S1-S11.

Hussamy DJ, Wortman AC, McIntire DD, et al. Closed incision negative pressure therapy in morbidly obese women undergoing cesarean delivery: a randomized controlled trial. *Obstet Gynecol*. 2019 Oct;134(4):781-789.

Iheozor-Ejiofor Z, Newton K, Dumville JC, et al. Negative pressure wound therapy for open traumatic wounds. *Cochrane Database Syst Rev*. 2018 Jul 3;7:CD012522.

Johnson EK, Vogel JD, Cowan ML, et al.; Clinical Practice Guidelines Committee of the American Society of Colon and Rectal Surgeons. The American Society of Colon and Rectal Surgeons' clinical practice guidelines for the management of pilonidal disease. *Dis Colon Rectum*. 2019 Feb;62(2):146-157.

Kanapathy M, Mantelakis A, Khan N, et al. Clinical application and efficacy of negative pressure wound therapy with instillation and dwell time (NPWTi-d): A systematic review and meta-analysis. *Int Wound J*. 2020 Dec;17(6):1948-1959.

Kirsner R, Dove C, Reyzelman A, et al. A prospective, randomized, controlled clinical trial on the efficacy of a single-use negative pressure wound therapy system, compared to traditional negative pressure wound therapy in the treatment of chronic ulcers of the lower extremities. *Wound Repair Regen*. 2019 Sep;27(5):519-529.

Lavery LA, Davis KE, Berriman SJ, Braun L, et al. WHS guidelines update: Diabetic foot ulcer treatment guidelines. *Wound Repair Regen*. 2016 Jan-Feb;24(1):112-26.

Liu Z, Dumville JC, Hinchliffe RJ, et al. Negative pressure wound therapy for treating foot wounds in people with diabetes mellitus. *Cochrane Database Syst Rev*. 2018 Oct 17;10:CD010318.

Liu S, He CZ, Cai YT, et al. Evaluation of negative pressure wound therapy for patients with diabetic foot ulcers: Systematic review and meta-analysis. *Ther Clin Risk Manag*. 2017 Apr 18;13:533-544.

Mantyh C, Silverman R, Collinsworth A, et al. Closed incision negative pressure therapy versus standard of care over closed abdominal incisions in the reduction of surgical site complications: A systematic review and meta-analysis of comparative studies. *Eplasty*. 2024 May 30;24:e33. PMID: 38846511; PMCID: PMC11155374.

Marston W, Tang J, Kirsner RS, Ennis W. Wound Healing Society 2015 update on guidelines for venous ulcers. *Wound Repair Regen*. 2016 Jan-Feb;24(1):136-44.

Moreno Gijón M, Suárez Sánchez A, de Santiago Álvarez I, et al. The efficacy of negative-pressure wound therapy (NPWT) in the prevention of surgical site occurrences in open abdominal surgery: A randomized clinical trial. *Surgery*. 2025 Feb;178:108920.

National Institute for Health and Care Excellence (NICE). Medical Technologies Guidance. MTG54. The VAC Veraflo Therapy system for acute infected or chronic wounds that are failing to heal. January 2021a.

National Institute for Health and Care Excellence (NICE). NICE Guideline. NG192. Caesarean birth. March 2021b.

National Institute for Health and Care Excellence (NICE). Medical Technologies Guidance. MTG43. PICO negative pressure wound dressings for closed surgical incisions. May 2019.

National Pressure Injury Advisory Panel (NPIAP), European Pressure Ulcer Advisory Panel and Pan Pacific Pressure Injury Alliance. Prevention and treatment of pressure ulcers/injuries: Clinical practice guideline. Emily Haesler (Ed.). 2019.

Norman G, Goh EL, Dumville JC, et al. Negative pressure wound therapy for surgical wounds healing by primary closure. *Cochrane Database Syst Rev*. 2020 Jun 15;6(6):CD009261.

Norman G, Shi C, Goh EL, et al. Negative pressure wound therapy for surgical wounds healing by primary closure. *Cochrane Database Syst Rev*. 2022 Apr 26;4(4):CD009261.

Pan A, De Angelis G, Nicastrì E, et al. Topical negative pressure to treat surgical site infections, with a focus on post-sternotomy infections: A systematic review and meta-analysis. *Infection*. 2013 Dec;41(6):1129-35.

Rayman G, Vas P, Dhatariya K, et al.; International Working Group on the Diabetic Foot (IWGDF). Guidelines on use of interventions to enhance healing of chronic foot ulcers in diabetes (IWGDF 2019 update). *Diabetes Metab Res Rev*. 2020 Mar;36 Suppl 1:e3283.

Rys P, Borys S, Hohendorff J, et al. NPWT in diabetic foot wounds - a systematic review and meta-analysis of observational studies. *Endocrine*. 2020 Apr;68(1):44-55.

Şahin E, Rizalar S, Özker E. Effectiveness of negative-pressure wound therapy compared to wet-dry dressing in pressure injuries. *J Tissue Viability*. 2022 Feb;31(1):164-172.

Scalise A, Calamita R, Tartaglione C, et al. Improving wound healing and preventing surgical site complications of closed surgical incisions: a possible role of incisional negative pressure wound therapy. A systematic review of the literature. *Int Wound J*. 2016 Dec;13(6):1260-1281.

Singh DP, Gabriel A, Parvizi J, et al. Meta-analysis of comparative trials evaluating a single-use closed-incision negative-pressure therapy system. *Plast Reconstr Surg*. 2019 Jan;143:41S-46S.

Strugala V, Martin R. Meta-analysis of comparative trials evaluating a prophylactic single-use negative pressure wound therapy system for the prevention of surgical site complications. *Surg Infect (Larchmt)*. 2017 Oct;18(7):810-819.

Tansarli GS, Vardakas KZ, Stratoulis C, et al. Vacuum-assisted closure versus closure without vacuum assistance for preventing surgical site infections and infections of chronic wounds: A meta-analysis of randomized controlled trials. *Surg Infect (Larchmt)*. 2014 Aug;15(4):363-7.

Tuuli MG, Liu J, Tita ATN, et al. Effect of prophylactic negative pressure wound therapy vs standard wound dressing on surgical-site infection in obese women after cesarean delivery: A randomized clinical trial. *JAMA*. 2020 Sep 22;324(12):1180-1189.

Vig S, Dowsett C, Berg L, et al.; International Expert Panel on Negative Pressure Wound Therapy [NPWT-EP]. Evidence-based recommendations for the use of negative pressure wound therapy in chronic wounds: Steps towards an international consensus. *J Tissue Viability*. 2011 Dec;20 Suppl 1:S1-18.

Virani SR, Dahapute AA, Bava SS, Muni SR. Impact of negative pressure wound therapy on open diaphyseal tibial fractures: A prospective randomized trial. *J Clin Orthop Trauma*. 2016 Oct-Dec;7(4):256-259.

Wang N, Li SS, Liu YP, et al. Comparison of negative pressure wound therapy and moist wound care in patients with diabetic foot ulcers: A protocol for systematic review and meta-analysis of randomized controlled trials. *Medicine (Baltimore)*. 2022 Aug 5;101(31):e29537.

Wu Y, Shen G, Hao C. Negative pressure wound therapy (NPWT) is superior to conventional moist dressings in wound bed preparation for diabetic foot ulcers: A randomized controlled trial. *Saudi Med J*. 2023 Oct;44(10):1020-1029.

Zhang J, Hu ZC, Chen D, et al. Effectiveness and safety of negative pressure wound therapy for diabetic foot ulcers: a meta-analysis. *Plast Reconstr Surg*. 2014 Jul;134(1):141-51.

## Policy History/Revision Information

Date	Summary of Changes
04/01/2026	<ul style="list-style-type: none"> <li>Retired policy; Louisiana plan membership disenrolled on Apr. 1, 2026</li> </ul>
08/01/2025	<p><b>Coverage Rationale</b></p> <ul style="list-style-type: none"> <li>Revised list of contraindications to negative pressure wound therapy (NPWT):               <ul style="list-style-type: none"> <li>Added “anticoagulation therapy”</li> <li>Replaced “uncontrolled soft tissue infection or osteomyelitis within vicinity of wound” with “uncontrolled soft tissue infection or <i>untreated</i> osteomyelitis within vicinity of wound”</li> </ul> </li> </ul> <p><b>Supporting Information</b></p> <ul style="list-style-type: none"> <li>Updated <i>Clinical Evidence</i> and <i>References</i> sections to reflect the most current information</li> <li>Archived previous policy version CS157LA.B</li> </ul>

## Instructions for Use

This Medical Policy provides assistance in interpreting UnitedHealthcare standard benefit plans. When deciding coverage, the federal, state or contractual requirements for benefit plan coverage must be referenced as the terms of the federal, state or contractual requirements for benefit plan coverage may differ from the standard benefit plan. In the event of a conflict, the federal, state or contractual requirements for benefit plan coverage govern. Before using this policy, please check the federal, state or contractual requirements for benefit plan coverage. UnitedHealthcare reserves the right to

modify its Policies and Guidelines as necessary. This Medical Policy is provided for informational purposes. It does not constitute medical advice.

UnitedHealthcare may also use tools developed by third parties, such as the InterQual® criteria, to assist us in administering health benefits. The UnitedHealthcare Medical Policies are intended to be used in connection with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice.