

Gastrointestinal Disorders Diagnostic Procedures (for New Mexico Only)

Policy Number: CS046NM.D
Effective Date: April 1, 2026

[Instructions for Use](#)

Table of Contents	Page
Application	1
Coverage Rationale	1
Definitions	1
Applicable Codes	2
Description of Services	2
Clinical Evidence	3
U.S. Food and Drug Administration	12
References	13
Policy History/Revision Information	15
Instructions for Use	15

Related Policies
<ul style="list-style-type: none"> Bariatric Surgery (for New Mexico Only) Minimally Invasive Procedures for the Treatment of Upper Gastrointestinal Diseases (for New Mexico Only)

Application

This Medical Policy only applies to the state of New Mexico.

Coverage Rationale

The following procedures are unproven and not medically necessary due to insufficient evidence of efficacy:

- [Magnetic Resonance Imaging \(MRI\) Defecography](#) for evaluating [Constipation](#) and [Anorectal](#) or pelvic floor disorders
- Cutaneous, mucous, or serosal [Electrogastrography](#), electroenterography, or body surface gastric mapping for diagnosing intestinal or gastric disorders including [Gastroparesis](#)
- [Esophageal Mucosal Integrity Testing](#) by electrical impedance for the diagnosis of gastroesophageal reflux disease (GERD), eosinophilic esophagitis (EoE), and nonacid reflux disease (non-GERD), or for the monitoring of treatment response in GERD and EoE
- [Functional Lumen Imaging Probe \(FLIP\)](#) technology for diagnosing [Achalasia](#)

Definitions

Refer to the federal, state, or contractual definitions that supersede the definitions below.

Achalasia: A primary esophageal motor disorder of unknown etiology characterized by degeneration of the myenteric plexus, which results in impaired relaxation of the esophagogastric junction (EGJ), along with the loss of organized peristalsis in the esophageal body [American Society of Gastrointestinal Endoscopy (ASGE)].

Anorectal Disorders: Structural or functional abnormalities of the anorectum or pelvic floor (Patcharatrakul and Rao, 2018).

Constipation: Infrequent or hard-to-pass bowel movements, hard stools, or incomplete bowel movement sensation; infrequent means less than three bowel movements a week (Bharucha et al., 2013a).

Electrogastrography (EGG): A non-invasive method for the measurement of gastric myoelectrical activity using cutaneous electrodes placed on the abdominal skin over the stomach (Yin and Chen, 2013).

Esophageal Mucosal Integrity Testing: An adjunct procedure to routine endoscopy to speed the diagnostic differentiation of GERD from non-GERD and EoE that is intended to be an alternative to 24- to 48- hour-long pH monitoring and biopsies (Hayes, 2022; updated 2024).

Fecal Incontinence (FI): The inability to control bowel movements causing stool (feces) to leak unexpectedly from the rectum; also called bowel or anal incontinence (Bharucha et al., 2013a).

Functional Lumen Imaging Probe (FLIP): A catheter-based endoscopic system designed to assess esophageal motility and the distensibility index at the esophagogastric junction; the measurement provides insights into the compliance of the lower esophageal sphincter (Alabbas et al., 2025).

Gastroparesis: A digestive disorder in which the motility of the stomach is either abnormal or absent; it is also known as delayed gastric emptying (Camilleri, 2013, updated 2022).

Magnetic Resonance Defecography: A noninvasive test that uses magnetic resonance imaging to obtain images at various stages of defecation to evaluate how well the pelvic muscles are working and provide insight into rectal function (RadiologyInfo.org); it can evaluate pelvic floor anatomy, dynamic motion, and rectal evacuation simultaneously (Rao and Patcharatrakul, 2016).

Applicable Codes

The following list(s) of procedure and/or diagnosis codes is provided for reference purposes only and may not be all inclusive. Listing of a code in this policy does not imply that the service described by the code is a covered or non-covered health service. Benefit coverage for health services is determined by federal, state, or contractual requirements and applicable laws that may require coverage for a specific service. The inclusion of a code does not imply any right to reimbursement or guarantee claim payment. Other Policies and Guidelines may apply.

CPT Code	Description
0779T	Gastrointestinal myoelectrical activity study, stomach through colon, with interpretation and report
0868T	High-resolution gastric electrophysiology mapping with simultaneous patient-symptom profiling, with interpretation and report
43499	Unlisted procedure, esophagus
72195	Magnetic resonance (e.g., proton) imaging, pelvis; without contrast material(s)
72196	Magnetic resonance (e.g., proton) imaging, pelvis; with contrast material(s)
72197	Magnetic resonance (e.g., proton) imaging, pelvis; without contrast material(s), followed by contrast material(s) and further sequences
76498	Unlisted magnetic resonance procedure (e.g., diagnostic, interventional)
91132	Electrogastrography, diagnostic, transcutaneous
91133	Electrogastrography, diagnostic, transcutaneous; with provocative testing

CPT® is a registered trademark of the American Medical Association

Description of Services

Several gastrointestinal disorders, such as Constipation, Fecal Incontinence, and Gastroparesis, may require testing before a diagnosis can be made.

Symptoms of Constipation, one of the most common digestive problems, are extremely common. The prevalence of Constipation is approximately 16% in adults overall and 33% in adults over 60. If symptoms do not improve, investigations to diagnose rectal evacuation disorders and slow-transit Constipation are sometimes performed, such as digital rectal examination, anorectal structure, and function testing (including the balloon expulsion test, anorectal manometry or defecography) or colonic transit tests (such as the radiopaque marker test, scintigraphy) (Camilleri et al., 2017). While in most cases, Constipation is benign and due to dietary and lifestyle factors, Constipation is sometimes due to disordered colonic and/or pelvic floor/anorectal function.

Fecal Incontinence (FI) is the inability to control bowel movements causing stool to leak unexpectedly from the rectum. Continence requires the rectum, anus, and nervous system to be working normally. FI is commonly caused by altered stools (generally diarrhea, but also Constipation) or conditions that affect the ability of the rectum and anus to hold stool.

Individuals with Gastroparesis may experience symptoms of frequent nausea and vomiting, early satiety, bloating, postprandial fullness, and epigastric pain and burning. Although Gastroparesis can occur with no obvious cause, patients with diabetes frequently develop this condition. If Gastroparesis causes nausea and persistent vomiting, it can lead to frequent hospitalization for hypoglycemia, hyperglycemia, acidosis, dehydration, pseudo-obstruction, electrolyte dyscrasias, or other complications. The diagnosis of Gastroparesis requires objective evidence of clearly delayed gastric emptying in symptomatic patients. Scintigraphy is the reference standard for measurement of gastric emptying. Protocols for standardized meals prior to scintigraphy have been recommended, however for interpretation of test results, it has to be considered that clinical utility depends on complete consumption of adequate test meals and adequate duration of imaging. For all gastrointestinal function tests, adherence to adequately validated, standardized study protocols is crucial (Keller et al., 2018).

Electrogastrography (EGG) is a non-invasive technique for recording gastric myoelectrical activity using cutaneous electrodes placed on the abdominal skin over the stomach. The surface recording obtained using electrography is called the electrogastrogram. Gastric myoelectrical activity may be altered or become abnormal in diseased states or upon provocative stimulations or even spontaneously. Abnormal gastric myoelectrical activity includes gastric dysrhythmia, abnormal slow wave propagation, and electro-mechanical uncoupling. In the stomach, there is lack of one-to-one correlation between spikes and contractions, and thus this abnormality cannot be accurately detected from the in vivo myoelectrical recording. In individuals with gastrointestinal motility disorders or individuals with functional gastrointestinal diseases, EGG is used to identify the pathophysiology of the diseases associated with gastric slow waves or dysrhythmia (Yin and Chen, 2013). Electroenterography is a similar procedure that records myoelectrical activity from the intestines and body surface gastric mapping (BGSM) uses high resolution electrode arrays, along with bioelectronics, automated artifact rejection, and analytics, to measure and map gastric myoelectric activity.

Anorectal Disorders present with a variety of symptoms and result from either structural or functional disorders. Clinical correlation is essential before labeling an abnormal finding as clinically significant. Together with a detailed history, a thorough physical and digital rectal examination and appropriate testing, in most patients the underlying cause and type of anorectal disorder can be correctly identified, and treatment can be tailored (Patcharatrakul and Rao, 2018).

Defecatory disorders are primarily characterized by impaired rectal evacuation from inadequate rectal propulsive forces and/or increased resistance to evacuation; the latter may result from high anal resting pressure (“anismus”) and/or incomplete relaxation or paradoxical contraction of the pelvic floor and external anal sphincters (“dyssynergia”) during defecation. Structural disturbances (e.g., rectocele, intussusception) and reduced rectal sensation may coexist.

Magnetic Resonance Imaging (MRI) Defecography is being studied as an imaging tool that may provide an enhanced view of the bowel movement process including the underlying anatomic and pathophysiologic background of pelvic floor disorders. It can evaluate pelvic floor anatomy, dynamic motion, and rectal evacuation simultaneously (Rao and Patcharatrakul, 2016).

Endoluminal Functional Lumen Imaging Probe (FLIP) technology is used for esophageal function testing, such as testing for Achalasia or Diffuse Esophageal Spasm. FLIP is performed alone or in conjunction with other diagnostic tests; EndoFlip™ is one such device. A catheter surrounded by a balloon filled with liquid extends from the esophagus into the stomach or from the stomach into the small intestine. The device then applies pressure while providing continuous diameter measurements along its length, giving information about esophageal and/or gastric function (Hayes, 2022).

Clinical Evidence

Functional Lumen Imaging Probe (FLIP)

While FLIP technology is used to enhance current procedures such as endoscopy, the evidence is insufficient for use in assessing esophageal and gastric disorders. The clinical utility, in addition or instead of standard care, is yet to be defined for the diagnostic work-up, procedure guidance or post-procedure evaluation of clinical outcomes. Additional research involving larger RCTs with long-term outcomes is needed to establish its safety and efficacy.

In a systematic review and meta-analysis, Fehervari et al. (2025) evaluated the intraoperative use of functional luminal imaging probes (EndoFLIP™) during surgical procedures for achalasia focusing on the distensibility index (DI) and treatment outcomes. A total of 32 studies with 2681 patients were included in the qualitative synthesis, and 17 studies with 1734 patients were included in the quantitative synthesis. The authors concluded the EndoFLIP™ enhances achalasia treatment by providing real-time feedback on DI, which allowed more tailored interventions and improved outcomes. Limitations include the need for a standardization protocol for EndoFLIP™ usage to further validate its clinical utility ensuring consistent, comparable measurements. Additionally, the majority of studies were retrospective or prospective

with no randomized controlled trials identified. The study doesn't address the clinical impact of the technology on patient-centered outcomes.

An Evidence Analysis Research Brief was published by Hayes (2024) to summarize the newly published peer-reviewed literature and publications of the Endolumenal Functional Lumen Imaging Probe (EndoFLIP; Medtronic Inc.) for the evaluation of esophageal and gastric disorders. Since 2019, seven new studies were published. None of these studies were case-control studies evaluating EndoFLIP, randomized controlled trials (RCTs) comparing treatment guided by EndoFLIP™ or systematic reviews with or without meta-analysis. Based on the eight position states or identified guidelines, all appear to confer no/unclear support for the use of EndoFLIP™ specifically and impedance planimetry generally, to evaluate gastric and esophageal disorders. Limitations included lack of high-quality RCTs and this evidence analysis research brief is based on a review of study abstract only and is therefore not intended to evaluate the safety or efficacy of the health technology.

An ECRI (2024) revised Executive Summary was published evaluating the EndoFLIP Impedance Planimetry System (Medtronic plc.) for identifying gastrointestinal (GI) motility disorders. A total of 12 studies reporting on approximately 2100 individuals were reviewed to determine the clinical utility and clinical validity of the device for identifying GI disorders and guiding a plan of care. The evidence from the two cohort studies, five case-control studies and five comparative cohort studies was determined to be limited in quality to permit conclusions on EndoFLIP's clinical utility and validity for the evaluation of GI motility disorders using high-resolution manometry (HRM) or timed barium esophagram (TBE) as reference standards. Limitations include small study sample sizes, very-low quality evidence and the studies being at a high risk of bias due to retrospective design, lack of blinding and single-center focus. Four ongoing trials were identified involving EndoFLIP™ but none that may address the evidence gaps identified in this summary.

Rafeeqi et al. (2023) analyzed the utility of EndoFLIP™ in diagnosis and management of achalasia in pediatric patients. The authors theorized that the distensibility index (DI) may aid diagnosis and treatment of pediatric achalasia. Thirty-three (21 male, 12 female) patients underwent POEM and EndoFLIP™. The authors were able to show that certain measurements, such as Eckardt scores, had a direct correlation to achalasia symptoms. The higher the Eckardt score, the lower DI. While manometry is the current gold standard test for the diagnosis of achalasia, it is not tolerated well in children which makes the EndoFLIP™ a potential alternative for the pediatric population. The authors concluded that EndoFLIP™ could be beneficial in the pediatric population in the diagnosis and management of achalasia. The DI could be used to guide the extent of myotomy and monitor the results of any lower esophageal sphincter LES focused interventions, however further studies are necessary to better support the use of EndoFLIP™ as a diagnostic tool. Limitations included small sample size, lack of manometry data for comparison and lack of long-term data. Furthermore, the study did not address how the technology improves patients' outcomes compared to standard of care.

Benitez et al. (2021) conducted a retrospective study of pediatric individuals with achalasia, comparing those who underwent functional luminal imaging probe (FLIP) before and immediately after balloon dilations compared to a non-achalasia cohort. The primary goals of the study were to evaluate esophagogastric junction (EGJ) distensibility using FLIP measurements between the two groups and also correlate changes in EGJ distensibility with changes in Eckardt Symptom Score (ESS). The ESS uses a four-point self-reporting scale assessing the degree of dysphagia, regurgitation, weight loss and chest pain. The ESS is the most commonly utilized tool for assessing symptoms in individuals with achalasia. Thirty individuals with achalasia (mean age, 15.2; 40% female), including fourteen treatment-naïve and thirteen controls (mean age, 7.9 years old; 61% female) were identified. All individuals underwent esophageal manometry studies in order to diagnose achalasia prior to intervention. Additionally, 86.7% had an upper GI series or esophagram. Balloon dilations were performed under general anesthesia utilizing pneumatic dilator sizes of 30, 35 or 40 mm based on each pediatric individual's clinical history and endoscopist' preference in the individuals with achalasia. FLIP measurements for EGJ distensibility index (EGJ-DI), diameter and intraballloon pressures were obtained before and after dilations at 20, 30 and 40 ml FLIP balloon inflation volume in all thirty achalasia individuals. This study demonstrates pediatric individuals diagnosed with achalasia present with decreased EGJ-DI similar to adult individuals with achalasia and a significant increase in EGJ-DI immediately after balloon dilations in treatment-naïve and previously treated pediatric individuals with achalasia. Additionally, the authors were able to detect a significant increase in Eckardt score at one month following the balloon dilation, indicative of the ability dilations can improve symptoms. The authors conclude that balloon dilations remain an effective therapeutic method in achalasia with significant increase in EGJ-DI; assessed by improvement in symptoms and assessed by FLIP. Limitations include small sample size and a limited quantity of previous high-quality studies assessing pediatric EGJ distensibility data. The need for larger studies of pediatric individuals with achalasia would further assess the clinical utility of FLIP technology in the pediatric population and evaluate the long-term clinical impact and safety of this technology on pediatric individuals with achalasia.

A Hayes Health Technology Assessment (2019, updated 2022) focused on the use of the Endolumenal Functional Lumen Imaging Probe (EndoFLIP®) as a device for the evaluation, prognosis and guidance in the management of achalasia,

eosinophilic esophagitis and gastroparesis. Of the 13 studies evaluated, 11 reviewed the clinical utility of EndoFLIP® for detecting and predicting disease course, or treatment response. The other two studies evaluated the clinical utility for guidance and management of gastric disorders including achalasia. While the body of evidence for EndoFLIP® was large for a variety of topics, the quality of evidence was low with a large range of uses. The evidence is insufficient to support any conclusions for the use of the device for any one gastrointestinal disorder and found no improvement or benefit to health outcomes. Limitations included lack of comparison groups, lack of systematic evaluation and lack of long-term outcomes.

In a systematic review, Desprez et al. (2020) summarized the available data in the literature on the use of the EndoFLIP® system in the gastrointestinal tract. A search was conducted using MEDLINE-PubMed, Cochrane Library, and Google Scholar databases. A total of 95 studies were found and 32 of these reported results in patients with achalasia. The authors found the EndoFLIP® system demonstrated its relevance in the diagnosis of atypical achalasia and in the prediction of the treatment outcome in achalasia, but the precise esophago-gastric junction (EGJ) distensibility index (DI) threshold associated with success and prolonged response remains to be determined and therefore warrants additional studies. In addition, a review of thirteen studies was completed which investigated EGJ in GERD patients. Patients were assessed with upper GI endoscopy and esophageal pH monitoring in addition to symptomatic evaluation. The authors found the patients with GERD seemed to have a higher EGJ DI than the healthy volunteers tested, but the relevance of the EndoFLIP® system in the prediction of the efficacy of fundoplication remained unconfirmed. Limitations included no standard protocol found when using the EndoFLIP® in the esophagus along with cost and availability in daily practice. In conclusion, while the authors found EndoFLIP® might have some promise, its use in the body of the esophagus, other esophageal diseases (GERD, eosinophilic esophagitis), and other sphincter regions (anal canal, pylorus) necessitates the need for additional confirmatory studies.

Yoo et al. (2019) assessed the utility of EndoFLIP in fifty-two patients with achalasia treated with POEM and hypothesized that improvement in the distensibility index (DI) correlated with the postoperative clinical outcome of POEM. EndoFLIP was performed before and after POEM by well-trained endoscopists in an outpatient setting.

Esophagogastroduodenoscopy (EGD) was done just before EndoFLIP to measure esophageal length, four days following the procedure and again in outpatient clinic to evaluate the development of reflux esophagitis. The DI of the esophagogastric junction (EGJ) was measured at the point of narrowest perimeter of the functional lumen image during volumetric distension of the saline bag. After the probe was inserted, the DI and cross-sectional area (CSA) of the EGJ were measured at the site of the narrowest point. DI and CSA were measured twice during volumetric distension of the saline bag, each of them with 30 mL and 40 mL. The authors' conclusion was that EndoFLIP was useful in predicting the treatment response and post procedure reflux of POEM in patients with achalasia. Limitations included lack of comparison groups, lack of long-term outcomes and small sample size.

Clinical Practice Guidelines

American College of Gastroenterology (ACG)

In a 2021 ACG published clinical guideline (Katz, et al. 2021) for the diagnosis and management of GERD, the following recommendations are cited:

- Recommend antireflux surgery as an option for long-term treatment of patients with objective evidence of GERD, (strong recommendation; moderate level of evidence).
- Recommend consideration of magnetic sphincter augmentation (MSA) as an alternative to laparoscopic fundoplication for patients with regurgitation who fail medical management (strong recommendation, moderate level of evidence).
- Consideration of transoral incisionless fundoplication (TIF) for patients with troublesome regurgitation or heartburn who do not wish to undergo antireflux surgery and who do not have severe reflux esophagitis or hiatal hernias (conditional recommendation; low level of evidence).
- Do not recommend radiofrequency energy (Stretta) as an antireflux procedure due to inconsistent data on the efficacy of the device (conditional recommendation; low level of evidence).

In the 2020 ACG published clinical guideline (Vaezi, MF, et al. 2020) for the diagnosis and management of Achalasia indicate functional lumen imaging probe (FLIP) may be useful in evaluating individuals who cannot tolerate or complete a standard manometry as FLIP is performed during endoscopy while the individual is sedated. The authors note, based on consistent but low-quality data, the role of FLIP in achalasia is evolving and further studies are required to determine whether FLIP can replace or reduce the number of manometry studies and barium esophagrams in the management of achalasia.

American Gastroenterological Association (AGA)

The AGA published a clinical practice update on the functional lumen imaging probe for the management of esophageal disorders based on an expert review. The functional luminal imaging probe (FLIP) is a Food and Drug Administration–approved measurement tool used to measure simultaneous pressure and diameter to guide management of various upper gastrointestinal disorders. FLIP has been available since 2009 as the endoluminal functional lumen imaging probe (EndoFLIP). The EndoFlip has had limited penetrance into clinical settings outside of specialized centers primarily because of a paucity of data supporting its utility in general practice and a lack of standardized protocols and data analysis methodology. The AGA lists best practices for the management of esophageal disorders regarding FLIP in the following order:

- “Clinicians should not make a diagnosis or treatment decision based on functional lumen imaging probe (FLIP) assessment alone.
- FLIP assessment is a complementary tool to assess esophagogastric junction opening dynamics and the stiffness of the esophageal wall.
- Utilization should follow distinct protocols and analysis paradigms based on the disease state of interest.
- Clinicians should not utilize FLIP in routine diagnostic assessments of gastroesophageal reflux disease.
- FLIP should not be used to diagnose eosinophilic esophagitis but may have a role in severity assessment and therapeutic monitoring” (Hirano, 2017).

MRI Defecography

There is insufficient evidence regarding the effectiveness and efficacy of MRI defecography. Existing data suggest that this approach is not superior and, in some cases, inferior to conventional defecography.

In a retrospective single-center test development study with 46 adults with chronic constipation, Thanaracthanon et al. (2023) evaluated the diagnostic performance of MR defecographic findings in diagnosis of dyssynergic defecation (DD). Study participants were divided into two groups based on the presence of DD in two of three diagnostic tests (ARM, BET and anal surface electromyography), with 24 in the DD group (37.5% female) and 22 in the non-DD group (81.8% female). All patients underwent MR defecography according to the institutional standard protocol with both static and dynamic MR defecography images obtained. The study included analysis by two radiologists of nine parameters: anorectal angle (ARA) and M line at rest, defecation, and change between two phases; anal canal width; prominent puborectalis muscle; abnormal evacuation. The authors reported that seven of the nine parameters showed statistically significant difference between the DD and the non-DD group with M line at defecation having the highest odds ratio, followed by ARA change, ARA defecation, M line change, prominent puborectalis muscle, abnormal evacuation, and anal canal width, respectively. The authors also reported that the ARA change and prominent puborectalis muscle had the highest specificity and that multivariate logistic regression revealed two significant findings in differentiating between DD and non-DD, including M line at defecation and ARA at defecation. Limitations include the single-center, retrospective study design, the small sample size, the lack of a comparison test, the heterogeneity of other diseases in the non-DD group that were not seen in the DD group, and the use of a consensus panel which may incorporate bias. The authors concluded that MR defecography had high diagnostic performance in diagnosis of DD with ARA change of less than 1.5 degrees and prominent puborectalis muscle having good specificity in DD diagnosis. The authors recommend additional prospective studies with normal healthy patients as a control group. The findings need to be validated in an independent group of patients and the clinical utility of the test needs to be defined.

Pääkkö et al. (2022) completed a single-center, retrospective review of both magnetic resonance defecography (MRD) and video defecography (VD) studies that were done on 64 women with defecation disorders who underwent both VD and MRD within a year to compare the findings of the two methods and to analyze the success rates. In 58 patients, the indication for the first study were symptoms of obstructive defecation with incontinence as the primary diagnosis for the remaining six patients. The indication for the second study was insufficient information from the first study in 48 patients and for preoperative planning to get more anatomical information in the remaining 16 cases, the second imaging was performed before operative treatment to get more anatomical information or to confirm the findings of the first study. Both studies were analyzed in consensus by two radiologists who were blinded to clinical patient data and radiology reports. The authors reported that 96.9% of the VD studies were technically fully diagnostic compared with 45.3% for MRD and that 1.6% of the VD studies were partially diagnostic versus 32.8% for MRD. They reported that 30 enteroceles were observed by VD compared with seven in MRD with moderate agreement, 53 intussusceptions were observed by VD compared with 27 by MRD with poor agreement, 47 cases of rectocele were diagnosed by VD versus 29 by MRD with moderate agreement, and dyssynergic defecation was observed in three patients by VD and in 11 patients by MRD with slight agreement. Limitations of the study included the variability of which study was done first, the retrospective nature of the study, the small sample size, and the variability of the amount and consistency of the gel used in the studies. The authors concluded that technical success and diagnostic capabilities of VD were better than those of MRD and that VD remains the method of choice in the imaging of defecation disorders.

A Cochrane Database systematic review and meta-analysis by van Gruting, et al. (2021) evaluated imaging modalities for the detection of posterior pelvic floor disorders in women with obstructed defecation syndrome (ODS). The review included 39 studies [including the Foti (2013), Poncelet (2017), van Iersel (2017), Vitton (2011), and Zafar (2017) studies previously included in this section] with 2,483 women that evaluated the diagnostic accuracy of evacuation proctography (EP), dynamic magnetic resonance imaging (MRI) and pelvic floor ultrasound for detecting posterior pelvic floor disorders. The meta-analysis was done using Bayesian hierarchical latent class analysis and the overall quality of evidence (QoE) was assessed using the GRADE approach for diagnostic test accuracy. The authors reported that the sensitivity of EP for diagnosis of rectocele was 98%, enterocele 91% and pelvic floor descent 98% while the specificity of enterocele was 96%, intussusception 92% and anismus 97%, all with high QoE. The sensitivity for anismus of 80% and the specificity for rectocele of 78% and pelvic floor descent 83% had a moderate to low QoE. The specificity of MRI defecography for diagnosis of rectocele was 90%, enterocele 99% and intussusception 97% with high QoE. The heterogeneity analysis completed in the study showed that sensitivity of MRI performed with evacuation phase was higher than without for rectocele (94% with and 65% without), and for enterocele (87% with and 62% without), while the sensitivity of MRI without evacuation phase was significantly lower than EP. The study also showed that the specificity of transperineal ultrasound (TPUS) for diagnosis of rectocele was 89%, enterocele was 98% and intussusception 96% while the sensitivity for anismus was 92%. The authors concluded that neither MRI defecography nor TPUS met the criteria to replace EP as the reference standard for diagnosis of posterior pelvic floor disorders although both met the criteria of a triage test as a positive test confirms the diagnosis of rectocele, enterocele and intussusception and a negative test rules out diagnosis of anismus. The results of the other ultrasound techniques including endovaginal ultrasound, dynamic anal endosonography, and echodefecography were of too low a quality of evidence to draw conclusions. The authors recommended more well-designed studies to define the role of MRI defecography in the diagnostic pathway of ODS.

Ramage et al. (2018) assessed whether MRI features indicative of pelvic floor dysfunction correlated with patient-reported symptom severity. Univariate and multivariate analyses were performed using pre-treatment questionnaire responses to the Birmingham Bowel, Bladder and Urinary Symptom Questionnaire (BBUSQ), Wexner Incontinence Score (WIS), and modified Obstructed Defecation Symptom (ODS) Score. 302 MRI proctograms were performed (n = 170). Patients with a rectocele larger than two centimeters (p = 0.003; OR 5.756) or MRD features suggestive of puborectalis syndrome (p = 0.025; OR 8.602) were more likely to report a higher ODS score on multivariate analysis. Lack of rectal evacuation was negatively associated with an abnormal WIS (p = 0.007; OR 0.228). Age > 50 (p = 0.027, OR 2.204) and a history of pelvic floor surgery (p = 0.042, OR 0.359) were correlated with an abnormal BBUSQ incontinence score. Lack of rectal evacuation (p = 0.027, OR 3.602) was associated with an abnormal BBUSQ constipation score. Age > 50 (p = 0.07, OR 0.156) and the presence of rectoanal intussusception (p = 0.010, OR 0.138) were associated with an abnormal BBUSQ evacuation score. The authors concluded that while MRD is a useful tool in aiding multidisciplinary decision making, overall, it is poorly correlated with patient-reported symptom severity, and treatment decisions should not rest solely on results. Limitations of this study included lack of a reference standard test and a questionnaire with questions directed at only females.

In a systematic review and meta-analysis of MRD versus clinical examination and fluoroscopy, Ramage et al. (2017) compared detection and miss rates of pelvic floor abnormalities with MRD versus clinical examination and traditional fluoroscopic techniques. Twenty-eight studies were included: 14 studies compared clinical examination to MRD, and 16 compared fluoroscopic techniques to MRD. Detection and miss rates with MRD were not significantly different from clinical examination findings for any outcome except enterocele, where MRD fared significantly better than clinical examination. However, when comparing MRD versus fluoroscopy, MRD have no better detection rate or lower miss rate of a structural abnormality than fluoroscopy. In some studies, fluoroscopy was considered the gold standard, and therefore, a distinct possibility exists that there was a degree of reporting bias with regards to the miss rates of fluoroscopy in particular. Limitations included the large variation in techniques employed during MRD along with numerous fluoroscopic techniques that were utilized across the different studies. Based on their analysis, the authors concluded that MRD has a role in the assessment of pelvic floor dysfunction. However, they advise that clinicians need to be mindful of the risk of under-diagnosis and consideration of the use of additional imaging.

Cappabianca et al. (2011) compared the diagnostic efficacy of dynamic MR defecography (MR-D) with entero-colpocysto-defecography (ECCD) in the assessment of midline pelvic floor hernias (MPH) in female pelvic floor disorders. 1,142 participants underwent MR-D with analysis. The results of the study indicated that MR-D shows lower sensitivity than ECCD in the detection of MPH development.

Clinical Practice Guidelines

American Gastroenterological Association (AGA)

The AGA guideline on constipation states that although ARM and a rectal BET generally suffice to diagnose or exclude a defecatory disorder, defecography, which is generally performed with barium, or at some centers with magnetic resonance imaging, is useful if results are inconclusive (Bharucha et al. 2013a).

American College of Gastroenterology (ACG)

The ACG clinical guideline for management of benign anorectal disorders notes that barium or MRD can identify structural causes of outlet obstruction if one is expected. They may also confirm or exclude the diagnosis of defecatory disorders (DD) when the clinical features suggest DD, but the results of ARM and BET are equivocal (moderate recommendation, moderate quality of evidence) (Wald et al., 2014, updated 2021).

The same ACG 2014 guideline also cites the advantages of MRI over defecography as being better resolution of soft tissue surrounding the rectum and anal canal, including the bladder, uterus, and small intestine during dynamic imaging; improved ability to visualize anal sphincter and levator ani muscles with endoanal MRI, and lack of radiation (Wald et al., 2014, updated 2021).

American Society of Colon and Rectal Surgeons (ASCRS)

The ASCRS clinical practice guideline for the evaluation and management of chronic constipation states individuals who exhibit no improvement with dietary changes, fiber therapy and osmotic laxatives should be evaluated for outlet obstruction. Anorectal testing or dynamic imaging by fluoroscopic defecography, MRI defecography may assist in identifying functional or structural causes related to an evacuation disorder. Although MRI, performed in the supine position, permits excellent assessment of all pelvic floor compartments and the surrounding musculature, fluoroscopic defecography performed in the seated position is considered the evacuation examination with the most construct validity (Alavi et al., 2024). (Grade of Recommendation: conditional recommendation based on low-quality evidence).

In an updated clinical practice guideline on the treatment of rectal prolapse, the ASCRS (Bordeianou et al., 2017) states that if prolapse is suggested but cannot be seen during physical examination, fluoroscopic defecography, MRI defecography, or BET may reveal the problem. Defecography may also reveal associated anterior pelvic floor support defects, such as cystocele, vaginal vault prolapse, and enterocele (Grade of Recommendation: strong recommendation based on moderate-quality evidence, 1B).

Electrogastrography (EGG)/Electroenterography/Body Surface Gastric Mapping (BSGM)

Despite a possible use in clinical research, the studies of electrogastrography and body surface gastric mapping fail to provide convincing evidence that this technique is accurate for diagnosis of gastric disorders such as gastric stasis in clinical practice or that it has a positive impact on patient management or disease outcome. Additional studies are needed to determine if EGG is a useful adjunctive test or alternative to radioscintigraphy for the diagnosis of gastric stasis. These studies should involve a standardized procedure for diagnosis of gastroparesis with electrogastrography including recording, analysis, and interpretation. No studies were found to indicate electroenterography has a positive impact on patient management or disease outcome.

ECRI (2024) published a Clinical Evidence Assessment on the Gastric Alimetry System and concluded that the published evidence available for review was very low quality. The report assessed one diagnostic cohort study that reported measures of diagnostic accuracy and one retrospective case series that reported test impact on clinical decision making. ECRI did not find any direct clinical utility studies that reported improvements in patient-relevant outcomes following Alimetry-guided clinical decision making. The report concluded that additional clinical validity studies using appropriate reference standards and larger sample sizes are needed to assess Alimetry's diagnostic accuracy and that clinical utility studies are also needed to determine whether the use of this device to guide treatment and clinical decision making will result in improvements in patient-relevant outcomes.

A systematic review and meta-analysis by Peralta-Palmezano et al (2024) examined the prevalence and range of abnormalities in gastric slow waves in adults with gastroparesis who underwent EGG. The systematic review included 31 prospective studies with 1545 patients with gastroparesis and 340 controls (reported in 14 studies). In the 23 studies that reported the sex of the patients, 71.1% were women. The majority of the studies (67.7%) were conducted in the United States, four were conducted in Germany, and one each in Taiwan, Turkey, the Netherlands, France, Israel, and Germany. The authors reported that patients with gastroparesis had less normogastria (fasting: 50.3% versus 65.8%) (post-stimulus: 54.3% versus 66.5%), more bradygastria (fasting: 37.7% versus 13%) (post-stimulus: 31.9% versus 16.3%), and more

tachygastric (fasting: 16.1% versus 4.6%) (post-stimulus: 18.3% versus 5.2%). Limitations of this systematic review include the heterogeneity of the included studies' test protocols, concomitant medications used, and total duration of recordings during the EGG procedures. Additionally, the study did not address the clinical utility of the test and whether its use improves patients' outcomes. The authors concluded that adults with gastroparesis had a significantly lower percentage of normogastric than the controls, while they also had a higher percentage of bradygastric and tachygastric. This systematic review and meta-analysis included the Al Kafee et al. (2022) and Gharibans et al. (2019) studies previously summarized in this policy.

Xu et al. (2024) conducted a test-validation cohort study to investigate gastric myoelectrical abnormalities and symptoms in patients after fundoplication using noninvasive BSGM. The study included 16 adults (median age 34.5 years; 37.5% female) who had undergone a previous fundoplication operation (median time since fundoplication was five years) and who had ongoing significant gastroduodenal symptoms and 16 adult matched controls (based on age, biological sex, and BMI). BSGM using the Gastric Alimetry device was performed on each study participant. The authors reported that six of the 16 participants (37.5%) who had previously undergone a fundoplication operation showed significant spectral abnormalities defined by unstable gastric myoelectrical activity (n = 2), abnormally high gastric frequencies (n = 3), or high gastric amplitudes (n = 1) and that these participants had higher Patient Assessment of Upper Gastrointestinal Disorders-Symptom Severity Index scores than the remaining 10 patients with normal BSGM spectrograms. According to the authors, two of the three patients with abnormally high gastric frequencies had presumed vagal nerve injury documented in their procedure notes. The authors also reported that seven of 16 participants had BSGM test results suggestive of gut-brain axis contributions and without myoelectrical dysfunction. The authors concluded that a significant number of patients with persistent post-fundoplication symptoms displayed abnormal gastric functioning on BSGM testing, which correlated with symptom severity. Limitations of the study include the small cohort size, the inclusion of patients with a single procedure and with multiple revisional procedures, the lack of blinding, and the consecutive sample patient selection methodology. Also, the study does not address the clinical utility of the test and whether its use improves outcomes of patients after fundoplication.

Patient-specific phenotyping using BSGM was compared to gastric emptying testing (GET) in an exploratory comparison study by Wang, et al. (2024) that included 75 adults (77% female, median age 43 years) with chronic gastroduodenal symptoms. All patients had undergone a clinical work-up by a gastroenterologist, including upper gastrointestinal endoscopy, to exclude alternative pathologies, had withheld any medications affecting gastrointestinal motility for 48 hours, had completed an overnight fast, and were asked to avoid caffeine, nicotine, opiates and cannabis the morning of to their testing. Each study participant underwent simultaneous GET and BSGM that consisted of a 30-minute baseline reading, consumption of a 99mTC-labelled egg meal, and 4-four-hour postprandial recording. Before motility testing was done, 56 patients met Rome IV Criteria for chronic nausea and vomiting syndromes (CNVS) [75%, with 52/56 also meeting functional dyspepsia (FD) criteria]. There were 14 participants who met FD criteria alone, and five did not meet either criteria, indicating a high chronic gastroduodenal symptom burden in the study cohort. The authors reported that motility abnormality detection rates were 22.7% for GET with 14 delayed and three rapid while BSGM spectral analysis was 33.3% with 14 low rhythm stability/low amplitude, five high amplitude, and six abnormal frequency. The authors also reported that, in patients with normal spectral analysis, BSGM symptom phenotypes included sensorimotor 17% (where symptoms strongly paired with gastric amplitude), continuous 30%, and other 53%. The authors reported that BSGM phenotypes showed superior correlations with the Gastroparesis Cardinal Symptom Index (GCSI), the Patient Assessment of Upper Gastrointestinal Symptom Severity Index, and anxiety scales while the Rome IV Criteria did not correlate with psychometric scores. The authors concluded that BSGM improves patient phenotyping in chronic gastroduodenal disorders in the presence and absence of motility abnormalities with increased correlation with symptoms and psychometrics compared with GET and Rome IV criteria. Whether use of BSGM improves patients' outcomes is however not addressed in this study.

Xu et al. (2023) performed a study using BSGM in people with longstanding type 1 diabetes (T1D) with and without symptoms to define phenotypes of gastric myoelectrical abnormalities. The study included 64 people, 32 of which had a medical history of T1D of more than 10 years (mean age of 50 years; 64% female), and 32 were controls. Of the 32 patients with T1D, 15 were noted to have a high symptom burden (12 with CNVS and FD, and three with FD only) based on their assessment using the Rome IV criteria for CNVS if they met at least one of the criteria. The 32 participants with T1D were then matched to a database of controls in a 1:1 ratio using the nearest neighbor based on age, sex, and body mass index. The authors reported that the patients with T1D with symptoms showed more unstable gastric myoelectrical activity and lower average special covariance compared to controls and that symptomatic patients also had a higher prevalence of peripheral neuropathy, anxiety/depression diagnoses, and higher mean hemoglobin A1C levels. They also reported that deviation in gastric frequency was positively correlated with symptoms of bloating, upper gut pain, nausea and vomiting and fullness. Limitations of the study included the lack of blood glucose monitoring for all the participants (due to device availability), different meals given to the study participants with T1D than the control group received, the ongoing development of reference values for the emerging spatial metrics and the small sample size. The authors

concluded that gastric symptoms in people with longstanding T1D correlated with myoelectrical abnormalities on BSGM testing, in addition to glycemic control, psychological comorbidities and peripheral neuropathy.

Schamberg et al. (2023) conducted a multi-center, retrospective observational study to compare BSGM and EGG to quantify performance differences. The study included EGG and BSGM data from 178 subjects [43 patients with nausea and vomiting (NVS), 32 patients with type 1 diabetes (T1D) and 110 healthy volunteers]. The study assessed the use of BSGM and EGG in the following three domains: group level-differences in measures of gastric activity, the relationship between gastric abnormalities and symptoms, and patient-level classifications of gastric health. The comparisons followed standard methodologies for each test including pre-processing, post-processing, and analysis. Statistical evaluations were done for group-level differences, symptom correlations and patient-level classifications. The authors reported that BSGM showed substantially tighter frequency ranges when compared to EGG in the control group and that both tests detected rhythm instability in NVS, but EGG showed opposite frequency effects in T1D. The authors also reported that BSGM showed an eight times increase in the number of significant correlations with symptoms and that BSGM accuracy for patient-level classification was 0.78 for patients when compared to controls and 0.96 when compared to a blinded consensus panel while EGG accuracy was 0.54 and 0.43. Limitations of the study include the automated EGG analysis methodology, the possibility that proprietary signal processing steps may exist but were not used, the lack of testing with other meal preparations, only using a single electrode configuration for EGG testing, the heterogeneity of EGG processing approaches, and the focus on only spectral analyses of BSGM and EGG. The authors concluded that EGG detected group-level differences in patients but lacked symptom correlations and showed poor accuracy for patient-level classification while BSGM demonstrated substantial performance improvements across all three domains.

In a study to evaluate CNVS pathologies, Gharibans et al. (2022) performed BSGM in 43 patients with NVS and 43 matched controls. The study participants were adults, primarily female (76.7%) with a median age of 33 years (range 26 to 44). Each participant underwent BSGM that entailed a fasting baseline, ingestion of a 482-kilocalorie meal and a four-hour postprandial recording, then spectral and spatial biomarker analyses. The authors reported that meal responses were impaired in NVS with multiple BSGM abnormalities compared to the study controls, impaired fed-fasting power ratios, and disorganized slow waves. The authors also reported that most patients (62%) had normal BSGM results with increased psychological comorbidities and anxiety scores while a smaller subgroup (31%) had markedly abnormal BSGM with biomarkers that correlated with symptoms and that patients with NVS shared overlapping symptoms but comprised distinct underlying phenotypes that correlated with symptoms. Limitations of the study include the small sample size that may have affected the subanalyses performed, and the use of a consensus panel classification which may introduce subjectivity and limit reproducibility. Other limitations of the study included the lack of a control group, the under-representation of patients with diabetes as only 7% of the participants had diabetes, and the inclusion of patients with BMI > 35 as a high BMI may result in overestimating the low rhythm stability phenotype due to declining signal-to-noise ratio. The authors concluded that the study showed that BSGM expanded the phenotyping of patients with chronic gastroduodenal disorders when compared to GET and that the results of BSGM could improve clinical management of these patients by separating those with gastric dysfunction from those with gut-brain dysregulation or other etiologies.

A systematic review and meta-analysis was completed by Bhat et al. (2021) involving electrogastronomy (EGG) use in adults with gastroesophageal reflux disease (GERD). After the published literature was reviewed, thirteen studies were included in the analysis with a total of 591 participants (427 with GERD; 164 healthy controls) who had completed an EGG procedure. The study found that patients with GERD spent significantly less time with normal gastric slow-wave activity compared to healthy controls. The authors noted that correlations between GERD symptoms and EGG recordings were inconsistently studied; EGG apparatus and techniques also varied across the studies. They also recognized the limitations of the studies available including the known limitations to low-resolution EGG methodologies (as high-resolution EGG is now available), and the inclusion of studies that relied on subjective symptom-based diagnostic criteria. They concluded that further investigation for the use of EGG in adults with GERD is warranted.

In an evaluation of 54 patients with FD, Russo et al. (2017) utilized the results of EGG to differentiate postprandial distress syndrome (PDS) with epigastric pain syndrome (EPS). Using a symptom questionnaire, 42 patients were classified as PDS and 12 as EPS, although an overlap between the symptom profiles of the two subgroups was recorded. The EGG parameters (the postprandial instability coefficient of dominant frequency, the dominant power, and the power ratio) were significantly different between the subgroups, whereas the gastric emptying time did not differ significantly. In addition, EPS was characterized by a different gut peptide profile compared with PDS. Finally, neurotensin polymorphism was shown to be associated with neurotensin levels. The authors concluded that this evidence deserves further studies into FD. This study, however, does not support the use of EGG in clinical practice or beyond its use for research.

Kayar et al. (2016) utilized transcutaneous EGG to compare patients with FD (n = 30) to control subjects (n = 30) in terms of motility abnormalities according to the EGG results. A high incidence of gastric motility and myoelectrical activity abnormalities was observed in patients with FD. The authors concluded that although still considered an experimental

method, EGG is an effective, dependable, and non-invasive method in differentiating the subgroups and may be an essential and irreplaceable test to diagnose and follow-up patients with FD with motor dysfunction.

Clinical Practice Guidelines

American Gastroenterological Association (AGA)

A position statement from the AGA (Parkman, et al. 2004) on the diagnosis and treatment of refractory gastroparesis does not recommend the use of EGG. In their 2022 Clinical Practice update (Lacy, et al), the AGA guideline does not focus on the etiology, pathophysiology or diagnostic testing for refractory gastroparesis and there is no longer any mention or direction for the use of electrogastrography for the diagnosis of gastroparesis.

Esophageal Mucosal Integrity Testing

The limited number of published clinical studies on the use of esophageal mucosal integrity testing by electrical impedance fail to provide convincing evidence that this technology is safe and effective for diagnosing GERD, eosinophilic esophagitis (EoE), or nonacid reflux disease (non-GERD), or for the monitoring of treatment response in GERD and EoE. Additional studies are needed to provide evidence of the efficacy of this technology.

Hayes (2022, updated 2024) published an Evolving Evidence Review on the safety and efficacy of MiVu Mucosal Integrity Testing (Diversatek, Inc.) for the diagnosis of GERD in adults. Following their review of three “very poor-quality” studies and one systematic review that compared several emerging diagnostic tests, Hayes stated that the system may distinguish previously diagnosed EoE from GERD at least as well as esophagogastroduodenoscopy (EGD) and potentially better than pH testing but that there was a lack of data from participants that did not yet have a definitive diagnosis. Hayes also stated that insufficient data was published addressing the clinical utility of the device in terms of impact on clinical management or outcomes to draw conclusions regarding level of support for that use. In their review of medical society guidelines, Hayes assigned a weak level of support to reflect that the guidelines they identified indicated there is “promise” in this device as an adjunctive diagnostic tool but that the societies did not recommend it as preferable to standard diagnostic methods. Hayes’ assessment recommended that the literature continue to be monitored, as MI testing technology is still developing. In their 2024 update, Hayes identified one relevant newly published single-center, case-controlled study since this report was published in 2022; however, Hayes stated that this study would not result in a change to their minimal level of support for this device as there was no new evidence regarding the safety of the device or regarding longer-term follow-up. The Choksi et al. (2018) and Patel et al. (2019) studies summarized below are included in this report.

ECRI (2021) published an Evidence Analysis on the efficacy of the MiVu Mucosal Integrity Testing System (Diversatek, Inc.) for help diagnosing GERD and its ability to obtain real-time measurements of esophageal epithelial impedance during an endoscopy. The device is not for use as a sole diagnostic screening tool. ECRI’s review identified three studies for inclusion. Based on their review of the abstracts, ECRI concluded that the available published evidence was limited and that additional clinical validity studies with larger sample sizes are needed to assess the diagnostic accuracy of this device. ECRI also stated that clinical utility studies are needed to determine whether the use of this device to guide treatment and clinical decision making will result in improvements in patient-relevant outcomes. The ECRI review included the Patel et al. (2019) and Choksi et al. (2018) studies summarized below.

Patel et al. (2019) conducted a prospective, multi-center test-validation study to evaluate the ability of a balloon mucosal impedance (MI) catheter system to detect and evaluate esophageal disorders such as GERD and EoE. The study included 69 adult participants (91.7% Caucasian) who underwent esophagogastroduodenoscopy (EGD) then balloon MI testing during endoscopy and prior to wireless pH monitoring or esophageal biopsies for suspected GERD. The participants were classified based on endoscopic, pH monitoring and pathology findings as having GERD (n = 24; median age 48; 54% female), EoE (n = 21; median age 33; 33% female), or non-GERD (n = 24; median age 62; 71% female). The authors reported that the MI pattern along the esophageal axis differed significantly among participants with GERD, EoE and non-GERD as the MI pattern for GERD was easily distinguished from that of EoE with low MI values in the distal esophagus and normalized values along the proximal esophagus in those individuals with GERD while the measurements were low in all segments of the esophagus in participants with EoE. The authors also reported that those with non-GERD had higher baseline MI values in the distal esophagus and MI that the values remained elevated along the esophagus. One adverse event was reported by the authors in a participant with EoE who showed a small distal esophageal mucosal tear similar to that seen during endoscopic dilation. The authors concluded that the balloon MI catheter system instantly detected changes in esophageal mucosal integrity during endoscopy which could potentially obviate the need for 24-to-48-hour ambulatory wireless pH monitoring or esophageal biopsies for histopathology. The authors also concluded that the MI was safe and effective in identifying patients with GERD, EoE, or non-GERD. Limitations of the study include the small sample size, the lack of blinding, the homogeneity of the study population, and the assumption that the participants must belong to one of the three diagnosis groups with an equal baseline prevalence of GERD, non-GERD and EoE.

Furthermore, the findings of this study need to be validated in an independent patient population and the clinical utility of the test in improving patients' outcome is not addressed.

Choksi et al. (2018) conducted a retrospective analysis of 91 adult individuals (80.32% Caucasian) with upper gastrointestinal symptoms (non-GERD, n = 30) who were referred for GERD (n = 38) and EoE (n = 23) diagnostic testing to quantify MI testing that measured epithelial integrity during EGD and to identify patterns that differentiated individuals with and without GERD from those with EoE. Additionally, the authors sought to determine whether MI values and patterns were sufficient in identifying individuals with EoE utilizing histologic findings as a reference. The individuals underwent initial endoscopy with MI measurements obtained. The authors utilized statistical modeling to identify MI patterns within the esophagus that were associated with GERD versus EoE. The authors reported there were no statistically significant differences within the groups regarding race. MI parameters were determined for distinguishing EoE from non-EoE conditions (GERD or normal). The authors reported that the data showed significant improvement in sensitivity (100% vs 86%) and specificity (96% vs 59%) when predicting a diagnosis of EoE using MI pattern. Findings were validated by the authors in a prospective cohort of 49 individuals, symptomatic for dysphagia with no prior diagnosis, who underwent EGD for dysphagia, to test the ability of MI patterns to identify individuals with versus without EoE. The authors reported individuals with EoE had a unique MI pattern and low values long with esophageal axis and that MI measurements differentiated the individual populations. The authors reported per the validation cohort, that the assessment of mucosal integrity by MI values and pattern alone without endoscopic or clinical presentation was sufficient with a high level of accuracy in providing the correct diagnosis. Limitations of the study include the use of only two impedance rings, forming a single MI sensing channel and that the catheter was manually repositioned to various sites of the esophagus which may have resulted in some variability. Additional limitations include the small sample size as well as the homogeneity of the study population.

Clinical Practice Guidelines

American College of Gastroenterology (ACG)

The ACG's Clinical Guideline addressing the diagnosis and management of GERD states that ACG expects new diagnostic tools and treatments to be developed and refined. The statement goes on to state that mucosal integrity testing is available commercially but that it is not developed sufficiently to warrant discussion in this guideline (Katz, et al. 2022).

U.S. Food and Drug Administration (FDA)

This section is to be used for informational purposes only. FDA approval alone is not a basis for coverage.

Instruments to perform cutaneous electrogastrography, electroenterography, and body surface gastric mapping are regulated by the FDA as Class II devices. Refer to the following website for more information (use product code MYE or FFX): <http://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfPMN/pmnmn.cfm>. (Accessed July 22, 2025)

Several radiopaque markers have been approved by the FDA for colonic transit testing. Refer to the following website for more information (use product code FFX): <http://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfPMN/pmnmn.cfm>. (Accessed July 22, 2025)

Defecography is a procedure and, therefore, is not subject to FDA regulation. However, any medical equipment, drugs or tests used as part of this procedure may be subject to FDA regulation. A general list of cleared magnetic resonance imaging systems for MRI defecography can be found by entering the code LNH into the "product code" window in the form at the following FDA 510(k) database website: <http://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfPMN/pmnmn.cfm>. (Accessed July 22, 2025)

The MiVu™ Mucosal Integrity Testing System (MiVu) (Diversatek Healthcare, Highlands Ranch, CO) received FDA clearance as a Class II de novo device on December 23, 2019 as a new approach to assessing esophageal mucosal integrity. FDA 510(k) Premarket Notification was received on April 25, 2023 under 510(k) Number K230056. Refer to the following website and search using either the product name or the Product Code of QIS for more information: <https://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfPMN/pmnmn.cfm>. (Accessed July 22, 2025)

The EndoFLIP® System is indicated for use in a clinical setting to measure pressure and dimensions in the esophagus, pylorus, and anal sphincters and received 510K approval on April 17, 2017. It is intended to be used as an adjunct procedure to other diagnostic methods as part of a comprehensive evaluation of patients with symptoms consistent with gastrointestinal motility disorders. Refer to the following website for additional information: <https://www.accessdata.fda.gov/scripts/cdrh/cfdocs/cfpmn/pmnmn.cfm?ID=K170833>. (Accessed July 22, 2025)

References

- Al Kafee A, Cilaci T, Kayar Y, et al. Electrogastrography in patients with functional dyspepsia, joint hypermobility, and diabetic gastroparesis. *Turk J Gastroenterol*. 2022 Feb 4.
- Alabbas M, Khoudari H, Ghosh G, et al. FLIP use in achalasia: comparing POEM and Heller myotomy outcomes: a systematic review and meta-analysis. *Surg Endosc*. 2025 Jul;39(7):4060-4075. doi: 10.1007/s00464-025-11776-4.
- Alavi K, Thorsen AJ, Fang SH, et al; Clinical Practice Guidelines Committee of the American Society of Colon and Rectal Surgeons. The American Society of Colon and Rectal Surgeons clinical practice guidelines for the evaluation and management of chronic constipation. *Dis Colon Rectum*. 2024 Oct 1;67(10):1244-1257.
- American Society for Gastrointestinal Endoscopy (ASGE). The role of endoscopy in the management of GERD. *Gastroenterology* 2015 June; 81(6):1305-1310.
- Benitez AJ, Budhu S, Burger C, et al. Use of the functional luminal imaging probe in pediatrics: A comparison study of patients with achalasia before and after endoscopic dilation and non-achalasia controls. *Neurogastroenterol Motil*. 2021 Dec;33(12):e14133.
- Bharucha AE, Dorn SD, Lembo A, et al. American Gastroenterological Association medical position statement on constipation. *Gastroenterology*. 2013a Jan;144(1):211-17.
- Bhat S, Varghese C, Carson DA, et al. Gastric dysrhythmia in gastroesophageal reflux disease: a systematic review and meta-analysis. *Esophagus*. 2021 Jul;18(3):425-435.
- Bordeianou L, Paquette I, Johnson E, et al. American Society of Colon and Rectal Surgeons. Clinical practice guidelines for the treatment of rectal prolapse. *Dis Colon Rectum*. 2017 Nov;60(11):1121-1131.
- Camilleri M, Ford AC, Mawe GM, et al. Chronic constipation. *Nat Rev Dis Primers*. 2017 Dec 14; 3:17095.
- Cappabianca S, Reginelli A, Lacobellis F, et al. Dynamic MRI defecography vs. entero-colpo-cysto-defecography in the evaluation of midline pelvic floor hernias in female pelvic floor disorders. *Int J Colorectal Dis*. 2011 Sep;26(9):1191-6.
- Choksi Y, Lal P, Slaughter JC, et al. Esophageal mucosal impedance patterns discriminate patients with eosinophilic esophagitis from patients with GERD. *Clin Gastroenterol Hepatol*. 2018 May;16(5):664-671.e1.
- Chu H, Lin Z, Zhong L, et al. Treatment of high-frequency gastric electrical stimulation for gastroparesis. *J Gastroenterol Hepatol*. 2012 Jun;27(6):1017-26.
- Desprez C, Roman S, Leroi AM, et al. The use of impedance planimetry (Endoscopic Functional Lumen Imaging Probe, EndoFLIP®) in the gastrointestinal tract: A systematic review. *Neurogastroenterol Motil*. 2020 Sep;32(9):e13980.
- ECRI. EndoFLIP Impedance Planimetry System (Medtronic plc.) for identifying gastrointestinal motility disorders. Plymouth Meeting PA. ECRI; 2018 October; Revised 2024 September. (Executive Summary).
- ECRI. Gastric Alimetry System (Alimetry Ltd.) for aiding diagnosis of gastric disorders. Clinical Evidence Assessment. 2024 Jan.
- ECRI. Gastric Alimetry System (Alimetry Ltd.) for aiding diagnosis of gastric disorders. Clinical Evidence Assessment. 2024 Jan.
- ECRI. MiVu Mucosal Integrity Testing System (Diversatek, Inc.) for diagnosing gastroesophageal reflux disease. Curated Literature Search. 2021 Jul.
- Fehervari M, Middleton I, McGlone, ER, et al. Endoscopic functional luminal imaging probe for intraprocedural guidance of achalasia treatment: a systematic review and meta-analysis. *International Journal of Surgery*. 2025 July; 111(7):4165-4176.
- Foti PV, Farina R, Riva G, et al. Pelvic floor imaging: comparison between magnetic resonance imaging and conventional defecography in studying outlet obstruction syndrome. *Radiol Med*. 2013 Feb;118(1):23-39.
- Gharibans AA, Calder S, Varghese C, et al. Gastric dysfunction in patients with chronic nausea and vomiting syndromes defined by a noninvasive gastric mapping device. *Sci Transl Med*. 2022 Sep 21;14(663):eabq3544.
- Gharibans AA, Coleman TP, Mousa H, et al. Spatial patterns from high-resolution electrogastrography correlate with severity of symptoms in patients with functional dyspepsia and gastroparesis. *Clin Gastroenterol Hepatol*. 2019 Dec;17(13):2668-2677.
- Hayes, Inc. Evidence Analysis Research Brief. Endolumenal functional lumen imaging probe (EndoFLIP; Medtronic Inc.) for evaluation of esophageal and gastric disorders. Hayes, Inc.; Lansdale, PA: December 2024.

Hayes, Inc. Evolving Evidence Review. MiVu mucosal integrity testing (Diversatek Inc.) for diagnosis of gastroesophageal reflux disease in adults. Hayes, Inc.; June 2022. Updated July 10, 2024.

Hayes, Inc. Health Technology Assessment. Endolumenal functional lumen image probe (EndoFLIP; Medtronic Inc.) for evaluation of esophageal and gastric disorders. Hayes, Inc; Lansdale, PA: June 2019; Updated June 2022.

Hirano I, Pandolfino JE, Boeckxstaens GE. AGA clinical practice Update on functional lumen imaging probe for the management of esophageal disorders: expert review from the clinical practice updates committee of the AGA Institute. *Clinical Gastroenterology and Hepatology*. 2017 March;15(3):325-334.

Katz PO, Dunbar KB, Schnoll-Sussman FH, et al. ACG clinical guideline for the diagnosis and management of gastroesophageal reflux disease. *Am J Gastroenterol*. 2022 Jan 1;117(1):27-56.

Kayar Y, Danalıoğlu A, Kafee AA, et al. Gastric myoelectrical activity abnormalities of electrogastrography in patients with functional dyspepsia. *Turk J Gastroenterol*. 2016 Sep;27(5):415-420.

Keller J, Bassotti G, Clarke J, et al. Expert consensus document: Advances in the diagnosis and classification of gastric and intestinal motility disorders. *Nat Rev Gastroenterol Hepatol*. 2018 May;15(5):291-308.

Pääkkö E, Mäkelä-Kaikkonen J, Laukkanen H, et al. X-ray video defaecography is superior to magnetic resonance defaecography in the imaging of defaecation disorders. *Colorectal Dis*. 2022 Jun;24(6):747-753.

Parkman HP, Hasler WL, Fisher RS; American Gastroenterological Association. American Gastroenterological Association medical position statement: diagnosis and treatment of gastroparesis. *Gastroenterology*. 2004 Nov;127(5):1589-91.

Pasricha PJ, Camilleri M, Hasler WL, et al. White paper AGA: gastroparesis: clinical and regulatory insights for clinical trials. *Clin Gastroenterol Hepatol*. 2017 Aug;15(8):1184-1190.

Patcharatrakul T, Rao SSC. Update on the pathophysiology and management of anorectal disorders. *Gut Liver*. 2018 Jul 15;12(4):375-384.

Patel DA, Higginbotham T, Slaughter JC, et al. Development and validation of a mucosal impedance contour analysis system to distinguish esophageal disorders. *Gastroenterology*. 2019 May;156(6):1617-1626.e1.

Peralta-Palmezano JJ, Escobar-Serna DP, Peralta-Palmezano FJ, et al. Electrogastrography in adult gastroparesis: A systematic review and meta-analysis. *Dig Dis Sci*. 2025 Jan;70(1):298-315.

Poncelet E, Rock A, Quinton J, et al. Dynamic MR defecography of the posterior compartment: Comparison with conventional X-ray defecography. *Diagn Interv Imaging*. 2017 Apr;98(4):327-332.

Rafeeqi T, Salimi-Jazi F, Cunningham A, et al. The utility of Endoscopic Functional Luminal Imaging (EndoFLIP) in the diagnosis and management of children with achalasia. *J Pediatr Surg*. 2023 Apr;58(4):639-642.

Ramage L, Georgiou P, Qiu S, et al. Can we correlate pelvic floor dysfunction severity on MR defecography with patient-reported symptom severity? *Updates Surg*. 2018 Dec;70(4):467-476.

Ramage L, Simillis C, Yen C, et al. Magnetic resonance defecography versus clinical examination and fluoroscopy: a systematic review and meta-analysis. *Tech Coloproctol*. 2017 Dec;21(12):915-927.

Rao SS, Patcharatrakul T. Diagnosis and treatment of dyssynergic defecation. *J Neurogastroenterol Motil*. 2016 Jul 30;22(3):423-35.

Russo F, Chimienti G, Clemente C, et al. Gastric activity and gut peptides in patients with functional dyspepsia: postprandial distress syndrome versus epigastric pain syndrome. *J Clin Gastroenterol*. 2017 Feb;51(2):136-144.

Schamberg G, Calder S, Varghese C, et al. Comparison of Gastric Alimetry® body surface gastric mapping versus electrogastrography spectral analysis. *Sci Rep*. 2023 Sep 11;13(1):14987.

Shada A, Nielsen A, Marowski S, et al. Wisconsin's Enterra Therapy Experience: A multi-institutional review of gastric electrical stimulation for medically refractory gastroparesis. *Surgery*. 2018 Oct;164(4):760-765.

Thanaracthanon P, Sasiwimonphan K, Sunthornram A, et al. Diagnostic performance of dynamic MR defecography in assessment of dyssynergic defecation. *Abdom Radiol (NY)*. 2023 Nov;48(11):3458-3468.

Vaezi, MF, Pandolfino, JE, Yadlapati, RH, et al. ACG Clinical Guidelines: Diagnosis and management of achalasia. *The American Journal of Gastroenterology*. 2020 September; 115(9):1393-1411.

van Gruting IM, Stankiewicz A, Thakar R, et al. Imaging modalities for the detection of posterior pelvic floor disorders in women with obstructed defaecation syndrome. *Cochrane Database Syst Rev*. 2021 Sep 23;9(9):CD011482.

van Iersel JJ, Formijne Jonkers HA, Verheijen PM, et al. Comparison of dynamic magnetic resonance defaecography with rectal contrast and conventional defaecography for posterior pelvic floor compartment prolapse. *Colorectal Dis*. 2017 Jan;19(1):O46-O53.

Vitton V, Vignally P, Barthet M, et al. Dynamic anal endosonography and MRI defecography in diagnosis of pelvic floor disorders: comparison with conventional defecography. *Dis Colon Rectum*. 2011 Nov;54(11):1398-404.

Wald A, Bharucha AE, Limketkai B, et al. ACG clinical guidelines: management of benign anorectal disorders. *Am J Gastroenterol*. 2021 Oct 1;116(10):1987-2008.

Wang WJ, Foong D, Calder S, et al. Gastric Alimetry® expands patient phenotyping in gastroduodenal disorders compared to gastric emptying scintigraphy. *Am J Gastroenterol*. 2024 Feb 1;119(2):331-341.

Xu W, Gharibans AA, Calder S, et al. Defining and phenotyping gastric abnormalities in long-term type 1 diabetes using body surface gastric mapping. *bioRxiv*. 2023 Sep 21;2(8):1120-1132.

Xu W, Wang T, Foong D, et al. Characterization of gastric dysfunction after fundoplication using body surface gastric mapping. *J Gastrointest Surg [Internet]*. 2024 Mar 1;28(3):236-45.

Yin J, Chen JDZ. Electrogastrography: methodology, validation and applications. *J Neurogastroenterol Motil*. 2013 Jan;19(1):5-17.

Yoo IK, Choi SA, Kim WH, et al. Assessment of clinical outcomes after peroral endoscopic myotomy via esophageal distensibility measurements with the endoluminal functional lumen imaging probe. *Gut Liver*. 2019 Jan 15;13(1):32-39.

Zafar A, Seretis C, Feretis M, et al. Comparative study of magnetic resonance defaecography and evacuation proctography in the evaluation of obstructed defaecation. *Colorectal Dis*. 2017 Jun;19(6):0204-0209.

Policy History/Revision Information

Date	Summary of Changes
04/01/2026	<p>Title Change</p> <ul style="list-style-type: none"> Previously titled <i>Gastrointestinal Motility Disorders, Diagnosis and Treatment (for New Mexico Only)</i> <p>Coverage Rationale</p> <ul style="list-style-type: none"> Removed and relocated language pertaining to gastric electrical stimulation (GES) therapy; refer to the Medical Policy titled <i>Minimally Invasive Procedures for the Treatment of Upper Gastrointestinal Diseases (for New Mexico Only)</i> Revised list of unproven and not medically necessary procedures: <ul style="list-style-type: none"> Added “Functional Lumen Imaging Probe (FLIP) technology for diagnosing Achalasia” Removed list of examples of: <ul style="list-style-type: none"> Cutaneous, mucous, or serosal Electrogastrography, electroenterography, or body surface gastric mapping: Gastric Alimetry System, G-Tech Gut Tracker wireless patch system Esophageal Mucosal Integrity Testing by electrical impedance: MiVu™ Mucosal Integrity Testing System <p>Definitions</p> <ul style="list-style-type: none"> Added definition of: <ul style="list-style-type: none"> Achalasia Functional Lumen Imaging Probe (FLIP) <p>Applicable Codes</p> <ul style="list-style-type: none"> Removed CPT codes 43647, 43648, 43881, 43882, 64590, and 64595 <p>Supporting Information</p> <ul style="list-style-type: none"> Updated <i>Description of Services, Clinical Evidence, FDA, and References</i> sections to reflect the most current information Archived previous policy version CS046NM.C

Instructions for Use

This Medical Policy provides assistance in interpreting UnitedHealthcare standard benefit plans. When deciding coverage, the federal, state or contractual requirements for benefit plan coverage must be referenced as the terms of the federal, state or contractual requirements for benefit plan coverage may differ from the standard benefit plan. In the event of a

conflict, the federal, state or contractual requirements for benefit plan coverage govern. Before using this policy, check the federal, state or contractual requirements for benefit plan coverage. UnitedHealthcare reserves the right to modify its Policies and Guidelines as necessary. This Medical Policy is provided for informational purposes. It does not constitute medical advice.

UnitedHealthcare may also use tools developed by third parties, such as the InterQual[®] criteria, to assist us in administering health benefits. The UnitedHealthcare Medical Policies are intended to be used in connection with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice.